



University of Tasmania

Role of amyloid precursor protein in neural stem/progenitor cell proliferation and differentiation

Yanling Hu (MSc)

**Submitted in fulfillment of the requirements for the
degree of Doctor of Philosophy**

**Menzies Research Institute Tasmania
University of Tasmania
08/2015**

DECLARATION OF ORIGINALITY

This thesis contains no material which has been accepted for a degree or diploma by the University or any other institution, except by way of background information and duly acknowledged in the thesis, and to the best of my knowledge and belief no material previously published or written by another person except where due acknowledgement is made in the text of the thesis, nor does the thesis contain any material that infringes copyright.

AUTHORITY OF ACCESS

This thesis may be made available for loan and limited copying and communication in accordance with the Copyright Act 1968.

ETHICS STATEMENT

The research associated with this thesis abides by the international and Australian codes on animal experimentation, and the rulings of the Animal Ethics Committee of the University of Tasmania.

Yanling Hu

Statement of Co-Authorship

Yanling Hu has incorporated a version of the published open access paper Hu et al into Chapter 3 of the thesis.

This work was co-authored with Dr. Amos Hung, Dr. Hao Cui, Dr. Edgar Dawkins, Dr. Marta Bólós, Dr. Lisa Foa, Dr. Kaylene Young, and Prof David Small. Some of the research, data analysis was contributed by Dr. Amos Hung and Dr. Marta Bólós. The majority of the research, analysis, figures were generated and prepared by Yanling Hu herself.

Hu, Y., Hung, A. C., Cui, H., Dawkins, E., Bólós, M., Foa, L., Young, K. M., and Small, D. H. (2013). Role of cystatin C in amyloid precursor protein induced proliferation of neural stem/progenitor cells. J Biol Chem 288, 18853-18862.

The contribution of authors to the publication is as follows:

Yanling Hu was the primary author, conducted data analysis, was involved in laboratory experiments and study design and manuscript preparation; Dr. Amos Hung designed and performed part of the neural stem progenitor cell assay; Dr. Hao Cui and Dr. Edgar Dawkins provided technical advice for immunocytochemistry and immunoblotting; Dr. Marta Bólós performed real time PCR; Dr. Lisa Foa contributed to manuscript preparation and study advice; Dr. Kaylene Young provided technical support and experimental design suggestions. Prof. David Small assisted with the scientific direction and provided input into research design. All authors contributed to the final manuscript preparation.

Signed:

Prof David Small
Supervisor
Menzies Institute for Medical Research
University of Tasmania

Prof Alison Venn
Director
Menzies Institute for Medical Research
University of Tasmania

Date : 7/4/2016

7/4/2016

Table of Contents

Abstract	IX
Acknowledgements	XIII
Publication and presentation record	XV
List of abbreviations	XVII

Chapter 1 Introduction	1
1.1 Alzheimer's disease (AD)	1
1.1.1 Clinical features of AD.....	1
1.1.2 Impact and prevalence of AD.....	3
1.1.3 Pathology of AD.....	4
1.1.3.1 Neurofibrillary tangles.....	5
1.1.3.2 Amyloid plaques.....	6
1.1.3.3 Cerebral amyloid angiopathy (CAA).....	9
1.1.3.4 Synapse loss.....	10
1.2 β - Amyloid ($A\beta$).....	10
1.2.1 Amyloid precursor protein processing and $A\beta$ production.....	11
1.2.2 Aggregation of $A\beta$	13
1.2.3 $A\beta$ toxicity.....	14
1.2.4 Possible physiological functions of $A\beta$	17
1.3 Biology of amyloid precursor protein.....	18
1.3.1 Structure of the APP family.....	18
1.3.2 APP trafficking and processing.....	21
1.3.3 APP secretases.....	23
1.3.3.1 α – Secretase.....	23
1.3.3.2 β – Secretase.....	24
1.3.3.3 γ – Secretase.....	26
1.3.4 Function of APP.....	28
1.3.4.1 Role of APP in cell adhesion and neurite outgrowth.....	29
1.3.4.2 Role of APP in synaptogenesis and synaptic plasticity.....	30
1.3.4.3 Non- neuronal functions of APP.....	33

1.3.4.4	APP signalling.....	34
1.3.4.4.1	APP may act as a cell -surface receptor.....	34
1.3.4.4.2	Role of APP in intracellular signalling.....	35
1.4	Risk factors for AD.....	38
1.4.1	Familial AD (FAD).....	39
1.4.1.1	APP mutations.....	39
1.4.1.2	Presenilin mutations.....	40
1.4.2	Late onset AD (sporadic AD).....	41
1.4.2.1	Apolipoprotein E (apoE)- the major risk factor.....	42
1.4.2.2	TREM2.....	45
1.4.2.3	CST3.....	48
1.5	APP transgenic and knockout mice.....	49
1.5.1	Transgenic mouse model of AD.....	49
1.5.2	APP knockout mice.....	51
1.6	Current and prospective therapeutic approach for AD.....	52
1.6.1	Current therapeutic treatment for AD.....	52
1.6.1.1	Acetylcholinesterase inhibitors.....	52
1.6.1.2	Memantine.....	53
1.6.2	Prospective therapeutic strategies to delay disease progression.....	54
1.6.2.1	A β - based approaches.....	54
1.6.2.1.1	A β immunotherapy.....	56
1.6.2.2	Stem cell therapy for AD.....	58
1.7	Biology of neural stem and progenitor cells (NSPCs).....	60
1.7.1	Embryonic stem cells (ESCs).....	60
1.7.2	Adult neural stem/progenitor cell (NSPCs) and neurogenesis.....	62
1.7.2.1	Adult neurogenesis in V-SVZ.....	64
1.7.2.2	Adult neurogenesis in subgranular zone of dentate gyrus (DG- SGZ).....	71
1.7.2.3	Function of adult neurogenesis.....	75
1.7.2.4	Factors influencing NSPC proliferation and differentiation.....	78
1.7.2.4.1	Growth factors.....	79
1.7.2.4.1.1	Epidermal growth factor (EGF).....	79
1.7.2.4.1.2	Fibroblast growth factor-2 (FGF-2).....	80
1.7.2.4.1.3	Insulin and insulin-like growth factors (IGFs).....	82

1.7.2.4.2	Other factors affecting NSPC proliferation and differentiation.....	84
1.7.2.4.2.1	Cystatin C.....	86
1.7.2.4.2.2	Neurogenin 2.....	89
1.7.2.4.3	An in - vitro model for studying NSPCs: neurospheres.....	90
1.8	Neurogenesis in neurodegenerative disease.....	92
1.8.1	Neurogenesis in Alzheimer's disease.....	93
1.8.1.1	Role of PS on adult neurogenesis.....	94
1.8.1.2	Role of tau in neurogenesis.....	96
1.9	Stem cell therapy for neurodegenerative diseases.....	96
1.10	APP and neurogenesis.....	100
1.11	Hypothesis and aims of the study.....	104
1.11.1	Hypothesis.....	104
1.11.2	Aims.....	105
Chapter 2	Materials and Methods	106
2.1	Materials.....	107
2.1.1	Buffers, solutions and cell culture media.....	107
2.1.2	Antibodies used in this study.....	107
2.2	Methods.....	107
2.2.1	Neural stem/progenitor cell (NSPC) culture.....	107
2.2.2	AlamarBlue Assay.....	108
2.2.3	EdU incorporation assay of cell proliferation.....	109
2.2.4	NSPC differentiation and cell fixation.....	109
2.2.5	Immunoblotting.....	110
Chapter 3	Role of CysC in amyloid precursor protein - induced of neural proliferation stem/progenitor cells	116
3.1	Introduction.....	117
3.2	Materials and Methods.....	119
3.2.1	All materials used in this chapter were shown in Table 2.1.....	119
3.2.2	In vitro measurement of cell number and cell proliferation.....	119
3.2.3	Effect of conditioned medium on NSPC proliferation.....	119

3.2.4	Determination of the levels of sAPP α and CysC.....	120
3.2.5	Immunodepletion of sAPP α , A β or CysC.....	120
3.2.6	Real-time PCR.....	120
3.3	Results.....	121
3.3.1	Analysis of APP levels and proliferation of NSPCs derived from Tg2576 mice and background strain C57Bl/6 \times SJL mice.....	121
3.3.2	Analysis of proliferation of NSPCs derived from APP knock-out...	124
3.3.3	Effect of neurosphere conditioned medium on NSPC proliferation..	124
3.3.4	Effect of sAPP α and A β peptides on NSPC proliferation.....	126
3.3.5	Levels of CysC and expression of CysC mRNA.....	128
3.3.6	Role of secreted CysC in stimulating NSPC proliferation.....	132
3.4	Discussion.....	135
Chapter 4	Effect of cysteine protease inhibitors on NSPC proliferation	140
4.1	Introduction.....	141
4.2	Materials and Methods.....	142
4.2.1	Materials.....	142
4.2.2	NSPC proliferation.....	143
4.2.3	Preparation of CysC – Sepharose.....	143
4.2.4	Affinity purification and analysis of fraction.....	143
4.3	Result.....	144
4.3.1	Effect of protease inhibitors on NSPC proliferation.....	144
4.3.2	CysC affinity purification of proteases in conditioned medium.....	146
4.4	Discussion.....	149
Chapter 5	Role of APP in neural stem/progenitor cell differentiation	154
5.1	Introduction.....	155
5.2	Materials and Methods.....	157
5.2.1	Materials used in this study are shown in Table 2.1.....	157
5.2.2	NSPC differentiation.....	157
5.2.3	Immunocytochemistry	157
5.2.4	Image capture and data analysis.....	159

5.3	Results.....	159
5.3.1	Differentiation of NSPC.....	159
5.3.2	CysC is not involved in APP induced differentiation.....	163
5.3.3	sAPP α , A β ₄₀ or A β ₄₂ do not influence neuronal and astrocytic differentiation.....	163
5.4	Discussion.....	164
Chapter 6	Discussion	173
	References	184

Abstract

Alzheimer's disease (AD) is a progressive neurodegenerative condition that commonly affects people over the age of 65. There are currently no effective treatments which prevent or delay the progression of AD. The disease is characterized by two major pathological hallmarks in the brain, extracellular amyloid plaques and intracellular neurofibrillary tangles.

The β -amyloid protein ($A\beta$) is the major component of the amyloid plaques and is considered to play a central role in AD pathogenesis. $A\beta$ is produced by proteolytic processing of the amyloid precursor protein (APP). However, the normal function of APP remains unclear despite numerous studies. Understanding the biological function of APP may improve our understanding into the molecular basis of AD pathology.

APP expression has been reported to increase in neurons during embryogenesis. In addition, enhanced neurogenesis has been observed in the brains of AD patients and in transgenic mice which overexpress APP. However, other studies have reported impaired neurogenesis in APP transgenic mice. Thus, the studies presented here were aimed at investigating the role of APP in neural stem/progenitor cell (NSPC) proliferation and differentiation.

To test the effect of APP on NSPC proliferation, NSPCs derived from human APP overexpressing transgenic (Tg2576) mice and APP knockout (APP KO) mice and their corresponding background strain wild type (WT) mice were cultured in a proliferation medium containing growth factors and NSPC growth was measured. The study found that Tg2576 NSPCs proliferated more rapidly than NSPCs from WT

control mice. In contrast, NSPCs derived from APP KO mice proliferated less readily than the corresponding background strain mice.

The secreted fragments of APP, A β and sAPP α , have been reported to affect neurogenesis. The molecular basis of the effect is uncertain. To examine whether sAPP α or two major isoforms of A β , A β ₄₀ and A β ₄₂, were responsible for APP - induced NSPC proliferation, the WT NSPCs were grown in proliferation medium containing sAPP α , A β ₄₀ or A β ₄₂ and the cell proliferation was measured by examining cell viability. No significant difference in cell viability was found between the non - treatment groups and the groups treated with sAPP α , A β ₄₀ or A β ₄₂. Besides, immunodepletion of secreted fragments of APP (sAPP α , A β) from Tg2576 conditioned medium did not lower NSPC proliferation, indicating that neither sAPP α nor A β contributed to the proliferation effect.

To examine whether other secreted factors might be involved in the proliferation effect, the ability of cell conditioned medium to stimulate proliferation was tested. NSPC conditioned medium from Tg2576 cultures was found to increase proliferation while conditioned medium from APP KO cultures was found to have a lower effect on proliferation. The effect on proliferation was found to be due to a secreted factor, cystatin C (CysC), which has previously been reported to promote NSPC proliferation. Immunodepletion of CysC from the Tg2576 conditioned medium removed the stimulatory effect of APP on NSPC proliferation. mRNA levels of CysC in APP KO cells were lower than the wild-type control cells. Therefore, CysC is likely to be a major mediator of APP-associated NSPC proliferation.

CysC is well characterized as a cysteine protease inhibitor. Proteases have been reported to play a role on cell proliferation, thus regulation of proteases activities may affect NSPC proliferation. To examine whether the effect of CysC on NSPC proliferation was due to an effect on one or more cysteine proteases, NSPCs were treated with the broad spectrum protease inhibitor, chymostatin, the specific cysteine protease inhibitor, E-64, the aspartic protease inhibitor, pepstatin A and the serine and cysteine protease inhibitor, antipain, after which cell proliferation was examined. The cysteine protease inhibitor, E-64 and the serine/cysteine protease inhibitor, antipain, both of which are cell membrane impermeable were found to stimulate NSPC proliferation at low concentrations. However, no extracellular cysteine proteases were identified bound to CysC in the conditioned medium by the CysC affinity chromatography. Thus more studies may be required in the future to identify the specific target of CysC.

To test the effect of APP on NSPC differentiation, NSPCs prepared from Tg2576, and APP KO and corresponding background strain wild type mice were cultured in a differentiation medium lacking of growth factors. Cultures were immunostained for markers of neurons (β III tubulin), astrocytes (GFAP) and oligodendrocyte progenitors (NG2) various days after incubation to assess cell differentiation. APP overexpressing (Tg2576) NSPC cultures displayed more β III tubulin⁺ and GFAP⁺ cells than the wild type cultures, while APP KO NSPC cultures had fewer β III tubulin⁺ and GFAP⁺ cells compared to the corresponding wild type cultures. No significant difference was found in NG2 expression.

Previously published studies on the effect of sAPP α , A β ₄₀ and A β ₄₂ on NSPC differentiation have yielded conflicting results. Therefore, one of the aims of this study was to examine whether CysC, sAPP α , A β ₄₀ and A β ₄₂ influence NSPC differentiation. The APP KO NSPCs were grown in differentiation medium containing CysC, sAPP α , A β ₄₀ and A β ₄₂, and the cells were fixed and immunostained for markers of neurons (β III tubulin), astrocytes (GFAP) and oligodendrocyte progenitors (NG2). No difference in the expression of the neuronal and glial markers between the control group and groups treated with CysC, sAPP α , A β ₄₀ and A β ₄₂ was observed. It was concluded that neither CysC nor sAPP α , A β influence NSPC differentiation.

In summary, the study found that APP can promote NSPC proliferation *in vitro*, and the effect is mediated by secreted CysC. However, the molecular basis of CysC action requires further studies. In addition, APP was also found to play a role on NSPC differentiation *in vitro*. Adult neurogenesis has been reported to play important roles in learning, memory and mood regulation. Therefore, therapeutic strategies focusing on anti – A β production by alteration APP processing should be considered carefully. Besides, APP - directed neuronal differentiation of NSPCs could perhaps contribute to stem cell therapy for neurodegenerative disease treatments including AD.

Acknowledgements

Firstly, I would like to thank my supervisors, Professor David Small, Dr. Lisa Foa, Dr Kaylene Young for letting me to join their group and be their student. I would like to thank David for his patience, guidance, support, great encouragement and for pushing me when I needed to be pushed. Thanks to Lisa for offering me support and good advice. Thanks to Kaylene for giving me good suggestions and for training me in neural stem cell culture techniques.

I would also like to give my special thanks to Dr. Amos Hung and Dr. Marta Bolós. Thanks to Dr. Amos Hung for his patience in training me in the stem cell culture techniques when I started my degree. Thanks to Dr Marta Bolós for her friendship and help, and for being willing happy to collaborate.

My thanks also go to Dr. Edgar Dawkins for advice on tissue preparation and immunocytochemistry, to Dr. Hao Cui with immunoblotting, to Dr. Claire Dickson and Dr. Dave Gell for affinity chromatography, and Dr. Katherine Southam for risk assessment writing and very good advice on how to give a final PhD presentation. Thanks to Dr. Adele Vincent for kind help when I joined the lab, Thanks to Dr. Robert Gasperini for fluorescence microscopy, and Fiona Stennard for proofreading my thesis. Thanks to Jenny Smith and Claire Hadrill for keeping the lab running efficiently.

Thanks to past team members, Dr. Camilla Mitchell, Dr. Lila Landowski and thanks to my two nice desk neighbors Adrian Thompson and Macarena Pavez for their help

and support. I would like to thank new team members, Agnieszka Zbela, Dr. Carlie Cullen, Daniel Johnson, Daniela Achatz, Dr. Dino Premilovac, Lachlan Brown, Loic Auderset, Megan O' Rourke, Richard Pinferi, Dr. Shiwei Wang and Yilan Zhen for their friendship and support.

Finally, I would like to thank my parents and friends, although they are miles away from me. I would like to my dear uncle who took care of me when I was young although he has passed away. It is my regret that I was not able to be with him at the last time of his life.

Publication record

Peer-reviewed publications:

Hu, Y., Hung, A. C., Cui, H., Dawkins, E., Bolós, M., Foa, L., Young, K. M., and Small, D. H. (2013). Role of cystatin C in amyloid precursor protein induced proliferation of neural stem/progenitor cells. *J Biol Chem* 288, 18853-18862.

Bolós M., **Hu, Y.**, Foa, L., Young, K. M., and Small, D. H. (2014). Neurogenin 2 mediates amyloid- β precursor protein-stimulated neurogenesis *J Biol Chem* 289, 31253-31261.

Dawkins, E., Gasperini, R., Cui, H., **Hu, Y.**, Bolós, M., Vincent, A.J., Foa, L., Young, K.M. and Small, D.H. (2014) The N-terminal region of the β -amyloid precursor protein of Alzheimer's disease binds to PIP-rich microdomains on the surface of hippocampal neurons. *J. Neurosci. Res.* 92, 1478-1489.

Small, D. H., **Hu, Y.**, Bolós, M., Dawkins, E., Foa, L., and Young, K. M. (2014). Beta-Amyloid Precursor Protein: Function in Stem Cell Development and Alzheimer's Disease Brain. *Neurodegener Dis* 13, 96-98. (Review).

Manuscript in preparation:

Hu Y., Bolós M., Foa, L., Young, K. M., and Small, D. H. (2015) Role of amyloid precursor protein in neural stem/progenitor differentiation.

Conference publication:

Hu Y., Hung A.C., Cui H., Dawkins E., Foa L., Young K.M. and Small D.H. APP stimulates neural stem/progenitor cell proliferation by increasing cystatin C secretion, (2013), 33th annual meeting of Australian neuroscience society, Melbourne.

Hu, Y., Bolós, M., Dawkins, E., Foa, L., Young, K.M. and Small, D.H. Cystatin C regulates neural stem or progenitor cell proliferation, (2014), 34th annual meeting of Australia neuroscience society Adelaide.

Bolós M., **Hu Y.**, Foa L., Young K.M. and Small D.H. The role of neurogenin-2 in APP stimulated neural stem/progenitor cell differentiation, (2014), 34th annual meeting of Australia neuroscience society Adelaide.

Bolós M., **Hu Y.**, Young KM., Foa L., Small D.H. APP regulates neural stem or progenitor cell differentiation through neurogenin-2 expression, (2014) FENS Forum of neuroscience, Milan, Italy.

Hu Y., Bolós M., Young KM., Foa L., Small D.H. Role of amyloid precursor protein in neural stem/progenitor cell proliferation and differentiation, (2015), 9th international Brain Research Organization (IBRO) congress, Rio de Janeiro, Brazil.

List of Abbreviations

AChE	Acetylcholinesterase
AD	Alzheimer's disease
ADAM	A disintegrin and metalloproteinase
AICD	APP intracellular domain
ALS	Amyotrophic lateral sclerosis
AMPA	α -amino-3 hydroxy-5-methyl-isoxazolepropionic acid
AMPA	AMPA receptor
ANOVA	Analysis of variance
APLP1	APP – like protein 1
APLP2	APP – like protein 2
apoE4	Apolipoprotein E4
APH-1	Anterior pharynx-defective 1
APP	Amyloid precursor protein
APP KO	APP knockout
APP dKO	APP/APLP2 knockout
APP tKO	APP/APLP1/APLP2 triple knockout
APP ^{sw}	APP carrying the Swedish mutation (K670N/K671L)
BACE	β – site APP cleaving enzyme
BACE KO	BACE knockout
BBB	Blood - brain barrier
BDNF	Brain derived neurotrophic factor
BIN1	Bridging integrator 1
BSA	Bovine serum albumin
bHLH	basic helix -loop - helix
BMP	Bone morphogenetic protein
CAA	Cerebral amyloid angiopathy
CalB	Calbindin
CALHM1	Calcium homeostasis modulator
CCR	Chemokine receptors
CNS	Central nervous system
CNBr	Cyanogen bromide

CSF	Cerebrospinal fluid
CysC	Cystatin C
CTS3	Gene coding CysC
CuBD	Copper – binding domain
DAPI	4',6-diamidino-2-phenylindole
DCX	Doublecortin
DG-SGZ	Subgranular zone of dentate gyrus
ECM	Extracellular matrix
EdU	5-ethynyl-2'-deoxyuridine
EGC	Embryonic germ cells
EGF	Epidermal growth factor
EGFR	EGF receptor
DMEM	Dulbecco's modified Eagle's medium
DMSO	Dimethyl sulfoxide
EOAD	Early - onset AD
ESC	Embryonic stem cell
FAD	Familial AD
FBS	Fetal bovine serum
FGF	Fibroblast growth factor
FGFR	FGF receptor
FTD	Frontotemporal dementia
GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
GC	Granule cells
GDNF	Glial cell line-derived neurotrophic factor
GFAP	Glial fibrillary acid protein
GFLD	Growth factor like domain
GLAST	Glutamate aspartate transporter
GSK-3 β	Glycogen synthase kinase-3 β
GWAS	Genome-wide association study
HCCAA	Hereditary cystatin C amyloid angiopathy
HBD	Heparin binding domain
HD	Huntington's disease
ICC	Immunocytochemistry
IF γ	Interferon gamma

IGF	Insulin growth like factor
iPSC	Induced pluripotent stem cells
KPI	Kunitz protease inhibitor domain
LDL	Low-density lipoprotein
LDLR	Low density lipoprotein receptor
LRP	Lipoprotein protein receptor
LTD	Long term depression
LTP	Long term potentiation
LOAD	Late - onset AD
MAP	Microtubule - associated protein
MAPK	Mitogen activated protein kinase
MCI	Mild cognitive impairment
mGluR7	Metabotropic glutamate receptor 7
MSC	Mesenchymal stem cell
NCT	Nicastrin
NGF	Nerve growth factor
Ngn2	Neurogenin2
NICD	Notch intracellular domain
NMDA	N-methyl-D-aspartate
NMDAR	NMDA receptor
NSC	Neural stem cell
NSPC	Neural stem/progenitor cell
NFT	Neurofibrillary tangles
NMJ	Neuromuscular junction synapses
OB	Olfactory bulb
OPC	Oligodendrocyte progenitor
PBS	Phosphate-buffer saline
PD	Parkinson's disease
PEN	Presenilin enhancer
PDGF-APP ^{sw} ,Ind	APP carrying the Swedish (KM670N/671L) and Indiana (V717F) mutation
PGC	Periglomerular cell
PID	Phosphotyrosine interacting domains

Pin1	Peptidyl-prolyl cis-trans isomerase NIMA-interacting1
PS	Presenilin
PSA-NCAM	Polysialylated neural cell adhesion molecule
PTB	Phosphotyrosine binding domain
PVDF	Polyvinylidene difluoride
RMS	Rostral migratory stream
sAD	Sporadic AD
SDS-PAGE	Sodium dodecylsulfate polyacrylamide gel electrophoresis
SGZ	Subgranular zone
Shh	Sonic hedgehog
siRNA	Small interfering RNA
SNAP25	Synaptosomal-associated protein 25
TACE	Tumour necrosis factor α converting enzyme
TBS	Tris buffered saline
TBS-T	Tris buffered saline Tween -20
TFs	Transcriptional factors
TH	Tyrosine hydroxylase
Tip60	Tat-interactive protein 60
TLX	A member of the nuclear receptor family of intracellular transcription factors
TNF α	Tumour necrosis factor α
TREM2	Triggering receptor expressed on myeloid cells 2
Trk	High affinity nerve growth factor receptor
UR	Unstructured region
VEGF	Vascular endothelial growth factor
V-SVZ	Ventricular – subventricular zone
Wnt	Wingless
WT	Wild type

Chapter 1 Introduction

1.1 Alzheimer's disease (AD)

The study presented in this thesis was focused on elucidating one of the functions of amyloid precursor protein (APP) of Alzheimer's disease (AD). Because of the importance of APP for AD, this chapter first reviews AD and the central role of APP in AD pathogenesis.

AD is a neurodegenerative disease and the leading cause of dementia in the elderly. Approximately 2% of the population between the ages of 65 and 69 years suffer from AD. However, this figure dramatically increases to 25 to 30% between the ages of 80 to 85 (Vandenberghe & Tournoy 2005). The pathology of AD was first described in a 51 year old female patient by the German psychiatrist, Alois Alzheimer (Alzheimer 1907). There is still no cure, or even an effective treatment, to delay the progression of AD.

1.1.1 Clinical features of AD

During the progression of AD, brain cells are damaged or become dysfunctional, which gradually leading to the loss of memory, thinking, cognition and other physiological functions regulated by the brain, including motor activity (Tarawneh & Holtzman 2012). The combination of memory decline, language disturbance, motor activity and cognitive impairment clinically describes dementia (Tarawneh & Holtzman 2012). However, a diagnosis of AD only can be confirmed by post mortem pathological analysis. Therefore, ante-mortem diagnosis of AD depends on the analysis of clinical symptoms (Tarawneh & Holtzman 2012). At an early stage of dementia, diagnosis of AD is difficult because some memory decline is normally associated with aging. At an early stage of AD, patients exhibit mild cognitive

impairment (MCI) and a deficiency in short-term memory and information storage (Bayley et al 2000, McCormick et al 1994, Morris et al 2001). Besides aggressive behavior, the majority of MCI patients show mild psychological symptoms including anxiety, depression, delusions, hallucinations and mood changes (Gauthier et al 1997, Mega et al 1996, Weiner et al 2005). Moreover, patients at this stage have problems with calculation, insight and judgment, but neurological function usually is normal (Tarawneh & Holtzman 2012). When AD develops to a moderate stage, intensive or permanent supervision may be required because patients experience cognitive behavioral problems, somatic symptoms, impairment of recent memories, sleep disturbances and loss of emotional control (Forstl & Kurz 1999, Rebok et al 1991). In the final (severe) stage of AD, patients potentially lose the ability to control bodily functions and have profound circadian rhythm disturbances (Forstl & Kurz 1999, Volicer et al 2001). Therefore, full-time nursing care is required (Herrmann & Gauthier 2008). Clinical diagnosis of dementia mostly occurs during the mild to moderate stages of the disease and death usually occurs 5-8 years after diagnosis (Forstl & Kurz 1999). Patients with dementia most frequently die of bronchopneumonia, cerebrovascular or cardiovascular disease (Brunnström & Englund 2009).

1.1.2 Impact and prevalence of AD

AD accounts for 50%-70% of all cases of dementia. Patients with a moderate to severe stage of AD rely heavily on caregivers, and at a severe stage of the disease usually require institutionalisation. Therefore, the total cost of dementia is a major challenge for the healthcare system. In Australia in 2009, there are more than 321,000 people living with dementia, and the cost of the dementia and aged care sector is at

least USD \$4.9 billion annually, or around 1% of GDP (Access Economics 2009, Australian Institute of Health and Welfare 2012). In the United State of America, 5.2 million people have AD including 200,000 individuals (Hebert et al 2013) under 65 who have early-onset AD (Alzheimer's Association 2006). The total cost of care for people living with AD and other dementias in USA is estimated to be USD \$214 billion (Alzheimer's Association 2014). Furthermore, more than 36 million people worldwide live with dementia and the total estimated worldwide cost is estimated to be around 604 billion (Alzheimer's Disease International 2012). However, the number of people living with dementia in the population around the worldwide is predicted to be 115 million by 2050 (Alzheimer's Disease International 2012), with spending on dementia treatment and care estimated to outnumber that of any other health condition by 2050 (Access Economics 2009). Therefore, it is crucial to find therapies that can block or delay the progression of AD. Development of a therapy for AD would not only relieve the enormous burden on the health care system, but more importantly, it would improve the quality of life for individuals with AD and their families.

1.1.3 Pathology of AD

The development of lesions in the vulnerable sites of the brain is thought to be the central pathological processes in many neurological diseases, including AD (Braak et al 2006). Thus, investigation into the role that pathological lesions play in disease pathogenesis is important. A number of pathological lesions are typically found in the post-mortem AD brain. Pathological lesions such as extracellular plaques, intracellular neurofibrillary tangles and adipose inclusions were originally identified using silver-staining techniques by Alois Alzheimer (Alzheimer 1907, Alzheimer et al 1995). Neuron and synapse loss (Terry et al 1991b, Whitehouse et al 1982), and

amyloid deposits in brain blood vessels (cerebral amyloid angiopathy; (Revesz et al 2003)) were identified in later histopathological investigations of AD brain sections.

1.1.3.1 Neurofibrillary tangles

One of the hallmarks of AD pathology is the intracellular accumulation of neurofibrillary tangles (NFTs) in neurons. The NFTs develop in the cell body and can spread into the dendrites. Once nerve cells have been destroyed, the NFTs convert to an extraneuronal structure which can eventually be taken up and degraded by astrocytes (Braak & Braak 1991b). In the brain of AD patients, NFTs generally exhibit a common and highly predictable pattern of distribution with only minor inter-individual variations (Braak & Braak 1991b). NFTs develop in a temporal progression that correlates with the clinical symptoms of AD (Bierer et al 1995). The progression of NFT pathology has been characterized in six stages (Braak & Braak 1991b). In stages I and II, NFTs accumulate in the transentorhinal region and in the CA1 region of the hippocampus (transentorhinal stage). NFTs develop severely in transentorhinal and entorhinal regions but moderately in the hippocampus during stage III and IV (limbic stage). Finally, numerous NFT aggregates can be seen in the hippocampus and isocortex in stage V and VI. The isocortex is severely affected in these stages. Therefore, stages V and VI can be interpreted as an isocortex stage (Braak & Braak 1991b). Stage I and II are a nonclinical pathology, and can be found in individuals as young as 20. The limbic stage (Stage III and IV) is associated with emerging AD pathology, and can be found in 10% of adults of aged 50 or over, and 50% of adults 80 years old or more (Braak et al 2011). The isocortex stage (stage V and VI) is regarded as fully developed AD, which is present in 10% of adults aged 80, and 20% of adults aged 90 (Braak & Braak 1995).

NFTs are composed of paired helical filaments that consist of an abnormally hyperphosphorylated form of the microtubule-associated protein (MAP) tau (Grundke-Iqbal et al 1986, Kosik et al 1986, Wood et al 1986). Tau is a type II MAP, that modulates microtubule stability inside axons and facilitates the assembly of microtubule bundles within the nerve cell (Mandelkow & Mandelkow 1995). Normally, tau binds to the outer ridge of the neuroprotofilaments and works as an anchor to stabilise microtubule bundles or regulates the dynamics of microtubule assembly (Al-Bassam et al 2002). When tau is hyperphosphorylated, it detaches from protofilaments and loses the ability to promote assembly of, and maintain the structure of, microtubules (Grundke-Iqbal et al 1986). In addition, the cytosolic abnormally hyperphosphorylated tau can also cause normal tau and other MAPs to dissociate from microtubules and to block microtubule assembly (Iqbal et al 2008). Moreover, the level of abnormally hyperphosphorylated tau in the AD brain is significantly higher than the normal brain (Khatoon et al 1992). Therefore, hyperphosphorylated tau may lead to disruption of the microtubule assembly and may cause affected neurons to undergo a slow but progressive neurodegeneration (Iqbal et al 1994). However, the mechanism causing the development of abnormal tau aggregates needs clarification.

1.1.3.2 Amyloid plaques

Amyloid plaques are focal lesions that are another hallmark of AD (Braak & Braak 1991b). Generally, extracellular amyloid plaques are present in all six layers of the isocortex, but are most abundant in layer V. Unlike NFTs, amyloid plaques develop in an unpredictable pattern of distribution pattern with predominant individual variation. The progression of amyloid plaques can be roughly classified into three stages. In

stage A, a low density of amyloid deposits is found in the isocortex, particularly in the basal portions of the frontal, temporal and occipital lobe but deposits are almost absent from the hippocampus (Braak & Braak 1991b). However, a few amyloid plaques can be found in the junction between the hippocampus and isocortex. In stage B, amyloid plaques are encountered in nearly all the isocortical association areas and are also present in the white matter underlying the cortex. The hippocampus is mildly affected, but the primary sensory region and the primary motor area remain devoid of amyloid plaques at this stage. In stage C, all isocortex areas are loaded with amyloid in a laminar distribution. Some regions outside the cerebral cortex including the striatum, thalamus, hypothalamus, subthalamic nucleus and red nucleus also exhibit deposition of amyloid, but the hippocampus still harbours comparatively fewer deposits (Braak & Braak 1991b). Clinical pathological studies indicate that amyloid burden does not correlate with the severity and duration of dementia. Furthermore, the size of the plaques does not increase with the progression of AD (Arriagada et al 1992, Bierer et al 1995, Giannakopoulos et al 2003, Gomez-Isla et al 1997, Hyman et al 1993). However, preliminary data from longitudinal amyloid PET imaging studies of living patients suggest that the total amount of amyloid deposits in the cortical mantle increases during the clinical course of AD, and that amyloid plaques spread according to the stages described above (Jack et al 2009).

Amyloid plaques are characterized by accumulation and deposition of amyloid beta peptides ($A\beta$) that are produced from the normal proteolysis of the β -amyloid precursor protein (APP) (Kang et al 1987). $A\beta$ can aggregate into amyloid fibrils that subsequently deposit to form amyloid plaques. Although $A\beta$ is the major ingredient of amyloid plaques (Glenner & Wong 1984, Masters et al 1985, Selkoe et al 1986), a

number of other components in the plaque can also be identified including apoE, clusterin and cystatin C (CysC) (DeMattos et al 2001, May et al 1990, Sanan et al 1994, Yamada et al 1989). Current methods to classify different types of plaques in AD brains are generally based on morphological characteristics (D'Andrea & Nagele 2002). Amyloid plaques can be distinguished into two types depending on the β -pleated sheet conformation of the insoluble fibrous protein aggregates (β -amyloid aggregate) that can be specifically stained with Congo Red or Thioflavin-S (Sipe et al 2010). Plaques that are Thioflavin positive are dense core plaques, while those that accumulate immunoreactive A β , but which are Congo Red or Thioflavin negative, are diffuse plaques (Tagliavini et al 1989, Yamaguchi et al 1988). Plaque types correlate with disease. The Thioflavin-S positive dense core plaques consist of a central mass of amyloid filaments that radially project to the periphery where they interact with neuronal, astrocytic and microglial processes (Serrano-Pozo et al 2011). Several studies indicate that dense core plaques closely correlate with dystrophic neurites (neuronal processes), neuron loss, synaptic loss, and astrocytic and microglial activation (Itagaki et al 1989, Knowles et al 1999, Pike et al 1995, Urbanc et al 2002, Vehmas et al 2003). Dense core plaques, especially those associated with dystrophic neurites, are always present in the brains of AD patients (Sheng et al 1998, Urbanc et al 2002), whereas amorphous diffuse plaques can be found commonly in the brains of elderly individuals with intact cognition (Morris et al 1996, Yamaguchi et al 1989). The diffuse plaques are not typically related to dystrophic neurites and synapse loss (Masliah et al 1990, Morris et al 1996). Plaque-associated neuritic dystrophy provides evidence that amyloid plaques contribute to neurodegeneration in AD. Moreover, the phosphorylation of tau can be induced by the major component of the amyloid plaque – A β (Chabrier et al 2012, Gotz et al 2001, Jin et al 2011, Tokutake et al 2012).

Therefore, tau aggregates are most probably a downstream pathology mediated by A β .

1.1.3.3 Cerebral amyloid angiopathy (CAA)

Amyloid deposited in the wall of cerebral and leptomeningeal vessels is called cerebral amyloid angiopathy (CAA). CAA pathology commonly appears in individuals with dementia, and can be observed in approximately 80% of AD patients (Jellinger 2002). Furthermore, post-mortem longitudinal studies reveal that CAA may play a role in cognitive decline in AD (Arvanitakis et al 2011, Greenberg et al 2004, Pfeifer et al 2002). Cerebral amyloid deposits can often be found in leptomeningeal and cortical arteries, and arterioles, but less frequently is present in capillaries and veins of the central nervous system (CNS) (Pezzini et al 2009). After deposition in the outer membrane, and between smooth muscle cells of the blood vessel, CAA can weaken and destroy blood vessels leading to intracerebral haemorrhage and cerebral infarction (Pezzini et al 2009). The major ingredient of the cerebral amyloid deposits in AD-related CAA is A β (Herzig et al 2004). However, other proteins (CysC, prion protein, transthyretin, and gelsolin) can form deposits in the walls of cerebral blood vessels to form CAA in other diseases (Pezzini et al 2009). For instance, in hereditary CysC amyloid angiopathy (HCCAA), CysC accumulation in the vascular media and adventitia causes progressive loss of smooth muscle cells leading to cerebral haemorrhage (Wang et al 1997).

Studies reveal that vascular lesions may contribute to CAA pathology of AD (Gorelick et al 2011, Thal et al 2003). However, based on autopsy samples based from the Honolulu Asia Aging Study, it is suggested that the CAA burden in AD is

independent of vascular lesions (Launer et al 2008, Pfeifer et al 2002). Therefore, vascular disease may make little contributions to the formation of CAA in AD (Launer et al 2008). Formation of CAA pathology in AD is probably more closely associated with amyloid plaque deposition.

1.1.3.4 Synapse loss

Synaptic transmission is essential for cognition. Therefore, synapse loss is thought to be the major correlate of cognitive impairment in dementia including AD (Hamos et al 1989, Terry et al 1991b). Loss of synaptic contacts in both the neocortex and hippocampus is one of the pathological features of AD, and has been demonstrated to be an early event in AD before neuron loss and tangle formation (Masliah 1995, Scheff et al 2006). Synapse loss occurs predominantly in areas surrounding senile plaques, suggesting that amyloid plaques may contain a reservoir of A β that causes synaptic toxicity (Moolman et al 2004, Tsai et al 2004).

1.2 β - Amyloid (A β)

Extracellular amyloid plaques are primarily composed of A β . The hypothesis that A β is the main cause of dementia in AD is central to the majority of research in both academia and industry, although the direct relationship between A β , memory loss and cognition impairment in dementia has not been fully elucidated. However, considerable evidence from studies using A β overproducing mice supports the view that overproduction of A β leads to amyloid plaque lesions that are related to AD (Citron et al 1992, Citron et al 1997, Selkoe 1993). Furthermore, various studies indicate that various forms of A β aggregates are harmful to neuronal and synaptic

function (Jin et al 2011, Kullmann & Lamsa 2007, Lefterov et al 2009, Shankar et al 2008).

1.2.1 Amyloid precursor protein processing and A β production

A β is derived by proteolytic processing of amyloid precursor protein (APP) (Sisodia et al 1990). APP is a transmembrane protein, and the A β sequence extends from the ectodomain of APP into the membrane spanning-domain (Kang et al 1987). During proteolytic processing, APP can be cleaved by several proteases called secretases. APP is mainly processed by an α -secretase cleavage pathway that is non-amyloidogenic pathway (Sisodia et al 1990) (Fig. 1.1). In this pathway, APP is processed consecutively, at first by α -secretase within the A β sequence. This releases a 100 kDa extracellular soluble form of APP named sAPP α , leaving an 83 amino acid residue fragment containing the transmembrane and intracellular C-terminal region of APP (C83) (Esch et al 1990, Sisodia 1992). C83 spans the plasma membrane and projects into the intracellular space. C83 can be further cleaved by γ -secretase to produce a small fragment of APP (P3) and the intracellular domain of APP (AICD). In the β -secretase pathway (amyloidogenic pathway), APP is initially cleaved by β -secretase to liberate a N-terminal ectodomain (sAPP β) into the extracellular space. The remaining 99 amino acid residue C-terminal fragment (C99) is then further cleaved by γ -secretase to liberate A β and the APP intracellular domain AICD. The A β can be cut by γ -secretase into a peptide ranging from 37 to 42 amino acid residues in length (Qi-Takahara et al 2005). Different variants of A β peptide can be detected as




Figure 1.1 Diagram showing the β -secretase cleavage and α -secretase cleavage pathways of APP proteolytic processing. In the β -secretase pathway, APP is cleaved by BACE1 to generate a secreted ectodomain, sAPP β , and the C - terminal membrane spanning domain, named C99. C99 can be further cleaved by γ -secretase to produce an intracellular domain fragment (AICD) and the secreted A β . The α -secretase pathway is a major processing pathway of APP. In this pathway, APP is cleaved by α -secretase to release a soluble extracellular domain fragment, sAPP α . The C-terminal domain remaining in the membrane, termed C83, is further cleaved by γ -secretase to generate a secreted fragment, p3, and the intracellular domain (AICD).

a soluble component of plasma cerebrospinal fluid and other tissues, even in normal physiological conditions (Iwatsubo et al 1994, Seubert et al 1992). However, A β 40 and A β 42 are the most frequent variants that can be found in amyloid plaques (Iwatsubo et al 1994), and A β 42 is more common in CAA than A β 40 (Roher et al 1993). Because of the two extra amino acids, A β 42 is more prone to aggregate and form fibrils, which are suggested to be more toxic than those formed by A β 40 (Dahlgren et al 2002, Jarrett et al 1993). Moreover, an increased amount of A β 42, or an enhanced ratio of A β 42/ A β 40, correlates with hereditary forms of AD (Golde et al 2000, Scheuner et al 1996), whereas a decrease in A β 42 concentration is associated with a lower risk of AD (Weggen et al 2001).

1.2.2 Aggregation of A β

A β contains a hydrophobic C-terminal region that contributes significantly to β -sheet conformation. The region in the N terminal domain is important for α -helical or β -strand structural switching (Soto et al 1994, Zagorski & Barrow 1992). A β predominately forms an α -helical conformation in organic solvents, while in aqueous solution it is more prone to be present in a β -sheet configuration that is closely associated with A β fibrillogenesis (Lansbury 1999). However, the transition of A β structure is highly dependent on pH and hydrophobicity, which may influence its aggregation and amyloid deposition in the AD brain (Soto et al 1994, Wood et al 1996b).

A β containing 39 to 40 amino acid residues is kinetically soluble for hours to days whereas the peptide containing 42 to 43 amino acid residues aggregates immediately (Jarrett et al 1993). In general, multiple A β aggregates can be found in amyloid

plaques in the AD brain: fibrils, dimers, monomers, tetramers and the more soluble oligomeric forms of A β (Shankar et al 2009). The mechanism of A β aggregation to form A β deposits and amyloid plaques is still unclear. However, *in vitro* studies on synthetic A β proteins suggest several models. Amyloid fibrils are possibly formed through a nucleation-dependent process in which A β aggregation requires monomeric A β to convert slowly in a nucleation phase to form A β oligomers. The addition of A β monomers to these oligomers, which are often referred to as ‘seed’, results in fibril elongation to form longer A β aggregates. The reaction eventually reaches a dynamic equilibrium between A β aggregates and A β monomers. A nucleation-dependent mechanism is identified by a high concentration dependence on ‘seed’ (nucleus) formation, which leads to A β aggregation (Jarrett et al 1993). Esler and his colleagues (Esler et al 2000) reported that A β aggregation is propagated by a template dependent ‘dock-lock’ mechanism that is mediated by two distinct kinetic processes. In the ‘dock’ phase, A β binds to the pre-existing amyloid template in a reversible manner. However, A β can, in a time-dependent transition, interact with the template in an irreversible manner. In this ‘lock’ phase, the deposited ‘dock’ A β interacts irreversibly with the template and is now referred to as ‘lock’ A β . The locked A β may then become a new template to which ‘dock’ A β may bind (Esler et al 2000). This provides a template driven mechanism for a transition from A β peptides to A β aggregates. However, recent structural analysis suggests possible mechanisms of A β aggregation. These mechanisms have been recently reviewed (Tycko & Wickner 2013).

1.2.3 A β toxicity

A β is naturally produced by healthy cells. However, mutations within or around the A β sequence lead to elevated A β production or changes in A β isoforms that have been found to correlate with early-onset AD (Selkoe 1993). Results from many studies suggest that A β plays a central role in AD progression, although the underlying mechanism remains to be elucidated (Braak & Braak 1991a, Small et al 2001, Terry et al 1991a).

A β induces impaired long-term potentiation (LTP) and thereby alters synaptic transmission. This may contribute to memory and cognitive decline in AD. LTP induces growth of dendritic spines, whereas long-term depression (LTD) facilitates spine shrinkage and synaptic loss (Kullmann & Lamsa 2007). LTP is thought to be closely related to learning and memory (Otto et al 1991), thus excessive accumulation of A β may inhibit LTP and lead to memory decline. The memory of a learned behaviour by a normal rat was disrupted after intracerebroventricular injection of the extracted A β oligomers (Shankar et al 2008). At high concentrations, soluble A β oligomers extracted from the cortex of AD subjects were found to inhibit synaptic LTP and to promote LTD of synaptic transmission (Shankar et al 2008). A β -induced LTD can lead to endocytosis of synaptic α -amino-3 hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors (Hsieh et al 2006, Shankar et al 2008). A reduction of AMPA receptors is necessary and sufficient to induce loss of spines and synaptic (N-methyl-D-aspartate) NMDA responses (Hsieh et al 2006). Recent studies also suggest that A β triggers a reduction in the number of cell-surface NMDA receptors (Hsieh et al 2006, Snyder et al 2005). Increased A β levels are reported to block uptake of glutamate, which leads to the accumulation of glutamates in the

synaptic cleft (Li et al 2009). Synaptic transmission can requires activation of NMDA receptors. However, persistently elevated glutamate can desensitize NMDA receptors and thereby disrupt synaptic transmission (Li et al 2009). Therefore, pathologically increased levels of A β may possibly change NMDA and AMPA receptor signalling pathways, induce LTD, impair LTP, and cause synaptic transmission dysfunction.

A β is also suggested to cause functional and structural damage to neurons. Numerous studies show that high levels of synthetic A β_{40} and A β_{42} can bind to cell membranes and increase membrane permeability, leading to destruction of cellular structures (Arispe et al 1993, Demuro et al 2005, Lin et al 2001). In addition, A β_{42} internalization through the endosome/lysosome pathway can lead to lysosomal leakage (Arispe et al 1993, Yang et al 1998). However, natural oligomers of secreted A β isolated from culture medium or from brain tissues at nanomolar concentrations have little effect on lipid membrane permeability. This may be explained by the fact that A β oligomers induce neuronal dysfunction rather than perturb the membrane (Sanchez-Mejia et al 2008). For example, soluble A β oligomers purified from AD cortex are able to induce tau hyperphosphorylation, which further triggers progressive collapse of the microtubule cytoskeleton (Jin et al 2011). Thus, A β oligomers contribute to neuronal dysfunction whereas A β_{40} or A β_{42} probably induce structural destruction.

Calcium homeostasis dysregulation is considered to be another possible mechanism of neuronal dysfunction. A β increases the basal levels of calcium and therefore renders neurons more vulnerable to glutamate-induced excitotoxicity eventually leading to neuronal death (Mattson et al 1992, Price et al 1998). A β is also reported to

form calcium channels in the cell membrane (Demuro et al 2005). In addition, A β has been reported to induce Ca²⁺ influx through binding to NMDA receptors (De Felice et al 2007) or via stimulation of L-type Ca²⁺ channels (Ho et al 2001, Sberna et al 1997). A recent paper indicates that A β may impair metabotropic glutamate receptor 7 (mGluR7) on cholinergic neurons, which indirectly affects calcium influx (Gu et al 2014). This implies that disruption of calcium homeostasis may cause selective degeneration of cholinergic neurons, which is one of the pathological features in AD (Gu et al 2014). Moreover, a polymorphism in the calcium homeostasis modulator (CALHM1) 1 gene correlates with an increased risk of AD (Dreses-Werringloer et al 2008). This polymorphism of CALHM1 increases A β levels through alteration of Ca²⁺ permeability and cytosolic Ca²⁺ levels (Dreses-Werringloer et al 2008).

However, it is worth mentioning that A β -induced neuronal death cannot be thought as the sole cause of cognitive decline in AD (Small et al 2001). In fact, dysfunctional synaptic transmission or the loss of synapses are more correlated with cognitive loss than deposition of A β plaques and cell death (Braak & Braak 1991a, Terry et al 1991a).

1.2.4 Possible physiological functions of A β

A β has been intensively studied for its role in the pathogenesis of AD. However, its physiological functions have attracted less attention. Several studies have suggested that A β could play a neuroprotective or neurotrophic role at low concentrations (Giuffrida et al 2009, Whitson et al 1989). Low doses of A β could enhance learning and memory through facilitating induction and maintenance of LTP (Morley et al 2010). A small increase in A β levels by suppression of extracellular A β degradation

could lead to increased release of synaptic vesicles and enhanced neuronal activity in neuronal culture (Abramov et al 2009). Furthermore, increased A β levels may promote spontaneous excitatory postsynaptic currents without changing the inhibitory currents (Abramov et al 2009). Another study found that application of synthetic A β_{42} at picomolar concentrations potentiated synaptic transmission, but at nanomolar concentrations A β_{42} depressed synaptic transmission (Puzzo et al 2008). However, A β -induced synaptic effects are dependent on an optimal concentration of A β , and higher or lower concentrations lead to synaptic transmission dysfunction (Abramov et al 2009). This is also supported by other reports in which abnormal reduction of A β levels in mice deficient for APP (Seabrook et al 1999), PS1 (Saura et al 2004) or BACE1 (Laird et al 2005) were correlated with synaptic transmission dysfunction. Thus, A β may have an important physiological function in the brain. However, more work will be needed to determine whether effects that are seen are due to physiological or pathophysiological actions of A β .

1.3 Biology of amyloid precursor protein

1.3.1 Structure of the APP family

Amyloid precursor protein (APP) is a type I integral membrane protein and widely expressed in the mammalian brain (Kang et al 1987). The human APP gene, which is located on the long arm of chromosome 21, spans about 240 kb and consists of 18 exons (Lamb et al 1993, Yoshikai et al 1990). The A β encoding region is found in the 16th and 17th exons (Lemaire et al 1989, Suh & Checler 2002). APP undergoes alternative mRNA splicing to generate 8 different isoforms of APP comprising from 365 (APP365) to 770 (APP770) amino acid residues (Jacobsen et al 1991, Kitaguchi et al 1988). APP770, APP 751 and APP695 are the most common isoforms expressed

either in brain or peripheral tissues. APP770 and APP751, which contain a 56 amino-acid Kunitz protease inhibitor domain (KPI), are abundantly produced in glial cells (Donnelly et al 1988, Palmert et al 1988). APP770 differs from APP751 by the insertion of a 19 amino-acid residue OX-2 domain adjacent to the KPI domain of APP751. The OX-2 domain is only present in the APP770 isoform (Donnelly et al 1988). The shorter isoform, APP695, is normally produced in neurons, while less is produced in non-neuronal cells (Haass et al 1991). APP695 is the primary source of APP in the brain (Palmert et al 1988, Sisodia et al 1993). Interestingly, expression of APP751/770 isoforms is dramatically increased in astrocytes and microglia after injury (Siman et al 1989) and a substantial increase in APP695 expression can be detected during neuronal differentiation (Zheng & Koo 2011). However, the mechanism that regulates the expression of alternative APP isoforms is still poorly understood.

Two APP homologues have been identified in mammals: the amyloid like protein-1 (APLP1) and the amyloid like protein-2 (APLP2) (Wasco et al 1992, Wasco et al 1993). Generally, APLP1 is restricted to neurons whilst APP and APLP2 are ubiquitously expressed in all tissues (Zheng & Koo 2011). The APP family members have a relatively conserved structure with a longer N-terminal extracellular region (ectodomain), a single membrane-spanning domain and a shorter C-terminal intracellular domain (cytoplasmic domain) (Dyrks et al 1988).

The extracellular region of APP contains several structural domains for which the crystal structure is available (Fig. 1.2). These are the E1 domain and the E2 domains (Dahms et al 2010, Rossjohn et al 1999). The E1 domain, which is close to the N-




Figure 1.2 Domain structure of APP and its homologues. D1, heparin - binding domain; D2, metal binding domain including copper - binding domain and zinc - binding domain; Ac, acidic region; KPI: Kunitz-type protease inhibitor domain; E2: heparin binding domain; UR: unstructured region. YENPTY: adaptor protein binding region.

terminus of APP, contains a heparin binding domain (HBD) that forms a part of a growth factor like domain (GFLD) rich in cysteine (Small et al 1994), and a metal (copper and zinc) binding domain (Bush et al 1993, Hesse et al 1994, Reinhard et al 2005, Rossjohn et al 1999). Between the E1 domain and the E2 domain, there is an acidic region that is rich in aspartic and glutamic acid, and the KPI domain. The KPI domain is not found in APLP1. However, the OX-2 domain is located between the acidic region and KPI domain in APP770. The E2 domain is closer to the C-terminus and comprises another HBD/GFLD region and an unstructured region (UR) that contains α - and β - secretase cleavage sites (Mok et al 1997). The ectodomain of APP is followed by the transmembrane domain that has the γ -secretase cleavage site (Dyrks et al 1988, Weidemann et al 2002). The cytoplasmic domain of APP contains of a YENPTY motif that is proposed to be involved in transcriptional regulation and to interact with adaptor proteins (Cao & Sudhof 2001, Sabo et al 2003). Like the E1 and E2 domains of the extracellular region, the YENPTY motif of the intracellular region is highly conserved in APP family members (Jacobsen & Iverfeldt 2009). However, the A β sequence is not conserved. It is only present in APP, but not in APLP1 and APLP2 (Kang et al 1987).

1.3.2 APP trafficking and processing

After expression, APP is translocated into the endoplasmic reticulum (ER) and is post-translationally modified in the Golgi apparatus by glycosylation, sulfation, phosphorylation and palmitoylation (Bhattacharyya et al 2013, Selkoe 2001). Immature APP species undergo N- and O- glycosylation in the Golgi (Oltersdorf et al 1990, Weidemann et al 1989) where the N- and O- glycosylated mature APP is then trafficked to the cell surface (Koo et al 1996). At the cell surface, APP can be

internalized through clathrin-induced endocytosis via interactions with the YENPTY motif in the cytoplasmic domain of APP (Yamazaki et al 1996). After internalization, APP is delivered to the endosome, after which it can be recycled back to the cell surface, or it can be delivered to the lysosome where it can be further degraded (Perez et al 1999, Yamazaki et al 1996).

APP undergoes γ -secretase cleavage after the ectodomain shedding by α or β cleavage (Sisodia 1992). The α -secretase cleavage of APP to produce sAPP α may occur at the cell surface (Koo et al 1996), but some α -secretase cleavage may occur in the secretory and endocytic vesicles (Parvathy et al 1999, Tomita et al 1998). The β -site APP cleaving enzyme-1 (BACE1), which is the β -secretase (Cai et al 2001, Sisodia et al 1990), is mainly distributed in membranes of the Golgi and endosomes (Koo & Squazzo 1994). This suggests that APP can be cleaved in both secretory and endocytic steps. Therefore, β -secretase processing and A β production may commonly occur in acidic compartments in which β -secretase is most activated (Koo & Squazzo 1994, Vassar et al 1999). The site of intracellular γ -secretase cleavage of APP has not been well characterized, but several studies indicate that the γ -secretase complex is located, and has activity in, multiple cell compartments including the ER, Golgi, cell membrane and endosomal/lysosomal system (Buxbaum et al 1988, Kaether et al 2002, Pasternak et al 2004, Tarassishin et al 2004).

APLP1 and APLP2 are thought to undergo similar proteolytic processing to APP (see section 1.2.1) although they lack the A β peptide region. Soluble forms of APLPs could be detected in human brain and cerebrospinal fluid (Webster et al 1995). In addition, Eggert (et al 2004) found that both homologues of APP were cleaved via α

and β secretase like-pathways. Furthermore, p3 and A β -like fragments of APLP2, and a p3-like fragments of APLP1, were detected in the conditioned medium of stably transfected SH-SY5Y cells (Eggert et al 2004, Minogue et al 2009). Moreover, increased levels of soluble APLP2 and its C-terminal fragments were found in ADAM10 (α -secretase) over-expressing transgenic mice (Endres et al 2005). APLP1 processing was reported to be independent of BACE 1 (β -secretase) reactivity because BACE1 inhibitors had no effect on APLP1 shedding (Eggert et al 2004, Minogue et al 2009). However, a recent study on BACE1- knockout and BACE1- overexpressing mice showed that BACE 1 tightly regulates the level of full length and soluble APLP1 and APLP2 (Sala Frigerio et al 2010). Deficiency in a functional part of γ -secretase (presenilin1 (PS1)), not only leads to failure of A β secretion in neurons, but also causes accumulation of C-terminal fragments of APP and APLP1 in neurons (Naruse et al 1998). These studies support the view that processing of APP homologues is similar to that of APP.

1.3.3 APP secretases

APP is cleaved by proteases, called secretases. Three APP secretases have been designated as the α -, β - and γ -secretases.

1.3.3.1 α - Secretase

The “non-amyloidogenic” pathway of APP processing involves α -secretase cleavage. This pathway is the primary pathway through which APP undergoes constitutive shedding. APP is cleaved within the A β sequence by α -secretase on the C- terminal side of amino acid residue 16 of the A β sequence (Sisodia 1992, Sisodia et al 1990). Thus, the primary route of APP processing precludes the generation of intact A β . The

enzymes that participate in α -secretase cleavage are members of the disintegrin and metalloproteinase (ADAM) family. Tumor necrosis factor- α converting enzyme (TACE) was the first identified ADAM, and is known as ADAM17 (Buxbaum et al 1998). One year after the identification of ADAM17 as an α -secretase, ADAM9 and ADAM10 were also reported to function as α -secretases (Koike et al 1999, Lammich et al 1999). ADAM10 overexpression leads to a major increase in α -secretase cleavage of APP (Lammich et al 1999, Postina et al 2004), a result that was later confirmed by Kuhn et al (2010). In addition, ADAM17-knockout mice show reduced production of sAPP α , although small inhibitory RNA (siRNA) mediated knockdown of ADAM17 did not show a reduction in α -secretase dependent APP processing (Buxbaum et al 1998, Kuhn et al 2010). However, mice with an ADAM9 deletion had neither major abnormalities nor a significant difference in α -secretase associated APP processing (Kuhn et al 2010, Weskamp et al 2002). Therefore, the major constitutive α -secretase activity may be contributed by ADAM10, with ADAM9 and 17 are more likely only to be involved only in regulation of α -secretase cleavage (Kuhn et al 2010, Moss et al 2011).

Besides APP processing, ADAMs play a physiological role in various types of cellular processing. ADAMs are suggested to act as sheddases that catalyze the release of soluble extracellular domains from proteins (Huovila et al 2005). This process is essential for the release of cytokines and growth factor ligands. For example, ectodomain shedding by ADAM10 can lead to release of epidermal growth factors (EGFs) and β cellulin (Sahin et al 2004). A recent paper indicates that senescence-associated release of transmembrane proteins requires proteolytic processing by ADAM17 (Effenberger et al 2014). ADAM17 can regulate the release

of tumour necrosis factor- α by acting as a sheddase (Black et al 1997). ADAMs-induced ectodomain shedding participates in cellular processing of transmembrane proteins, thus ADAMs may cleave APP to produce soluble APP fragments that play a role in downstream cellular signalling (Demars et al 2011, Hoey et al 2009, Huovila et al 2005).

1.3.3.2 β -Secretase

In the amyloidogenic pathway, APP is cleaved by β -secretase at the N-terminal end of the A β sequence to produce the soluble fragment sAPP β , and a membrane bound fragment C99 which can be subsequently cleaved by γ -secretase to produce A β . The β -site APP cleaving enzyme 1 (BACE1) is a transmembrane aspartic protease that has all the characteristics of a β - secretase (Hussain et al 1999, Lin et al 2000, Vassar et al 1999, Yan et al 1999). BACE1 is primarily expressed in the brain and contributes the essential role of a β -secretase (Hussain et al 1999, Lin et al 2000). A homologue of BACE1, known as BACE2, can cleave APP in the middle of the A β domain and is expressed widely in peripheral tissues, but has a lower expression levels in brain than BACE1 (Fluhrer et al 2002, Lin et al 2000).

BACE1 has two aspartate domains that form the active site; mutation of either aspartate residue leads to inactivation of BACE1 (Hussain et al 1999, Knops et al 1995). In AD patients, both the expression and activity of BACE1 is elevated (Fukumoto et al 2002, Holsinger et al 2006, Yang et al 2003). BACE1 was reported to be the enzyme that contributes to the majority of A β production by neurons, and BACE1 knockout (BACE1 KO) mice show no significant neuronal A β and sAPP β production (Dominguez et al 2005, Luo et al 2001). BACE2 KO mice also show a normal healthy phenotype, but mice with a double knockout of BACE1 and BACE2

display an increased incidence of lethality phenotype (Dominguez et al 2005). BACE1 deficiency did not affect A β production in glia, supporting the idea that BACE2 may be responsible for A β generation in glia in the brain of AD patients (Dominguez et al 2005). Thus, both BACE1 in combination with BACE2 may be needed for the maintenance of a healthy phenotype and normal APP processing.

BACE1 KO mice do not exhibit a major phenotypic difference to wild type mice (Roberds et al 2001). In addition, deletion of the *BACE1* gene rescues the memory deficiency in APP transgenic mice (Ohno et al 2006). This suggests that inhibition of BACE1 activity is a viable strategy for AD therapy. However, further studies have found that cognitive and emotional abnormalities, and synaptic dysfunction and spatial memory deficits appear in BACE1 KO mice (Laird et al 2005). Furthermore, studies on both BACE1 KO and over-expressing mice indicate that BACE1 contributes to myelination (Hu et al 2006, Willem et al 2006). Moreover, electroencephalographic recordings reveal that kainic acid-induced seizures occur more frequently in BACE1 KO mice than in wild type mice (Hu et al 2010). BACE1 can act as sheddase for a variety of substrates including neuregulin, the β -subunits of the voltage-gated sodium channels, interleukin-1 receptor 2, as well as the low-density lipoprotein (LDL) receptor-related protein (Cai et al 2012, Klaver et al 2010, Vassar et al 2009). Therefore, BACE1 plays a variety of roles in cellular processes, and it is thus essential to monitor the potential side effects of BACE1 inhibitor use as a therapeutic treatment for AD (Klaver et al 2010).

1.3.3.3 γ -Secretase

After ectodomain shedding by α - or β -secretases, the membrane bound fragments C99

and C83 are further cleaved by the γ -secretase to release AICD, A β or p3. γ - Secretase is a transmembrane protein complex that contains at least four protein subunits, including presenilins (PS1 or PS2), anterior pharynx-defective 1 (APH-1), presenilin enhancer 2 (PEN 2) and nicastrin (NCT) (Kimberly et al 2003b). The formation of the γ -secretase complex occurs in sequential steps, with APH-1 and NCT initially forming a stable complex with PS, followed by the addition of PEN 2 to form the active γ - secretase (Hu & Fortini 2003, LaVoie et al 2003).

All the components (PSs, APH-1, PEN 2 and NCT) are important for stability, maturation and activity of γ -secretase (Francis et al 2002, Goutte et al 2002, Kimberly et al 2003b). Deletion of a conserved hydrophilic domain of nicastrin inhibits γ -secretase and thereby inhibits A β production (Shirotani et al 2003, Yu et al 2000). Deletion of APH-1 causes a reduction in PS levels (Lee et al 2002, Shirotani et al 2004) whereas overexpression of APH-1 can cause accumulation of PS (Takasugi et al 2003). Therefore, APH-1 may act as a cofactor for PS that contributes to γ -secretase activity. PEN-2, a membrane-spanning component of the γ -secretase complex, coordinately regulates PS1 proteolytic processing (Luo et al 2003, Steiner et al 2002).

The PSs are suggested to be the major catalytically active components of γ -secretase (De Strooper et al 1998, Figueroa et al 2002, Wolfe et al 1999). Mammalian presenilins have two family members, presenilin 1 (PS1) and presenilin 2 (PS2). Cleavage of α - and β -secretase is not affected by *PS1* knockout, whereas cleavage by γ -secretase is completely blocked, causing accumulation of APP C-terminal fragments and a drop in A β levels (De Strooper et al 1998). PS mutations cause a

decrease in γ -secretase function. This leads to the incomplete digestion of the A β -peptide, which may contribute to increased vulnerability of the brain, and early onset of the inherited form of AD (De Strooper 2007).

Apart from APP, γ -secretase has several other substrates. One of the most notable substrates is the cell-surface receptor, Notch (De Strooper et al 1999). Cleavage of APP and Notch by γ -secretase appears to be similar (Kimberly et al 2003a). The Notch intracellular domain (NICD) is liberated by γ - secretase cleavage, resulting in its translocation to the nucleus where, it forms complexes with DNA binding proteins, and thereby acting as a regulator of downstream gene transcription (Jarriault et al 1995, Struhl & Adachi 1998). PS1 deficiency leads to a reduction in Notch proteolytic processing of Notch, and a decreased level of NICD. Thus the Notch signalling pathway is affected by PS1 (De Strooper et al 1999, Wong et al 1997). Notch signalling plays a variety of roles in CNS development including modulation of neural stem cell (NSC) proliferation, differentiation, maturation and survival (Lathia et al 2008). In addition, integrin regulates epithelial cell differentiation by modulating Notch activity (Gomez-Lamarca et al 2014). Therefore, inhibition of γ -secretase for the treatment of AD may lead to toxicity caused by reduced Notch signalling (De Strooper et al 1999).

1.3.4 Function of APP

The biological properties of APP have been studied intensively since its sequence was first reported in 1987 (Kang et al 1987). However, the physiological roles of APP still remain unclear (Zheng & Koo 2006). Studies on APP knockout (APP KO) mice suggest that APP may play a variety of interesting roles (Dinet et al 2011, Hung &

Selkoe 1994, Puig et al 2012). APP KO mice (Zheng et al 1996) display a number of physiological abnormalities (Magara et al 1999, Seabrook et al 1999, Steinbach et al 1998) compared to wild-type mice. However, it must be noted that the two homologous proteins, APLP1 and APLP2 may provide some functional compensation for APP deletion or deficiency (Heber et al 2000). Triple knockout of APP, APLP1 and APLP2 is lethal during embryonic development, suggesting a developmental function may be blocked when compensation by other family members is lost (Herms et al 2004). Taken together, various studies suggest that APP plays an essential role in the maintenance and normal physiology of the brain (Heber et al 2000, Zheng & Koo 2006). In this section, studies related to functions of APP in neurodevelopment are reviewed.

1.3.4.1 Role of APP in cell adhesion and neurite outgrowth

The E1 and E2 regions in the extracellular domain of APP have been shown to interact with extracellular matrix proteins: laminin (Kibbey et al 1993), collagen type 1 (Behr et al 1996) heparin sulfate proteoglycans (Mok et al 1997, Small et al 1994) and glypican-1 (Williamson et al 1996b), which suggests that APP plays a role in cell-substrate adhesion (Small et al 1999). APP and other family members can form homo- or hetero-dimers (Soba et al 2005), supporting the idea that APP is involved in cell- to cell contacts. Moreover, APP has been suggested to interact with proteins implicated in cell adhesion, such as integrins (Yamazaki et al 1997), neuron-glia cell adhesion molecule (Young-Pearse et al 2008), and transient axonal glycoprotein 1 (Ma et al 2008a). For example, APP and β 1-integrin were colocalized on the cell surface at contact sites in neural cells (Yamazaki et al 1997). Soluble APP was shown to induce neurite outgrowth by interacting with β 1-integrin (Young-Pearse et al 2008). siRNA mediated knockdown of APP expression causes defects in neuronal

migration that was associated with a cell adhesion deficiency (Young-Pearse et al 2007, Young-Pearse et al 2008).

The up-regulation of APP expression has been reported in growing neurites (Clarris et al 1995, Hung et al 1992), and down-regulation of APP seems to block neurite outgrowth (Allinquant et al 1995, Milward et al 1992). The effect of APP on neurite outgrowth may be driven by the physical cell adhesion characteristics of APP (Small et al 1994). APP was reported to bind to the neurite-promoting site of the transmembrane glycoprotein, laminin to promote neurite outgrowth (Kibbey et al 1993). Furthermore, the APP heparin binding domain that is responsible for cell adhesion (Small et al 1999), is also suggested to be involved in regulation of neurite outgrowth (Small et al 1994). Indeed, APP-induced neurite outgrowth can be blocked by application of a peptide homologous to the heparin-binding region (Small et al 1994, Williamson et al 1996a). Moreover, hippocampal neurons grown on a substrate of APP transfected CHO cells, showed short-term neuronal adhesion and longer-term neurite outgrowth (Qiu et al 1995). Therefore, the APP-stimulated neurite outgrowth may be attributed to an increase in cell-cell or cell-extracellular matrix interaction.

Several other studies indicate that APP can directly or indirectly affect neurite outgrowth and axonal path finding (Gakhar-Koppole et al 2008, Hasebe et al 2013, Hoareau et al 2008, Young-Pearse et al 2008). For instance, the soluble forms of APP, sAPP α and sAPP β , that are produced by α - and β -secretase cleavage respectively, can reportedly induce neurite outgrowth by binding to the p75 neurotrophin receptor (Hasebe et al 2013). The APP homologue, APLP2 was reported to have a role in the regulation of cell migration, neurite outgrowth and axonal path-finding (Thinakaran et al 1995). Thus, neurite outgrowth may be regulated by both APP and APLP2.

1.3.4.2 Role of APP in synaptogenesis and synaptic plasticity

APP is also suggested to participate in the regulation of synaptogenesis and synaptic plasticity. During development, the mitral cells of the olfactory bulb show dramatically increased in APP expression at a particular stage when the mitral cell dendrites are contacted by approaching neurites from olfactory receptor neurons (Clarris et al 1995). In neurons, APP is rapidly transported to the synaptic terminals after synthesis (Koo et al 1990, Moya et al 1994, Sisodia et al 1993) and its expression is predominantly up-regulated during the critical period of synaptogenesis (Clarris et al 1995, Wang et al 2009).

Synaptogenesis defects in APP KO mice may lead to a functional deficiency in neurotransmission and to long-lasting effects on synaptic plasticity and synaptogenesis (Priller et al 2006, Ring et al 2007, Zheng et al 1996). These defects may cause deficits in grip strength and locomotor activity in mice lacking APP (Ring et al 2007). For example, an in vitro study showed that deletion of APP can increase the number and cumulative strength of excitatory synapses during synaptogenesis (Priller et al 2006, Steinbach et al 1998), which may contribute to a hypersensitivity to kainate-induced seizure. APP-null mediated behavioural abnormalities are rescued by application of the soluble ectodomain form of APP, supporting the idea of a role for sAPP in synaptic function (Ring et al 2007). Endogenous A β is also proposed to be necessary for hippocampal synaptic transmission and plasticity at certain concentrations (Puzzo et al 2011, Puzzo et al 2008). Such a role may be one explanation for the cognitive impairment that occurs in APP-null mice (Dawson et al 1999, Seabrook et al 1999).

APP KO mice usually display mild phenotypes, as the functional deficiency may be masked by compensatory changes in APLP expression (Heber et al 2000, Korte et al 2012). The APP homologue, APLP2 is abundant in axon terminals in glomeruli and is present in pre- and postsynaptic compartments in the olfactory bulb (Thinakaran et al 1995). Studies from double knockout APP^{-/-} /APLP2^{-/-} (dKO) mice show that the APP family of proteins are essential for neuromuscular junction (NMJ) synapse patterning-APP/APLP2 dKO mice show impaired neuromuscular formation, with a reduced quantity of synaptic vesicles and impaired synaptic transmission (Ring et al 2007). APLP1/APLP2 dKO mice lack presynaptic nerve terminal sprouting, although they show normal endplate patterning, which means that APLP2 may play an essential role at NMJ synapses that cannot be compensated for by APP (Klevanski et al 2014). Therefore, APP and APLP2 may have distinct and divergent roles at NMJ synapses (Klevanski et al 2014).

APP has also been implicated in synaptic plasticity and neurotransmission (Hoe et al 2009a, Puzzo et al 2008, Yang et al 2009a). APP was reported to modulate levels and activities of Ca(v)1.2 L type calcium channels and to affect short-term plasticity through the regulation of calcium currents in GABAergic neurons (Yang et al 2009a). APP was also found to exert an effect on excitatory synaptic transmission via a change in two ionotropic glutamate receptors, AMPA receptor (AMPA) and NMDA receptor (NMDAR) trafficking (Hoe et al 2009a, Hoe et al 2012, Lee et al 2010). APP plays a role in expression of the GluA2 subunit of the AMPAR- GluA2, which in turn affects synaptic transmission and plasticity (Isaac et al 2007). Thus APP may affect synaptic function by altering GluA2 (Lee et al 2010). Moreover, APP also mediates

synaptic function through an alteration of cell-surface NMDAR expression (Hoe et al 2009a). The NMDR is a calcium channel, and calcium permeability of the synapse membrane is crucial for synaptic plasticity (Cousins et al 2009). However, it has been suggested that APP's effects on synaptic transmission may be partially mediated by endogenous A β produced by APP processing because certain concentrations of A β can potentiate synaptic transmission (Puzzo et al 2008).

1.3.4.3 Non- neuronal functions of APP

Several non-neuronal functions of APP and APLPs have been proposed. APP has been reported to participate in blood coagulation (Bush et al 1990, Smith et al 1990). The isoform of APP containing the KPI domain is produced and released by platelets (Bush et al 1990, Gardella et al 1990, Van Nostrand et al 1991a). After stimulation of coagulation, APP, sAPP and A β , which accumulate in α - granules of platelets, is released along with the blood clotting factors that are also stored in the α -granule vesicles (Blair & Flaumenhaft 2009, Bush et al 1990, Gardella et al 1990, Smith et al 1990, Van Nostrand et al 1991b). Due to the presence of the KPI domain, APP may act as a serine protease inhibitor to inhibit the activity of coagulation factor Xia which plays an essential role in the blood coagulation cascade (Scandura et al 1997, Smith & Broze 1992, Smith et al 1990, Van Nostrand et al 1990). Studies on APP KO and APP/APLP2 dKO mice suggest the possibility that APP is involved in the regulation of glucose and insulin homeostasis. APP KO mice have hypoglycaemia until adulthood, whereas the APP/APLP2 double knockout mice display hyperinsulinaemia and die 24 hours after birth (Needham et al 2008). In addition, enhanced expression of adipocyte APP has also been found in individuals with obesity, and this supports a role for APP in the development of adipose tissue inflammation and insulin resistance

in obesity (Lee et al 2009, Lee et al 2008b). Moreover, up-regulated APP expression has been identified in several types of cancer including oral squamous cell carcinoma, human pancreatic adenocarcinoma and colon cancer (Ko et al 2004, Venkataramani et al 2010). Interestingly, siRNA-mediated APP knockdown significantly reduces tumour cell growth. Thus, APP expression may be related to tumour cell growth (Venkataramani et al 2010). The fact that APP has a trophic role and can promote cell proliferation (Ayuso-Sacido et al 2010, Saitoh et al 1989) is consistent with this idea. However, another study shows that the intracellular domain of APP may activate transcription of the tumour suppressor gene *P53* (Sumioka et al 2005).

1.3.4.4 APP signalling

Activation of specific cell signalling transduction pathways by APP is presumably important for APP to exert physiological effects, such as neurite outgrowth (Allinquant et al 1995), neural stem cell proliferation and differentiation (Clarris et al 1995, Hu et al 2013, Masliah et al 1992) and cell viability (Murayama et al 1996). However, the mechanisms underlying these effects have not been well clarified. In this section, the role of APP as a putative cell-surface receptor is discussed, as is role of the intracellular domain of APP (AICD).

1.3.4.4.1 APP may act as a cell -surface receptor

APP has been proposed to be a cell-surface receptor, because it shares in structural, post-translational modification and proteolytic processing similarities to Notch, a cell surface receptor involved in cell growth (De Strooper et al 1999, Selkoe & Kopan

2003). APP is also proposed to be a G-protein coupled receptor in a ligand-dependent and ligand-specific signalling (Okamoto et al 1995). Furthermore, APP has been reported to activate serine/threonine kinases and to stimulate the mitogen activated protein kinase (MAPK) pathway that transfers signals from the cell surface to the nucleus (Murayama et al 1996). In addition, the extracellular matrix protein, reelin binds the E1 domain of APP, causing a reduction in A β production and promoting neurite outgrowth (Hoe et al 2009b, Hoe et al 2006). Moreover, the diffusible molecule netrin-1 reportedly can interact with APP and participate in the regulation of A β production (Lourenco et al 2009). More recently, APP was shown to act as a co-receptor in netrin-1 mediated neural navigation and commissural axon outgrowth (Rama et al 2012). Although APP interacts with extracellular matrix proteins, the idea that APP may act as a cell-surface receptor is supported by that F-spondin's ability to bind to the E2 domain of APP, APLP1 and APLP2 (Ho & Sudhof 2004). As a signalling glycoprotein secreted by neurons, F-spondin may exert effects on neuronal development and repair (Peterziel et al 2011). However, binding of F-spondin to APP can lead to blockage of β -secretase cleavage. Therefore, F-spondin is proposed to regulate APP processing (Ho & Sudhof 2004). Although several physiological ligands can interact with APP, APP induced intracellular signalling transduction is the strongest evidence for the idea that APP is a cell-surface receptor (Dawkins & Small 2014).

1.3.4.4.2 Role of APP in intracellular signalling

APP is a substrate for regulated intramembrane proteolysis. This is a mechanism that regulates membrane protein activity, and has been implicated in a wide range of biological processes (Brown et al 2000, Lichtenthaler & Steiner 2007). The C-

terminal region of proteins produced by the γ -secretase translocate to the nucleus and activate gene transcription (Ebinu & Yankner 2002). The intracellular domain of APP (AICD) is suggested to translocate to the nucleus, where it regulates transcriptional activation (Cupers et al 2001, Hebert et al 2006, Lichtenthaler et al 2011), although AICD easily undergoes degradation under normal conditions (Kimberly et al 2001). However, AICD may be stabilized by interaction with adaptor proteins (Kimberly et al 2001, Small et al 2005). AICD can be produced in different forms because the ϵ -cleavage by γ -secretase generates AICD fragments that start further towards the C-terminus (Sastre et al 2001). Caspase-dependent AICD cleavage starts at a C-terminal position upstream of the ϵ -cleavage site (Lu et al 2000).

There are three sequence motifs in AICD that have been suggested to have functional significance. The first one is the ⁶⁵³YTSI sequence, that is proposed to contribute to basolateral sorting of APP (Lai et al 1998), and participates in tyrosine-mediated and clathrin-based endocytic sorting (Bonifacino & Traub 2003). The second region, the ⁶⁶⁷VTPEER sequence motif of AICD, has been implicated in certain pathophysiological processes. For example, Thr668 in the sequence was found to undergo increased phosphorylation in AD patients (Lee et al 2003). Peptidyl-prolyl cis-trans isomerase NIMA-interacting 1 (Pin1), a prolyl isomerase, may affect the turnover of APP by halting GSK3 β -induced phosphorylation at Thr668, as Pin1 overexpression reduces A β whereas a knockout of Pin1 results in an enhanced A β yield (Ma et al 2012a, Pastorino et al 2006). The third functional sequence motif, YENPTY, has drawn the most attention and been studied intensively. The YENPTY region has been shown to interact with many adaptor proteins that have phosphotyrosine binding (PTB) or phosphotyrosine interacting domains (PID)

(Borg et al 1996). APP homologues all contain the YENPTY motif, which means that all APP family members may interact with similar adaptor proteins and may function similarly as transcriptional regulators (Bressler et al 1996, Zheng & Koo 2011). This is consistent with the evidence for functional redundancy among APP homologues from APP knockout mice studies.

The YENPTY motif has been shown to bind to a variety of adaptor proteins: X11 family proteins (Borg et al 1996), Fe65 (Fiore et al 1995), Disable-1 protein (Homayouni et al 1999), JNK interacting protein 1 (Scheinfeld et al 2002b), ShcA/C (Tarr et al 2002) and growth factor receptor-bound protein 2 (Zhou et al 2004). Fe65 was the first identified APP binding partner (Fiore et al 1995) and has been investigated intensively. Fe65 binds to the YENPTY motif in a manner that is independent of tyrosine phosphorylation (Scheinfeld et al 2002a). Fe65 overexpression results in enhanced plasma membrane translocation of APP, and increased production of sAPP α and A β (Sabo et al 1999). This demonstrates that the interaction of APP with FE65 probably affects APP processing and trafficking. However, FE65 expression may lead to inhibition of APP maturation as well as a reduction in A β production when thr668 has been phosphorylated (Ando et al 2001). Therefore, FE65 may bind to both ⁶⁶⁷VTPEER and YENPTY to mediate FE65-dependent gene transactivation (Cao & Sudhof 2001, Sumioka et al 2005).

The interaction of FE65 with AICD can recruit another protein, TIP60, and this AICD/FE65/TIP60 complex is suggested to be involved in transcriptional activation of several target genes (Kim et al 2003, Pardossi-Piquard et al 2005, Zhang et al 2007b). Studies of AICD's role in cell proliferation indicate that the

AICD/FE65/TIP60 complex may work as a negative regulator of neural stem/progenitor cell (NSPC) proliferation via down regulation of the epidermal growth factor receptor (EGFR) mediated gene transcription (Ayuso-Sacido et al 2010, Zhang et al 2007b). In addition, genes that participate in cell death and apoptosis, such as P53, are suggested to be a target of AICD-dependent activation (Alves da Costa et al 2006, Checler et al 2007, Ozaki et al 2006). Cell death and apoptosis occur in transfected cells expressing AICD (Konietzko 2012, Lu et al 2000), while those expressing mutant versions that contain no FE65 binding domain show little effect on cell survival. These findings demonstrate that the interaction of AICD with FE65 may be essential for mediating P53 - induced cell death (Konietzko 2012).

Apart from targets on EGFR or P53, AICD has also been reported to form multiple protein complexes affecting *APP* gene transcription (von Rotz et al 2004). In addition, the intracellular domains of APP and APLP are implicated in the regulation of neprilysin, which is an A β -degrading enzyme (Pardossi-Piquard et al 2005). Furthermore, APP and APLP2 have been shown to regulate cholesterol metabolism via AICD nuclear signaling that may involve modulation of the lipoprotein protein receptor 1 (LRP1) promoter (Liu et al 2007).

1.4 Risk factors for AD

AD can be separated into forms that can be distinguished according to the age of onset or their heritability. Early-onset AD (EOAD) is generally a familial AD (FAD), and presents before 65 years of age. Late-onset AD (LOAD) is generally not inherited, and is sometimes called sporadic AD (sAD), although this term may not be entirely appropriate. It usually occurs over the age of 65. FAD accounts for up to 5%

of AD cases, while LOAD contributes to the majority (~ 95%) of AD cases (Reitz et al 2011). Genetic studies suggest that an increase in A β production may be the cause of all forms of EOAD (Citron et al 1992, Duff et al 1996), whereas LOAD may be associated with a defect in removal of A β from the brain (Miners et al 2008), although this is not yet clearly demonstrated. Apart from A β production, metabolism as well as clearance, education, diet, physical activity, smoking, alcohol, and drugs are also suggested to correlate with the development of AD (Cummings et al 1998, Lannfelt 1996).

1.4.1 Familial AD (FAD)

FAD is an autosomal dominant disorder. Three genes have been firmly identified as responsible for the pathophysiology of FAD, with APP and presenilin genes (PS1 and PS2) all causing changes in APP processing and A β production. These three genes have been found to bear AD linked mutations that are diagnostic disease biomarkers given the very high penetrance (>85%) and mostly autosomal dominant inheritance (Reitz et al 2011).

1.4.1.1 APP mutations

FAD mutations in APP occur within or near the A β encoding region, or close to the cleavage sites of the three secretases. All known mutations account for <0.1% of AD cases (Ancolio et al 1999, Eckman et al 1997, Goate et al 1991, Reitz et al 2011). Most dominantly inherited APP missense mutations may lead either to an increase in total A β production or to an increased ratio of A β_{42} /A β_{40} , which accelerates progression of AD (Lichtenthaler et al 2011, Reitz et al 2011, Selkoe 2001).

Mutations in APP co-segregate with FAD. These mutations include those known as Dutch (E693Q) (Levy et al 1990), London (V717I) (Goate et al 1991), Indiana (V717F) (Murrell et al 1991), Swedish (K670N/M671L) (Mullan et al 1992), and Arctic (E693G) (Nilsberth et al 2001). The Swedish double mutation occurs close to the β -secretase cleavage position and results in approximately 6-fold more $A\beta$ generation compared to cells carrying the normal APP sequence (Citron et al 1992, Reaume et al 1996). APP that harbours mutations around the α -secretase cleavage site lowers α -secretase cleavage (Sahlin et al 2007) and results in more APP available to undergo β -secretase cleavage processing. Thus, $A\beta$ production is elevated. γ -secretase cleavage site-related mutations are proposed to increase production of longer and more neurotoxic forms, such as $A\beta_{42}$ (Suzuki et al 1994). In addition, a mutation at codon 715 of the APP770 gene that has been identified in an Indian family with FAD, was reported to decrease $A\beta_{40}$ production without altering $A\beta_{42}$ yield. Hence, the ratio of $A\beta_{42}$ to the total amount of $A\beta$ seems essential for AD development (Ancolio et al 1999).

1.4.1.2 Presenilin mutations

The presenilins form the main catalytic centre of γ -secretase and functionally participate in the γ -secretase-induced proteolytic cleavage of APP (Ahn et al 2010, De Strooper et al 1998, Figueroa et al 2002, Octave et al 2000). Mutations leading to FAD are found in the *PS1* and *PS2* genes, located on chromosome 14 and 1, respectively (Levy-Lahad et al 1995a, Levy-Lahad et al 1995b, Sherrington et al 1995). To date, approximately 180 different AD-linked polymorphisms in the *PS1* gene from 401 families have been characterized, but only 34 AD-associated mutations from 23 families have been found in the *PS2* gene

(<http://www.alzforum.org/mutations/search/genes>). Some *PS* gene mutations are proposed to be associated with the more rapidly progressing and aggressive forms of AD (Selkoe 2001).

Single-nucleotide substitutions have been suggested to account for the majority of AD-associated PS1 and PS2 mutations (Reitz et al 2011). Several reports indicate that point mutations in the *PS* gene lead to elevated production of A β ₄₂, which contributes to the symptoms of FAD (Bentahir et al 2006, Duff et al 1996, Scheuner et al 1996). While PS mutations exert an apparent gain-of-function effect on γ -secretase activity, De Strooper (2007) suggests that the role of presenilin mutations involve a loss or decrease of γ -secretase function. A decrease in γ -secretase leads to the incomplete digestion of A β -peptide, which may contribute to an increased vulnerability in the brain, and thus explain the early onset of the inherited form of AD (De Strooper 2007).

1.4.2 Late onset AD (sporadic AD)

Genetic risk factors for FAD exhibit clear influences on AD symptom development, however known genetic risk factors for sporadic AD do not display clear effects on AD progression. The precise mechanisms by which genetic risk factors lead to AD remain poorly understood (Finelli et al 2014). Several gene variations have been identified as contributing to LOAD risk. Over 20 common genetic variants linked to sporadic AD have been identified through genome-wide association studies (GWAS) (Chouraki & Seshadri 2014). The top ten AD risk genes are ranked and shown in Table (1.1). Several genes are implicated in lipid transportation and metabolism, neuroinflammation, cytoskeleton dynamics as well as endocytosis (Olgiati et al 2011).

Interestingly, down-regulation of any of the proteins expressed by these genes does not affect A β production, implying that the expression of these proteins may contribute to AD via a pathway unrelated to enhancement of A β production (Bali et al 2012).

1.4.2.1 Apolipoprotein E (apoE)- the major risk factor

The most prevalent genetic risk factor for LOAD is the ϵ -4 allele of the *apolipoprotein E (APOE ϵ 4)*, which is found in more than 50% of AD patients (Michaelson 2014). ApoE consists of three allelic variants that are determined by cysteine-to-arginine substitutions at positions 112 and 158 of the amino-acid sequence (Weisgraber et al 1981). The variants are named as apoE 2 (cys112, cys158), apoE 3 (cys112, arg158), and apoE 4 (arg112, arg158) which correspond to the alleles ϵ 2, ϵ 3 and ϵ 4 respectively (Zannis et al 1982). The small difference in amino - acid sequence gives rise to great differences in the structure and function of the isoforms (Mahley et al 2006). The ϵ - 4 allele is more frequently found in AD cases. People carrying two copies of the alleles have up to 12–15 fold increased risk of developing sporadic AD compared to non-carrier subjects. Even one copy of the ϵ 4 allele increases the risk of AD threefold (Bertram et al 2007).

ApoE is the major apolipoprotein expressed in brain. ApoE is also the constituent of several lipoproteins, such as the high and very low density lipoproteins and chylomicrons (Chouraki & Seshadri 2014). In the CNS, apoE is produced mainly in astrocytes and to a lesser extent in microglia (Grehan et al 2001, Pitas et al 1987). Production of apoE in neurons is normally is low, but can occur under certain

physiological and pathological conditions (Xu et al 1999). ApoE-containing lipoproteins are responsible for transport and delivery of CNS lipids and cholesterol to cells (Holtzman et al 2012), and for the activation of neuroinflammation (Lynch et al 2001). They also act as a ligand in receptor-mediated endocytosis of lipoprotein particles, such as in the uptake of cholesterol needed to support synaptogenesis and the maintenance of synaptic connections (Pfrieger 2003a, Pfrieger 2003b).

ApoE has been shown to colocalize with amyloid deposits in both AD and CAA (Namba et al 1991, Wisniewski & Frangione 1992). A positive relationship between plaque density and apoE 4 dose in AD has been shown in post-mortem brain studies (Rebeck et al 1993, Schmechel et al 1993). ApoE may affect the quantity of A β accumulation and deposition as well as the A β 42/A β 40 ratio in an isoform-specific manner. ApoE4 is the most potent in this regard, and apoE2 the least potent (apoE4 > apoE3 >> apoE2) (Fagan et al 2002, Fryer et al 2005, Holtzman et al 2000). Indeed, individuals with apoE2 have a lower risk for AD (Corder et al 1994). In contrast, several papers suggest that apoE is able to inhibit A β aggregation based on in vitro studies (Evans et al 1995, Naiki et al 1997, Wood et al 1996a, Yang et al 1999). These opposite results may be due to differences in A β or the apoE preparation (Holtzman et al 2012).

ApoE isoforms have been implicated in A β clearance. Binding of apoE to soluble A β forms an apoE-A β complex that may be taken up via receptor-mediated endocytosis into neurons, astrocytes and microglia, after which it may be degraded in lysosomes (He et al 2007, Jiang et al 2008, Li et al 2012a, Yamauchi et al 2002, Yang et al 1999, Zhao et al 2014). Receptor-mediated cellular uptake of A β bound to apoE has been

Table 1.1 Genetic risk factors for LOAD

Gene	Protein	Potential Functions
<i>ApoE 2/3/4</i>	Apolipoprotein E	Lipid transport
<i>BIN1</i>	Bridging-integrator 1	Tumour suppressor and cell endocytosis
<i>CLU</i>	Clusterin	Lipid transport
<i>ABCA7</i>	ATP-binding cassette, sub-family A, Member 7	Lipid homeostasis
<i>CRI</i>	Complement receptor 1	Regulation of immune response
<i>PICALM</i>	Phosphatidylinositol-binding clathrin assembly protein	Clathrin-mediated endocytosis
<i>MS4A6A</i>	Membrane-spanning 4-domians, subfamily A, member 6A	Unknown, expected to have receptor activity
<i>CD33</i>	Myeloid cell surface antigen CD33	Myeloid lineage development. Immune response
<i>MS4A4E</i>	Membrane-spanning 4-domians, subfamily A, member 4E	Unknown, cell component
<i>CD2AP</i>	CD2-associated protein	Regulation of actin cytoskeleton

This table shows 10 top risk genes for LOAD, published on <http://www.alzgene.org>; ranked by the AlzGene database meta – analysis.

reported to occur through a low density lipoprotein receptor (LDLR) - LRP1 dependent pathway (Kang et al 2000, Yang et al 1999), and has been considered to be a possible mechanism of A β clearance. Alternatively, A β removal from the brain into the circulation may be modulated by apoE via trafficking across the blood brain barrier (BBB) (Bell et al 2007, Deane et al 2008). A weaker binding affinity of apoE4 for A β may lower A β clearance from the brain (Deane et al 2008). More abundant A β aggregation may therefore be caused by a decrease in A β clearance. Notably, a recent study suggests that apoE4 may negatively affect A β removal by competing with A β for clearance receptors rather than by specific binding to A β (Verghese et al 2013). In vitro studies show that A β can be directly internalized by cells through the LDLR, which is the major receptor for apoE (Basak et al 2012). ApoE has been shown to enhance protease-mediated A β degradation by neprilysin, but apoE 4 shows less efficiency in facilitating proteolysis of A β (Jiang et al 2008). Finally, a more recent study shows that apoE may be involved in the modulation of microglia activation, with apoE enabling the conversion of macrophages to the anti-inflammatory M2 type (Aguzzi et al 2013, Baitsch et al 2011). In this case, apoE4 may possibly be less effective in its anti-inflammatory action (Zhu et al 2012). Studies on mice argue a role of apoE in A β clearance, as APP cross apoE knockout mice exhibits decreased amyloid deposition compared to APP mice with normal apoE (Bales et al 1997). Recently, another study also argues in favour of apoE having a function in A β removal-normal cognitive and neurological function, as well as normal CSF A β and tau proteins levels were described in a patient with a complete absence of apoE due to an APOE frame shift mutation (Mak et al 2014).

Although apoE is the main identified genetic risk factor for LOAD, neuropathological

alterations other than A β deposition, such as neuro-inflammation and synaptic plasticity, have also been found to associate with apoE, (Kuszczyk et al 2013, Zhu et al 2012). However, the precise mechanism by which apoE contributes to AD progression remains to be elucidated.

1.4.2.2 TREM2

Recent studies have identified a rare variant of the triggering receptor expressed on myeloid cells 2 (TREM2) as a significant risk factor for LOAD (Guerreiro et al 2013, Jonsson et al 2013). This unusual variant in the TREM2 gene has also been linked to EOAD in another study (Pottier et al 2013). A missense mutation (rs75932628) in the *TREM2* gene leads to a C to T base pair change that results in a histidine to arginine amino - acid substitution at position 47 (R47H) (Finelli et al 2014). This mutation and other mutations in TREM2 have been implicated not only in AD, but also in Parkinson's disease, frontotemporal dementia (FTD) and amyotrophic lateral sclerosis (ALS) by other independent studies (Cady et al 2014, Cuyvers et al 2014, Feng et al 2014, Rayaprolu et al 2013).

The TREM2 gene located on chromosome 6 encodes a protein comprising 230 amino - acid residues (Paloneva et al 2002). TREM2, is a transmembrane glycoprotein that belongs to the immunoglobulin (Ig) superfamily. The expressed protein has an extracellular domain that is followed by a membrane spanning domain that is joined to a cytosolic domain (Lue et al 2014a). TREM2 serves as an innate immune receptor that is expressed on the cell membrane of monocyte-derived dendritic cells like macrophages, osteoclasts as well as microglia in the CNS (Colonna 2003). Although generally thought of as a myeloid cell specific protein, TREM2 may be found in

neurons and possibly in oligodendrocytes (Guerreiro et al 2013, Kiialainen et al 2005, Sessa et al 2004). However, some studies have failed to detect TREM2 expression in oligodendrocytes (Chertoff et al 2013, Sessa et al 2004), thus warranting further investigation of its expression in this cell type.

Microglia significantly contribute to phagocytosis of apoptotic neurons and clearance of synapses during development and in adulthood (Miyamoto et al 2013, Neumann et al 2009, Wake et al 2011). In the CNS, TREM2 is expressed predominantly in microglia, and microglia have been found to scavenge apoptotic neurons via TREM2 without triggering inflammation (Takahashi et al 2005). Furthermore, microglial TREM2 is suggested to play an essential role in maintaining immune homeostasis of the CNS tissues (Neumann & Takahashi 2007). Therefore, gene variants of *TREM2* may be correlated with dysfunction of microglia.

AD pathological features such as mature and diffuse plaques have been reported in the patients carrying the R47H TREM2 mutation (Guerreiro et al 2013). Furthermore, post - mortem studies of AD brains and non-demented case controls reveal that elevated levels of TREM2 protein are associated with an increase in phosphorylated tau and the marker of apoptosis, activated caspase 3, as well as with the loss of the presynaptic protein SNAP25 (Lue et al 2014b). This result is seen in a transgenic mice model as well, with enhanced TREM2 mRNA and protein levels in microglia around plaques and neurons in APP transgenic mice brains compared to corresponding wild - type controls (Frank et al 2008, Guerreiro et al 2013). Microglia activation marker complement mRNA was detected in plaque-associated microglia as well (Melchior et al 2010). In addition, AD transgenic mice missing one *TREM2*

allele display no significant alteration in their amyloid plaque load, although they do exhibit a 40% reduction in microglia around amyloid plaques (Ulrich et al 2014). However, up-regulation of TREM2 in AD mice reduces amyloid plaque accumulation, attenuates neuroinflammation and decreases neuronal and synaptic loss (Jiang et al 2014). Enhanced A β phagocytosis has been observed in TREM2-overexpressing microglia (Lue et al 2014a). These results suggest that mutations in TREM2 may result in a loss of function leading to a deficiency in microglial-induced amyloid phagocytosis. However, further investigations are required to confirm this hypothesis.

1.4.2.3 CST3

Cystatin C (CysC), is a cysteine protease inhibitor that is found widely in CSF and other tissues (Turk et al 2000, Turk et al 2008). CysC is encoded by the *CST3* gene. A polymorphism in the *CST3* gene resulting in an Ala/Thr transition (G73A) has been identified, which is reported to increase the risk of AD (Bertram et al 2007, Crawford et al 2000), although this idea has been controversial (Dodel et al 2002, Monastero et al 2005). Different findings may be due to the use of different ethnic groups. In addition, allelic variation at the same position may be another reason for opposing results, with either Ala (allele A) or Thr (allele B) of *CST3* being reported as responsible for AD development (Crawford et al 2000, Finckh et al 2000, Kaur & Levy 2012).

CysC has been found to deposit with A β in amyloid-loaded vascular walls (Maruyama et al 1990), and in the core of amyloid plaques in the AD brain (Levy et al 2001). Compared to non-demented cases, much lower CysC levels can be detected

in the CSF of AD patients (Hansson et al 2009, Simonsen et al 2007), although specific neuronal cells in AD patients display higher CysC expression (Deng et al 2001, Levy et al 2001). Apart from AD, alterations in CSF CysC concentration in CSF has been implicated in other neurodegenerative diseases, such as amyotrophic lateral sclerosis (ALS) (Pasinetti et al 2006, Tsuji-Akimoto et al 2009, Urbizu et al 2014, Yang et al 2009b).

There are different views on the role of CysC in AD. CysC may be involved in exacerbation of neurodegeneration, or it may served a neuroprotective role in response to pathological stimulation (Gauthier et al 2011, Kaur & Levy 2012, Maruyama et al 1990, Suzuki et al 2014b, Urbizu et al 2014). In vitro studies show that CysC has a protective function in neuronal cells, as cells from CysC knockout mice (Huh et al 1999) are more vulnerable to in vitro-toxicity compared to cells isolated from wild-type mice (Tizon et al 2010). A role of CysC in CAA-induced haemorrhage has been proposed because CysC co-deposits with A β , and may increase sensitivity to cerebral haemorrhage (Maruyama et al 1990). APP overexpressing mice that express human CysC show a diminished A β level compared to APP mice with normal CysC (Kaeser et al 2007, Mi et al 2007). However, CysC knockout mice crossed with APP overexpressing mice also show enhanced A β degradation (Sun et al 2008).

1.5 APP transgenic and knockout mice

Transgenic mouse models have been widely used in studies of AD pathogenesis, although there are still no mice carrying a single AD associated gene allele that develop all of the features of human AD (Ashe & Zahs 2010). However, APP

overexpressing mouse models have exhibited some of the lesions and symptoms of AD. APP knockout mice probably exhibit biological deficits caused by APP deficiency, and may provide valuable insights into the function of APP (Ashe & Zahs 2010, Duyckaerts et al 2008).

1.5.1 Transgenic mouse model of AD

Transgenic animal models of AD provide strong evidence that A β is the major cause of AD (<http://www.alzforum.org/res/com/tra/default.asp>). The PDAPP mouse bearing a single valine to phenylalanine substitution at residue 717 (Indiana mutation) (Murrell et al 1991) was the first APP mouse model that developed and exhibited similar neuropathology to AD, including A β deposition, neuritic plaques, synaptic loss, astrogliosis as well as microgliosis (Games et al 1995). The Tg2576 mouse has been used widely for studying AD pathogenesis (Frautschy et al 1998, Hsiao et al 1996, Seo et al 2010). This mouse expresses high levels of human APP with the double Swedish mutation (APP_{K670N/M671L}) under the control of the hamster prion protein (PrP) promoter (Hsiao et al 1996). Apart from age-dependent changes in A β levels in the brain CSF, and plasma (Kawarabayashi et al 2001), Tg2576 mice have a 14-fold increase in A β ₄₂ levels and develop A β plaques, and display behavioural deficits and cognitive impairment after 9 months (Hsiao et al 1996). Moreover, Tg2576 mice also show A β -mediated microglial activation (Frautschy et al 1998), and aged Tg2576 mice exhibit severe motor neuron degeneration in the spinal cord (Seo et al 2010). Mice harbouring the FAD mutations in the *PS1* gene produce an elevated level of A β ₄₂, but do not make amyloid plaques (Duff et al 1996). However, mice with both *APP* and *PS* mutations develop aggressive amyloid pathology (Gordon et al 2001). The amyloid plaques that accumulate in APP/PS1 mice develop at a very early

stage - around 2 to 3 months (Gordon et al 2001). In addition, amyloid-associated pathology, including dystrophic synapses, hyperphosphorylated tau, gliosis, and increased microglia activity starts to appear from 1 to 8 months of age in APP/PS1 mice. Memory and cognitive deficits present at approximately 8 months of age (Arendash et al 2001, Radde et al 2006).

While APP transgenic mice exhibit A β plaques, it still remains unclear why they do not form neurofibrillary tangles (Radde et al 2008). However, transgenic mice bearing three mutations in APP, PS and tau respectively, progressively develop amyloid plaques and tangles and replicate certain aspects of AD neuropathology (Billings et al 2005, Oddo et al 2003). Nicotinic $\alpha 7$ receptors are lost selectively in the hippocampus and cortex and synaptic dysfunction is present prior to plaque and tangle pathology (Oddo et al 2005, Oddo et al 2003). Thus transgenic mice provide an opportunity to investigate some aspects of AD pathophysiology and to test certain therapies for AD, even though they do not recapitulate every pathological feature of the disease.

1.5.2 APP knockout mice

APP knockout (APP KO) mice were made by inactivation of the APP gene (Zheng et al 1996). APP KO mice are viable and fertile, and are relatively normal and healthy until 12 weeks of age (Zheng et al 1996). They exhibit decreased body weight (approximately 15%–20% smaller) and have a reduced brain weight (about 10% less) that is correlated with a reduced size of forebrain commissures. This supports the idea that APP may play a role in neurite outgrowth (Magara et al 1999, Zheng et al 1996). A deficiency in grip strength and a reduction in locomotor activity in APP KO mice (Zheng et al 1995) indicates a possible contribution of APP to neuronal or muscular

function. In addition, a profound reduction of proteins in the presynaptic terminal vesicle, and of dendritic proteins in the cortex and hippocampus, are observed in APP KO mice (Dawson et al 1999). This suggests that APP may contribute to maintaining synaptic function during ageing. In addition, an in vivo study describes a significant reduction in spine density, a decrease in the length of apical dendrites, and a decline in dendrite arborization in APP KO mice aged 12 to 15 months, but not at younger ages. This supports the suggestion that APP may be important for dendritic integrity and maintenance in the hippocampus (Tyan et al 2012).

Interestingly, to date, a mild somatic growth deficit is the only abnormality observed in APLP1-KO mice (Heber et al 2000), and no serious abnormalities have been described in APLP2-KO mice (von Koch et al 1997). The lack of phenotype may possibly be explained by functional redundancies of APP family proteins in transgenic mice (Muller & Zheng 2012). However, APLP2/APLP1 and APLP2/APP double knockout mice usually die shortly after birth (Heber et al 2000, von Koch et al 1997). In contrast, APLP1/APP-mice are viable and fertile, and have not been reported to have any additional abnormalities (Heber et al 2000). This suggests that there may not be compensation between APLP2 and other family members and it suggests a critical physiological role for APLP2 (Muller & Zheng 2012).

1.6 Current and prospective therapeutic approach for AD

At present, there are no clinically proven therapies that can block or even to delay progression of AD. Current therapeutic approaches only ameliorate the symptoms and arrest the rate of development of the disease's pathological symptoms.

1.6.1 Current therapeutic treatment for AD

1.6.1.1 Acetylcholinesterase inhibitors

Loss of cholinergic synapses has been considered for many years to be closely associated with AD because cholinergic neurons are profoundly lost in the basal forebrain of AD patients (Bartus et al 1982, Coyle et al 1983). Cholinergic neurons are essential for memory and concentration, as well as for other cognitive processes. Hence, improvement of AD symptoms can be achieved by prolonged cholinergic stimulation (Sarter & Bruno 2004). Acetylcholinesterase (AChE) is the enzyme that degrades acetylcholine in the synaptic cleft to halt cholinergic neurotransmission. Inhibition of AChE could extend the duration of cholinergic neuron transmission in the brain, leading to mild relief of AD symptoms (Giacobini 1998). Inhibition of AChE by three approved AChE inhibitors: donepezil, galantamine and rivastigmine has been found to decrease AD symptoms to a small extent in the short term (Prvulovic et al 2010, Wilkinson et al 2009). Unfortunately, drug resistance has been reported with AD progression and several side effects have been also reported (Takeda et al 2006). When moderate-to-severe AD patients were administered a combination of donepezil and a N-methyl-D-aspartate (NMDA) receptors antagonist, memantine, there was a lessening in several AD-associated symptoms, although the clinical relevance of this benefit is unclear (Riordan et al 2011).

1.6.1.2 Memantine

Chronic long-term activation of NMDA receptors causes pathological influx of Ca^{2+} , and may ultimately lead to excitotoxicity and neuronal cell death (Butterfield & Pocernich 2003, Dodd 2002, Holscher 1998). $\text{A}\beta$ was reported to disrupt the glutamatergic system and Ca^{2+} homeostasis (Mattson et al 1992, Mattson et al 1999,

Parameshwaran et al 2008). Thus, NMDA receptor induced excitotoxicity is considered to be one of the possible causes of neurodegeneration in AD (Danysz & Parsons 2012, Mattson et al 1999, Parameshwaran et al 2008, Parsons et al 1998).

Memantine is the only NMDA receptor antagonist clinically used for AD treatment, and acts via effects on the glutamatergic system (Danysz & Parsons 2012, Reisberg et al 2003, Riordan et al 2011). Memantine has a modest effect in moderate to severe AD at therapeutic concentrations (Kornhuber & Quack 1995). However, it showed little effects in mild AD (Schneider et al 2011). Nevertheless, administration of memantine has been reported to reduce A β generation in neuroblastoma cells, primary cortical cells and in APP and mutant PS transgenic animals (Alley et al 2010). AD like neuropathology was reduced in transgenic mice treated with memantine (Martinez-Coria et al 2010). Thus, pharmacologic manipulation of the NMDA receptors alone, or alternatively in combination with cholinergic pathway modification, may be a supplemental treatment to relieve AD symptoms to a small extent (Doraiswamy 2003, Prvulovic et al 2010).

1.6.2 Prospective therapeutic strategies to delay disease progression

In recent years, several other therapeutic strategies have been proposed for the treatment of AD.

1.6.2.1 A β - based approaches

A β is thought to play a central role in AD pathogenesis. Hence targeting A β production, aggregation and removal may be of therapeutic benefit. To inhibit A β production, drugs that target APP processing by inhibition of secretase activities are

being developed. Studies have also reported that up-regulation of α -secretase in APP processing decreases A β yield and improves cognitive behaviour in transgenic mice, implying that this may be a viable treatment for AD (Caccamo et al 2006, Lin et al 1999). In addition, APP undergoes γ -secretase cleavage to release A β , thus selectively inhibiting γ -secretase activity is also considered to be a viable therapeutic strategy. The applied dose of γ -secretase inhibitor would have to be carefully controlled to avoid blocking important signaling pathways, such as those downstream of Notch (Bergmans & De Strooper 2010). BACE1 (β -secretase), the other enzyme that is involved in A β generation has multiple substrates in vivo (Hu et al 2006, Kim et al 2007a) but fewer than γ -secretase. Thus, modification of BACE1 activities without interfering with important physiological signaling may be a more promising potential therapeutic strategy for AD.

Inhibition of A β aggregation has been proposed to be a prospective viable AD treatment, and indeed cognitive benefits have been reported both in animals and in humans who were given inhibitors of A β aggregation (Adlard et al 2008, Lannfelt et al 2008, Olcese et al 2009). However, one must be careful when considering AD by inhibition of A β aggregation because the exact A β aggregation species causing A β mediated neurotoxicity remains unknown. Blocking aggregation of A β could end up being more detrimental if small aggregates are more neurotoxic, or if aggregation inhibiting aggregation increases the formation of longer aggregates.

Inadequate removal of A β from the brain may contribute to amyloid accumulation in the CNS, thus therapeutic strategies aimed at increasing A β removal and clearance are being investigated. They may be a more promising approach given that both β - and

γ -secretase inhibition may have off-target effects. For example, APP transgenic mouse studies indicate that increased astrocytic lysosome biogenesis may promote uptake, trafficking and degradation of A β , which suggests that activation of astrocytic lysosomes may facilitate A β removal and attenuate amyloid induced AD pathogenesis (Xiao et al 2014). In addition, modulation of neprilysin expression, the most potent A β -degrading enzyme (Shirotani et al 2001) may provide an alternative opportunity for intervention (Nalivaeva et al 2012). Neprilysin may act on a wide range of peptide substrates with biological functions. Nevertheless, this problem may be overcome by modifying the neprilysin active site in such a way as to make it more specific to for A β degradation (Webster et al 2014). A variety of proteins such as apolipoprotein E and J have been described that interact with A β and regulate its ability to cross the BBB (Calero et al 2012). A β may be degraded in the circulatory system after transport into the blood by either A β -degrading enzymes or by immune response cells, such as monocytes that have been shown to clear vascular A β (Michaud et al 2013).

1.6.2.1.1 A β immunotherapy

A β immunotherapy is a potential AD treatment that has been advanced recently because of its ability to decrease brain A β accumulation (Frenkel et al 2000). A β immunotherapy can be achieved by active immunization, which is performed by injecting AD subjects with synthetic intact synthetic A β or synthetic A β fragments bound to a carrier protein to mediate an immune response that generate antibodies against A β (Bard et al 2000). The antibody bound A β may be removed via Fc receptor-mediated clearance by microglia (Bard et al 2000). A β immunization results in reduced CNS A β load and less neuritic dystrophy, as well as improved cognition

compared to non-treated controls (Bard et al 2000, Masliah et al 2005, Schenk et al 1999, Serrano-Pozo et al 2010). However, active immunotherapy of AD patients was terminated in phase II clinical trials due to an incidence of meningoencephalitis which may have been caused by T cell-induced inflammatory responses (Liu et al 2009, Orgogozo et al 2003).

Passive immunization is another A β immunotherapy that is currently being tested in clinical trials. In passive immunization, the antibodies directly against A β are administered to the patients. Using this approach, the antibody may interact with plaques thereby inducing a scavenging response in microglia (Bohrmann et al 2012). Nevertheless, clinical trials using anti-A β monoclonal antibodies (bapineuzumab and ponezumab) were discontinued due to the lack of cognitive improvement and to adverse side effects (vasogenic cerebral oedema) (Blennow et al 2012, Burstein et al 2013, Landen et al 2013, Salloway et al 2014). However, a humanized antibody, crenzumab, with the decreased Fc receptor affinity, was safely given at a high dose to patients with a good safety outcome in a phase I trials that has seen the drug moved to phase II trials (Adolfsson et al 2012). Another humanized monoclonal antibody, solanezumab, that recognizes the middle region of A β and binds soluble monomeric forms of A β , shows positive effects in mild AD (Lannfelt et al 2014). Nevertheless, phase III trials of solanezumab was failed in mild to moderate AD (Doody et al 2014). This may be because this treatment was administered too late in the course of AD (Laske 2014).

Taken together, these results indicate that immunotherapy for AD treatment is perhaps one of the most promising areas for future investigation. However, a major challenge for both active and passive immunization may be the delivery of synthetic

A β antibodies across the BBB into the CNS (Spencer & Masliah 2014). Besides, A β immunotherapy, especially active immunization in particular has been found to produce persistent autoimmune side effects in an AD mice model. This was manifested by microglial infiltration, which not only activated microglia to engulf A β but also importantly led to disruption and degeneration of local tissues (Liu et al 2009). Moreover, it has not been easy to identify any clinical benefits in A β immunotherapy in phase II – III clinical trials (Blennow et al 2014). Therefore, active A β immunotherapy for AD treatment needs to be more intensively investigated. However, ethics considerations may prohibit further clinical trials. Currently passive immunization strategies demonstrate the feasibility of A β clearance, however, more studies will be needed to determine dosage and the stage at which antibodies should be delivered.

1.6.2.2 Stem cell therapy for AD

During the late stage of AD, reduction or elimination of A β may not be adequate to overcome the functional loss of neurons and synapses, nor allow for rescue of neuronal dysfunction caused by AD progression (Spencer & Masliah 2014). Therefore, stem cell based therapies have been proposed as a way to repair and replace the deficit or loss of neurons. Stem cell therapeutic strategies could employ two approaches: one is to activate endogenous stem cells, and the other is to transplant regenerated cells or tissues into the damaged or injured sites.

Stem cell transplantation therapy for AD is challenging because many different types of neurons and neurotransmitter systems are affected in AD. In order to incorporate into the complex brain circuit system, stem cells have to migrate into multiple areas of

the brain, differentiate into different types of neuronal cells, and to establish afferent connectivities of physiological relevance with appropriate targets. For this reason, therapeutic strategies based upon cell replacement may not succeed for diffuse neural disorders like AD (Chen & Blurton-Jones 2012). Nonetheless, recent studies indicate that neural stem cell transplantation can improve cognition in AD transgenic mice (Blurton-Jones et al 2009, Yamasaki et al 2007). Thus stem cell therapy has potential for development as a treatment for AD.

Stem cell therapy can also be used to deliver growth factors to the brain. Production of several essential neurotrophins that may regulate synaptic strength and numbers in an activity-dependent manner can be decreased significantly in the early stages of AD, possibly contributing to synapse loss (Arancio & Chao 2007). Neurotrophins, have been reported to improve cognition in patients and they are produced abundantly in NSPCs (Blurton-Jones et al 2009, Kamei et al 2007, Sun et al 2003). Delivery of NSPCs into the AD brain may increase neurotrophin secretion and thereby exert a positive effect on cognition (Tuszynski et al 2005). However, it still unclear whether functional recovery is driven by stem-cell mediated neurotrophin delivery, as lesions or injuries also stimulate production of endogenous neurotrophins (Blurton-Jones et al 2009, Martinez-Serrano & Bjorklund 1996).

Chronic inflammation may play a crucial role in AD neurodegeneration (Akiyama et al 2000). NSPCs can produce anti-inflammatory factors such as interleukin-10 and prostaglandin E2 (Ylostalo et al 2012, Zhou et al 2011a), or can attenuate microgliosis and release of pro-inflammatory cytokine tumour necrosis factor α (TNF α) (Ryu et al 2009). Delivery of anti-inflammatory NSPCs was found to reduce neuroinflammation and result in improved cognition and in pathology in AD mice (Seo et al 2011).

Evidence showed that anti-inflammatory drugs lower AD risk (Andersen et al 1995, Côté et al 2012), although it may be too late for patients displaying AD symptoms (Aisen 2002). Nevertheless, suppression of the immune system may be a viable therapeutic target for AD, as NSPC transplantation may possibly relieve AD symptoms by decreasing inflammation (Bentham et al 2008, Green et al 2009).

NSPCs may be a promising way to deliver therapeutic proteins to the damaged areas of the brain because NSPCs can migrate throughout brain tissue to the injured and inflamed areas (Muller et al 2006). In addition, some stem cells can stimulate microglia to produce the A β degrading enzyme, neprilysin (Kim et al 2012). However, it is still unknown whether survival and differentiation of transplanted stem cells can be affected by AD-related pathology, and whether transplanted cells are influenced by endogenous NSPCs (Chen & Blurton-Jones 2012).

1.7 Biology of neural stem and progenitor cells (NSPCs)

Stem cells, either derived from embryonic or adult CNS, maintain an ability to undergo long-term self-renewal and to differentiation into more highly specialized brain cells. Therefore, stem cells are theoretically considered to be an inexhaustible supply of many cell types for regenerative medicine. Neurological diseases are usually associated with neuronal damage or loss, thus use of stem cells to replace destroyed neural is a potential supplementary therapy for neurological diseases including AD. Moreover, all major types of cells in the CNS are derived from stem cells, thus investigation of stem cell biology may provide new insights into understanding the pathogenesis of neurological disorders.

1.7.1 Embryonic stem cells (ESCs)

Mammalian embryonic stem cells (ESCs) develop from the inner cell mass of the primitive implanted embryo. They are pluripotent and capable of generating the three primary germ layers, ectoderm, mesoderm and endoderm, which will give rise to all types of cells in tissues later in embryogenesis including neural stem cells (Rippon & Bishop 2004). Neural differentiation occurs at an early stage during embryonic development, soon after the specification of the germ layer. Cells of the ectoderm, the outer embryonic germ layer, have the capacity to form both the CNS and the epidermis (Li et al 2013a), while the cells in the margin of ectoderm develop into peripheral nervous system (Marshak et al 2001). The ectoderm is divided into three sets of cells. Internal ectodermal cells form a closed neural tube with a central canal that subsequently develops into the CNS (brain and the spinal cord) (Kalyani et al 1997).

The neural tube is comprised of a single layer of proliferating and morphologically homogenous cells called neuroepithelial cells (Kalyani et al 1997). Neuroepithelial cells can divide symmetrically to generate two identical daughter cells. Alternatively, they can switch to undergo various asymmetrical divisions that produce two distinct daughter cells - a self-renewing neural stem cell and a differentiating neuroblast (Gotz & Huttner 2005). During their transition to neural tissues such as neocortex, early neuroepithelial cells develop into radial glia cells that are mitotically active throughout neurogenesis and undergo both symmetric and asymmetrical cell division (Mission et al 1991). The radial glia cells processes radial morphology with glial characteristics in contrast to epithelial cells (Hartfuss et al 2001, Malatesta et al 2000) and are not only capable of self-renewal, but also participate in the generation of

neurons, astrocytes, oligodendrocytes and ependymocytes during embryonic development (Malatesta et al 2003, Spassky et al 2005). Radial glial cells can serve as a neuronal progenitors in all regions of the CNS during development (Anthony et al 2004) and are present until the end of embryonic neurogenesis and neuronal migration (Noctor et al 2004). The remainder will be transformed into astrocytic adult neural stem/progenitor cells (NSPCs) postnatally residing in the subventricular zone (SVZ) (Merkle et al 2004). However, no matter whether the cells are radial glial or neuroepithelial cells, they are all embryonic neural stem cells.

In the developing CNS, neural stem cell (NSC) proliferation and differentiation are regulated by a number of growth factors. NSC proliferation occurs in response to both epidermal growth factor (EGF) and fibroblast growth factor (FGF) (Tropepe et al 1999). In addition, insulin and insulin – like growth factors (IGFs) are suggested to play a role in embryonic NSC development as insulin has been found to promote NSC proliferation and survival (Desai et al 2011, Freund et al 2008) while IGF-1 immunoreactivity can be detected in the developing brain (Garcia-Segura et al 1991). IGF-1 stimulates NSC proliferation in a way that is distinct from that of EGF and FGF (Arsenijevic et al 2001). Furthermore, IGF - 2 is involved in the modification of cell proliferation and survival in early embryonic stages (Burns & Hassan 2001), and IGF-2 mutant mice display defects in brain development (Baker et al 1993). In addition, brain-derived neurotrophic factor (BDNF) stimulates NSC proliferation (Islam et al 2009) and can promote early radial glial cell differentiation in the developing cerebral cortex by inducing bone morphogenetic protein (BMP) expression in embryonic neurons (Ortega & Alcantara 2010).

1.7.2 Adult neural stem/progenitor cell (NSPCs) and neurogenesis

Neurogenesis, the birth of new neurons, is a process that produces functionally integrated neurons from stem cells or progenitors. This process occurs mostly in the mammalian CNS during embryogenesis, but it also occurs in post-natal stages and throughout adulthood in mammals (Altman & Das 1965, Kaplan & Hinds 1977). A variety of studies have revealed that neurons are born, and neural stem/progenitor cell (NSPC) proliferate, during adult life, but only in selected areas of the brain (Alvarez-Buylla & Garcia-Verdugo 2002, Eriksson et al 1998, Reynolds & Weiss 1992, Temple 1989, Temple & Alvarez-Buylla 1999). However, apart from the two classic areas of neurogenesis, the ventricular-subventricular zone (V-SVZ) lining in the lateral ventricle (Alvarez-Buylla & Garcia-Verdugo 2002, Bedard & Parent 2004, Fuentealba et al 2012, Lois & Alvarez-Buylla 1993) and the subgranular zone of the dentate gyrus (DG-SGZ) in the hippocampus (Eriksson et al 1998), other major neurogenic regions may exist (Bordiuk et al 2014) in the adult mouse brain.

Adult neurogenesis in both the V-SVZ and the SGZ is a process that is tightly controlled and regulated. It involves in NSPC maintenance, activation and proliferation, as well as differentiation and migration of the intermediate progenitors to their final destination. Newborn neurons subsequently mature and integrate into the local neuronal network (Hsieh 2012, Taylor et al 2013). The V-SVZ and SGZ in adult brain contain multipotent NSPCs that are capable of self-renewal through cell division and differentiation into neurons and glia (Palmer et al 1999, Reynolds & Weiss 1992, Richards et al 1992). Thus, multipotent NSPCs may be a source of potential precursors useful for transplantation to damaged neural tissues, and thus for the treatment of neurological diseases. Furthermore, alteration of neurogenesis in

neurological injury or disease has been reported (Curtis et al 2003, Gao et al 2009, Hoglinger et al 2004, Jin et al 2004c, Miles & Kernie 2008, Parent et al 1997, Yu et al 2009). Therefore studying adult NSPC proliferation and differentiation may be helpful for understanding how endogenous adult-born neurons may be used for brain repair and restoration, and whether the manipulation of endogenous neurogenesis is feasible for the treatment of neurological disorders.

1.7.2.1 Adult neurogenesis in V-SVZ

Radial glia cells serve as progenitors for many neurons of CNS during embryonic development (Anthony et al 2004). However, they also give rise to adult NSPCs located in the V-SVZ of the lateral ventricle (Merkle et al 2004). The NSPCs in the V-SVZ are located adjacent to the ependymal cell layer, which lines the lateral ventricle separating the ventricular cavity from the V-SVZ (Doetsch et al 1999). In addition, NSPCs also interact with basal lamina projection from the local vasculature (Mirzadeh et al 2008).

Ependymal cells remain quiescent and have no NSPC properties (Doetsch et al 1999). NSPCs contain type B cells expressing astrocyte markers (Doetsch et al 1997, Platel et al 2009) including glial fibrillary acid protein (GFAP) (Bignami & Dahl 1974) and glutamate aspartate transporter (GLAST) (Lehre et al 1995). Expression of GFAP in type B cells is consistent with the idea that radial glia transformed into astrocytic NSPCs in V-SVZ at the end of histogenesis (Merkle et al 2004). Type B cells are relatively quiescent, but they also can exist in an activated state according to recent studies (Codega et al 2014, Mich et al 2014). Quiescent type B cells do not express the intermediate filament protein, nestin (Lendahl et al 1990), which has been thought

of as a NSPC marker (Lim & Alvarez-Buylla 2014). Nevertheless, nestin is expressed in activated type B cells that generate actively proliferating C cells, which serve as neurogenic intermediate progenitor cells (nIPC) or transiting amplified progenitors in the V-SVZ of adult brain (Doetsch et al 1999). Type C cells divide symmetrically or asymmetrically and are able to generate neuroblasts (type A cells) that migrate to the olfactory bulb through (RMS) (Fig. 1.3) (Belluzzi et al 2003, Lois & Alvarez-Buylla 1994). Type C and A cells are also labelled by specific molecular markers. Despite expression of the immature neuronal marker doublecortin (DCX), type A cells are positively stained by anti-PSA-NCAM (Rousselot et al 1995, Rousselot & Nottebohm 1995) and anti-TuJ1 antibodies (Doetsch et al 1997). PSA-NCAM is a protein expressed by migrating neuroblasts (Doetsch & Alvarez-Buylla 1996, Rousselot et al 1995) while the TuJ1 antibody labels β III tubulin in young neurons (Easter et al 1993, Moody et al 1996). However, ependymal cells, type B and type C cells do not express PSA-NCAM and β III tubulin (Doetsch et al 1997), whereas type A and C cells strongly express GFAP. Both type A and type C cell types are immunoreactive with nestin antibodies (Doetsch et al 1997).

Apart from producing type A cells, type B and C cells may contribute to the generation of oligodendrocytes (Menn et al 2006). NG2-expressing cells that are known as oligodendrocyte progenitors (OPCs) show a type C cell - like immunophenotype during early postnatal and adult stages in the V-SVZ (Aguirre et al 2004). In addition, a small sub-population of type C cells in the V-SVZ express oligodendrocyte lineage transcription factor 2 (Olig2) (Menn et al 2006) that guides astrocyte and oligodendrocyte formation in the V-SVZ (Marshall et al 2005). Furthermore, Olig2 - expressing type B cells can give rise to a small number of non-




Figure 1.3 Schematic illustration of adult neurogenesis in the adult ventricle - subventricular zone (V-SVZ). (1) The V-SVZ is located in the wall of the lateral ventricles. (2) The V - SVZ contains ependymal cells (ECs, white), comparatively quiescent type B cells (blue), transit-amplifying type C cells (red) and neuroblasts type A cells (green). (3) In the V-SVZ, GFAP-positive astrocytes (type B cells) act as neural stem/progenitor cells that are able to generate nestin positive transit-amplifying cells (type C cells) which differentiate into neuroblasts (type A cells). Neuroblasts generated in the V-SVZ migrate towards the olfactory bulb (OB) along the rostral migration stream (RMS).

myelinating NG2 expressing cells and myelinating mature oligodendrocytes (Menn et al 2006). Moreover, enhanced generation of new oligodendrocytes from NSPCs in SVZ migrating to injury sites can be detected in vivo after demyelinating lesions of the corpus callosum (Picard-Riera et al 2002). Therefore, the NSPCs in the SVZ may generate new oligodendrocytes under both normal and injury conditions in the adult brain (Menn et al 2006).

Neuroblasts (type A cells) originating from NSPCs in the SVZ migrate through the RMS for a distance up to 5 millimeters in the adult mouse brain (Doetsch & Alvarez-Buylla 1996). Neuroblasts undergo tangential directional migration to the olfactory bulb (OB) via a chain migratory manner to form elongated cell aggregates that are ensheathed by astrocytes (type B cells) (Lois et al., 1996). This directional chain migration of neuroblasts along the RMS is regulated by a number of factors (Belvindrah et al 2007, Cao et al 2013, O'Leary et al 2014, Platel et al 2009, Young et al 2010). For instance, neuroblasts survival and migration are controlled by SVZ astrocytes via glutamate signaling through NMDARs (Platel et al 2010). In the RMS, PSA-NCAM expressing cells (type A cells) are more abundant than GFAP-expressing type B cells, but no type C cells could be detected (Doetsch et al 1997). Most neuroblasts (type A cells) migrate through the RMS into the OB where they differentiate into interneurons (Lois et al 1996, Whitman & Greer 2007a). Beside the RMS, neuroblasts also migrate to populate the cortex and subcortex regions in the first few postnatal weeks (De Marchis et al 2004, Sanai et al 2011). In addition, in vivo studies suggest that glia migration may not follow the classical RMS route (Menn et al 2006, Picard-Riera et al 2002, Seri et al 2006). Moreover, SVZ - derived neuroblasts migrate toward lesion sites where they differentiate into mature neurons

(Kernie & Parent 2010, Sundholm-Peters et al 2005, Yamashita et al 2006). This highlights the fact that simulating neurogenesis in the SVZ may be a potential therapeutic strategy for neurological disease.

Upon reaching the center of the OB, neuroblasts detach from the chain, and turn radically and move toward the granular cell layer, and then to the periglomerular cell layer (Lois et al 1996). Upon arrival at the destination, neuroblasts start to differentiate into interneurons. Specification of OB interneurons has genetic (Kohwi et al 2005, Saino-Saito et al 2007, Waclaw et al 2006) spatial and temporal (Lledo et al 2006) contributory factors. NSPCs from the V-SVZ give rise to two main types of interneurons in the OB. One type reaches the granular cell layer where the cells differentiate into granular neurons, whereas the other moves into the periglomerular layer and develops into periglomerular neurons (Petreanu & Alvarez-Buylla 2002, Yang 2008). To integrate and survive, the newly generated neurons receive either GABAergic or glutamatergic synaptic inputs from surrounding neurons and axonal terminals (Eyre et al 2008, Laaris et al 2007).

Granule cells (GCs) comprise the majority of the OB. Most of the GCs in the OB are produced postnatally and are added constantly in adulthood (Bayer 1983, Kaplan & Hinds 1977). Petreanu & Alvaren-Buylla (2002) first described the maturation of GCs generated from adult. After arriving at the granule cell layer, newborn axonless GCs soon project basal and apical dendrites to form synaptic connections and functionally integrate with existing OB circuitry (Belluzzi et al 2003, Petreanu & Alvarez-Buylla 2002). However, the survival selection and synaptic development of adult generated GCs are determined by the level of activity they receive (Balu et al 2007, Whitman &

Greer 2007b) e.g. GABAergic (Eyre et al 2008, Pallotto et al 2012), glutamatergic (Laaris et al 2007), noradrenergic (Veyrac et al 2005) and cholinergic (Devore & Linster 2012, Kaneko et al 2006, Pressler et al 2007) stimulation occurring in critical time windows (Ge et al 2007, Kelsch et al 2009) when neurons have a high degree of plasticity (Petreanu & Alvarez-Buylla 2002, Yamaguchi & Mori 2005). Almost 50% of the GCs die after one month, and the rest are durably integrated into the OB (Petreanu & Alvarez-Buylla 2002, Winner et al 2002). Only less than 10% of newborn GCs survive more than 21 months. Nevertheless, the dead or older cells will be replaced by newborn GCs (Kaplan et al 1985).

Compared to GCs, newborn interneurons differentiate into periglomerular cells (PGCs) more slowly, with a delay of nearly one month. PGCs receive either GABAergic or glutamatergic input from olfactory sensory axons and from the apical dendrites of principal neurons in the OB (Grubb et al 2008). Survival and synaptogenesis of PGCs also process in a sensory activity - dependent manner (Grubb et al 2008, Mandaïron et al 2006). PGCs generally have axons, but axonless types may also exist (Kosaka & Kosaka 2011). Three subtypes of adult - born PGCs have been described based upon their immunoreactivity to tyrosine hydroxylase (TH), an enzyme required for dopamine synthesis, to the calcium - binding proteins calbindin (CalB) and calretinin (Parrish-Aungst et al 2007). However, PGCs can also be grouped into two types, depending upon whether they do or do not receive synapses from olfactory nerves (Kosaka et al 2001, Kosaka & Kosaka 2011). Compared to GCs, the newly generated PGCs display a similar decline in cell numbers during maturation, with the remaining PGCs survive up to 19 months (Winner et al 2002).

Although NSC proliferation and differentiation occur persistently in the V-SVZ-OB region (Lois & Alvarez-Buylla 1993, Morshead et al 1994), the volume of the OB does not seem to increase (Pomeroy et al 1990). Apoptosis could be detected in the layer into which the newly produced neurons flow. This not only maintains the OB structure but also essentially contributes to neural plasticity of olfactory information processing (Fiske & Brunjes 2001, Imayoshi et al 2008, Lepousez et al 2013, Sakamoto et al 2014).

Adult neurogenesis in the V-SVZ-OB system is affected and regulated by a number of factors. Type B cell bodies extend short processes through the ependymal layer via apical endings that directly contact cerebrospinal fluid (CSF). The CSF contains multiple soluble factors (eg neurotrophins, growth factors) that may regulate NSPC maintenance and behaviour (Mirzadeh et al 2008, Shen et al 2008). Several extrinsic signals including bone morphogenetic proteins BMPs, sonic hedgehog (Shh), and Wnt related integration site (Wnt) proteins are present in CSF and modulate neurogenesis in V-SVZ (Huang et al 2010, Lehtinen et al 2011). Ependymal cells growing a large number of long motile cilia into the lateral ventricle, and recent studies report that primary cilia are crucial for Shh signal transduction (Huangfu & Anderson 2005, Lehtinen et al 2011). In addition, blood vessels in the V-SVZ zone contain a substantial extracellular matrix (ECM) that is closely associated with type B and C cells (Shen et al 2008, Tavazoie et al 2008), and endothelial derived factors may also play an important role in NSPC proliferation and neuroblast production (Katsimpardi et al 2014, Shen et al 2004). Furthermore, V-SVZ neurogenesis is also affected by hormones (Lau et al 2007, Lopez-Juarez et al 2012) and different types of neurotransmitters especially at stages of integration and maturation, such as GABA

(Alfonso et al 2012, Liu et al 2005), dopamine (Kim et al 2010, O'Keefe et al 2009) and acetylcholine (Paez-Gonzalez et al 2014). Additionally, multiple transcriptional factors participate in the modulation of NSPC commitment and fate specification. SOX2, a sequence - specific DNA - binding transcription factor exerts effects in many essential processes of adult NSPC (Sarkar & Hochedlinger 2013, Shimosaki et al 2012). Sox 2 cooperates with TLX transcription factors, and thereby mediate type B cell proliferation (Ellis et al 2004, Shimosaki et al 2012). Expression of basic helix - loop - helix (bHLH) transcription factor-neurogenin 2 (Ngn2), found in some type C cells, induces generation of calretinin – positive OB neurons (Roybon et al 2009b). Interestingly, microglia are suggested to promote generation of neurons and oligodendrocytes in V-SVZ by the release of cytokines (Gonzalez-Perez et al 2010, Shigemoto-Mogami et al 2014).

1.7.2.2 Adult neurogenesis in subgranular zone of dentate gyrus (DG-SGZ)

The subgranular zone (SGZ) of the hippocampus is another major region that generates new neurons in adult mammalian brain (Altman & Das 1965, Eriksson et al 1998, Kaplan & Hinds 1977). The SGZ zone, where hippocampal neurons are born, is located at the interface of the dentate gyrus granular cell layer and the hippocampal hilus (Seri & Alvarez-Buylla 2002). Two types of dividing NSPCs, radial astrocytes and small cells with basophilic nuclei that stained darkly with hematoxylin, have been found in the SGZ (Altman & Das 1965, Cameron et al 1993, Kaplan & Hinds 1977, Palmer et al 2000). Classic anatomical studies suggest that adult radial astrocytes in the SGZ are derived from radial glial cells in the embryo (Altman & Bayer 1990, Eckenhoff & Rakic 1984).

Proliferating radial astrocytes and other precursors generate intermediate precursors, which in turn give rise to neuroblasts in the SGZ. The neuroblasts are generated from immature neurons that migrate to the granular cell layer and continuously differentiate into granule cells (Kempermann et al 2003, Kuhn et al 1996, Seri & Alvarez-Buylla 2002). The putative NSPC in the SGZ is a type of radial glia - like cell that expresses both the astrocyte marker GFAP and the neural precursor marker nestin. This cell is known as the type I cell (Filippov et al 2003, Seri et al 2001). In addition to acting as a neuronal precursor, type I cells also share astrocytic electrophysiology properties, and retain astrocyte features, such as the ability to interact with the vasculature (Filippov et al 2003, Fukuda et al 2003). Factors coming from blood vessels may affect NSPC behavior in the DG-SGZ (Palmer et al 2000). Type I cells, like GFAP expressing - type B cells in the SVZ, represent only 5% of the nestin-expressing cell population in SGZ (Filippov et al 2003, Kronenberg et al 2003, Lugert et al 2010). The slowly dividing quiescent type I cells do not generate neurons directly, but give rise to dark staining small cells, termed type II cells (Filippov et al 2003, Fukuda et al 2003).

Type II cells, have short separate processes and dense - shaped nuclei. They do not express GFAP and are highly amplified cells (Filippov et al 2003, Kempermann et al 2004, Kronenberg et al 2003). Type II cells are nestin - positive and are defined as transit amplified neurogenic intermediate progenitors. Type II cells can be classified into two subtypes depending on whether they express DCX or not (Brown et al 2003). Transit amplified type II cells can potentially generate astrocytes or neurons in the SGZ (Encinas et al 2011, Kempermann et al 2004, Kronenberg et al 2003). Type III cells are nestin - negative and DCX - positive, but still have the capacity to

proliferate. They express more neuronal features than type II cells. Type III cells are also referred as neuroblasts (Seaberg & van der Kooy 2003, Weissman et al 2001). Up – regulation of the interneuron marker calretinin and neuronal marker NeuN as well as down - regulation of DCX occurs during differentiation and development of immature neurons into mature glutamatergic granule neurons (Hsieh 2012).

Unlike migration in SVZ, newborn neurons only migrate a short distance in the SGZ before arriving at the granular cell layer. The newly generated granule cells in the dentate gyrus exhibit similar electrophysiology characteristics to mature granule cells, but they continually undergo morphological and physiological alterations (Ge et al 2007, Toni et al 2007), as they grow dendrites to the molecular layer and extend axons to the CA3 region through the hilus (Zhao et al 2006). The newly produced granule cells integrate into the existing circuitry, and are initially activated by ambient GABA produced by surrounding interneurons (Bhattacharyya et al 2008, Ge et al 2006). They subsequently receive GABAergic synaptic inputs and then glutamatergic synaptic inputs (Esposito et al 2005, Ge et al 2006, Schmidt-Salzmann et al 2014), and finally they form synaptic contacts with hilar and CA3 neurons through mossy fiber output (Faulkner et al 2008, Toni et al 2008). However, adult - generated neurons display hyperexcitability and increased synaptic plasticity during development at specific stages compared to the fully mature granular neurons (Ge et al 2008, Schmidt-Hieber et al 2004). Importantly, newborn neurons have to receive GABAergic synaptic and glutamatergic inputs in order to survive (Laplagne et al 2006, Laplagne et al 2007).

Adult neurogenesis in the DG-SGZ is also regulated by multiple physiological, environmental and pathological factors including exercise (Kempermann et al 1998) and neurological disorders (Monje et al 2003, Parent et al 1997). Micro-environmental signalling from cellular compartments such as vascular cells, glia cells as well as granular neurons (Ma et al 2005, Palmer et al 2000, Tavazoie et al 2008, Tozuka et al 2005), and from extracellular matrix (Wojcik et al 2009, Wojcik-Stanaszek et al 2011), exert effects on NSPC activity in the SGZ. In addition, growth factors (Cheng et al 2002), including neurotrophins (Colditz et al 2010), as well as cytokines (Cheng et al 2002) and hormones (Cameron et al 1998) also contribute to the regulation of adult neurogenesis in the DG-SGZ. The neurotransmitters, glutamate (Esposito et al 2005) and GABA (Ge et al 2006), acetylcholine are also suggested to directly modify migration, maturation, integration and survival of newborn neurons in the hippocampus (Cooper-Kuhn et al 2004, Vivar et al 2012). In addition, cell to cell signalling by Notch (Ables et al 2010), Shh (Ahn & Joyner 2005), BMP (Mira et al 2010) and Wnt (Lie et al 2005) potentially participate in hippocampal neurogenesis regulation. This may be achieved through a series of transcriptional factors, for example the transcriptional factor Sox2 (Lefebvre et al 2007) is involved in NSPC self renewal, and requires active Notch signalling to maintain expression in NSPCs (Ables et al 2010, Ehm et al 2010, Imayoshi et al 2010). Furthermore, a number of transcriptional factors are also reported to control neurogenesis at different stages (Hodge et al 2008, Lugert et al 2010, Roybon et al 2010). For instance, the bHLH transcriptional factor Hes5 was found to colocalize with Sox2 and thereby collaborate to regulate NSPC maintenance (Lugert et al 2010), whereas another bHLH-transcriptional factor Ngn2, is up-regulated expressing only during transition from type I cell to neuroblasts, and is suggested to be crucial for specifying neuronal fate

during hippocampal neurogenesis (Roybon et al 2009b). Microglia also play a role in the regulation of adult neurogenesis by phagocytosis of newborn neurons undergoing apoptosis (Sierra et al 2010).

1.7.2.3 Function of adult neurogenesis

The generation of new neurons from adult NSPCs in the DG-SGZ, V-SVZ and other neurogenic niches in the adult brain has implicated in learning (Dupret et al 2007) and memory (Akers et al 2014), mood regulation (Snyder & Cameron 2012, Snyder et al 2011) as well as in neural repair following brain dysfunction or damage (Lazarini et al 2014, Sun 2014). In this section, the role of neurogenesis in regulating these functions is reviewed.

Newly generated neurons are thought to participate in olfactory- and hippocampal-dependent learning and memory (Braun & Jessberger 2014, Moreno et al 2009). New neurons derived from V-SVZ can interconnect with olfactory circuits and contribute to circuit maintenance and olfaction (Cummings et al 2014). This integration triggers learned behaviour responses and thereby promotes long-term memory (Lepousez et al 2013, Moreno et al 2009, Scotto-Lomassese et al 2003, Sultan et al 2010). Ablation of newborn neurons in the OB specifically impairs odour associative long-term memory (Arruda-Carvalho et al 2011). Interestingly, either odour associative or non-associative odour learning has been proposed to promote survival of newly incorporated neurons in OB (Belnoue et al 2011, Lepousez et al 2014), which implies that V-SVZ olfactory neurogenesis may not only contribute to, but may also be regulated by learning and memory.

Newly generated neurons derived from adult hippocampal neurogenesis are also suggested to be involved in learning acquisition and long term - memory formation (Suarez-Pereira et al 2014). Disruption of adult neurogenesis in hippocampus by radiation treatment impairs contextual learning and memory formation (Hernandez-Rabaza et al 2009, Kee et al 2007, Stone et al 2011, Winocur et al 2006). However, other results suggest that the link between adult hippocampal neurogenesis and learning and memory may not be so clear (Groves et al 2013, Luu et al 2012, Saxe et al 2006). These inconsistencies may be caused by multiple variables such as incomplete block to neurogenesis, unexpected off-target effects and different tasks applied to test learning and memory abilities (Deng et al 2010, Hernandez-Rabaza et al 2009).

Adult neurogenesis is also suggested to influence mood, as stress-induced down-regulation of hippocampal neurogenesis has been postulated to contribute to depression (Lehmann et al 2013, Snyder et al 2011, Surget et al 2011). The hippocampus plays a role in modulating stress via the hypothalamic pituitary adrenal (HPA) axis by controlling the release of cortisol from the adrenal gland (Herman et al 1989, Mizoguchi et al 2003). Several studies show that reduced hippocampal neurogenesis occurs under conditions of increased cortisol levels or chronic stress (Gould et al 1998, Lagace et al 2010, Lehmann et al 2013, Murray et al 2008). Disrupted hippocampal neurogenesis also causes higher levels of anxiety as well as a longer lasting elevation in blood cortisol levels. This suggests that newly generated neurons may regulate the HPA axis via a feedback loop following stress, and that this may in turn regulate cortisol levels which affect neurogenesis (Snyder et al 2011, Surget et al 2011). In addition, several studies show increased numbers of adult

hippocampal NSPCs following chronic treatment with clinical antidepressants (Boldrini et al 2009, Duman 2004, Malberg et al 2000, Perera et al 2007), which supports the idea of a link between adult neurogenesis and depression. However, the relationship between neurogenesis and depressive behavior is also questioned by various studies showing that ablation of adult neurogenesis does not affect anxiety or depressive-like behaviors (Jayatissa et al 2009, Santarelli et al 2003, Wang et al 2008). The conflicting results from these groups may be attributed to inadequate inhibition of neurogenesis or different methodologies used to monitor anxiety or depression.

In addition the relief from depressive behaviour may not be due to enhanced neurogenesis, although exposure to antidepressant treatments has been reported to promote neurogenesis (Malberg et al 2000). For example, an antidepressant agent, fluoxetine was reported to stimulate NSPC proliferation (Malberg et al 2000). Nevertheless, fluoxetine can relieve depression by increasing synaptic serotonin availability through its action as a serotonin reuptake inhibitor (Rahmani et al 2013, Ranganathan et al 2001). Interestingly, serotonin was observed to promote granule cell proliferation (Brezun & Daszuta 2000, Rahmani et al 2013). Thus, inhibition of serotonin reuptake may increase NSPC proliferation. More importantly, no change has been reported in NSPC proliferation in subjects with depressive like symptoms (Reif et al 2006). Therefore, the potential role of adult neurogenesis in mood regulation including its role in clinical depression, needs further clarification.

In general, enhanced adult neurogenesis occurs in a number of pathological conditions such as during seizure (Parent et al 1997), schizophrenia (Reif et al 2006)

or traumatic brain injury (Gao et al 2009). However, increased adult neuron generation may not always contribute to symptom relief. Indeed, it may even make the pathological situation worse in some cases. For example, seizures induce increased number of dentate granule cells in the adult rat hippocampus, and these new granule cells are committed to abnormal network reorganization (Parent et al 1997). This may negatively affect local existing circuit in hippocampus. Nevertheless, some neural damage may benefit from enhanced adult neurogenesis. For instance, increasing neurogenesis via physical activity seems to improve behavioural deficits in a mouse model of schizophrenia (Wolf et al 2011). Therefore, the role of adult neurogenesis in generating functional neurons for repair and replacement of the damaged or dysfunctional cells needs to be clarified in specific neurological disorders.

1.7.2.4 Factors influencing NSPC proliferation and differentiation

To produce new functional neurons, astrocytes or oligodendrocytes, a NSPC must undergo proliferation, migration and differentiation and then integrate into existing cellular networks (Christie & Turnley 2012). The NSPC life cycle and biology in the adult brain under both normal or pathological conditions is tightly regulated by multiple factors including a variety of cell-signalling pathways (Huangfu & Anderson 2005, Lie et al 2005), growth factors (Cheng et al 2002, Teramoto et al 2003), cytokines and chemokines (Bajetto et al 2002, Turbic et al 2011), as well as extracellular matrix (ECM) molecules (Wojcik-Stanaszek et al 2011). In particular, proliferation and differentiation may be modified by these factors to promote cell production and the generation of specific neural cell types after CNS injury (Lu et al

2011, Schabitz et al 2007, Zhu et al 2011). In this section, several factors affecting cell proliferation and differentiation are reviewed.

1.7.2.4.1 Growth factors

Various growth factors including epidermal growth factor (EGF) (Ayuso-Sacido et al 2010), fibroblast growth factor (FGF) (Palmer et al 1999), nerve growth factor (NGF) (Scardigli et al 2014), brain derived neurotrophic factor (BDNF) (Ahmed et al 1995), glial cell line-derived neurotrophic factor (GDNF) (Boku et al 2013, Heuckeroth et al 1998), insulin-like growth factor 1 (IGF-1) (Arsenijevic et al 2001), and vascular endothelial growth factor (VEGF) (Calvo et al 2011) participate in the modulation of NSPC proliferation and differentiation. Increased expression of these growth factors has been found following ischemia and traumatic brain injury (Christie & Turnley 2012).

1.7.2.4.1.1 Epidermal growth factor (EGF)

EGF, a 5.5 kD peptide, is mitogen that is widely used to stimulate NSPC proliferation and to maintain NSPC growth in cell culture (Reynolds & Weiss 1992). EGF is a ligand that binds to the EGF receptor (EGFR) that is expressed by both type B and type C cells in the V-SVZ (Doetsch et al 2002) as well as by radial astrocytes in the SGZ (Jin et al 2002). Activation of EGFR tyrosine kinase activity promotes NSPC proliferation, differentiation, migration and survival (Aguirre et al 2010, Gonzalez-Perez & Quinones-Hinojosa 2010, Gonzalez-Perez et al 2009). Transit amplified NSPCs (type C cells) in V-SVZ express high levels of EGFR, and can be induced by EGF to form a multi-potent state capable of producing either neuronal or glial precursors (Doetsch et al 2002). The generation of EGF in the V-SVZ seems to be

essential for maintenance of the proliferative NSPC pool (Christie & Turnley 2012), while EGF infusion into the SGZ appears to promote NSPC proliferation in response to brain injury (Sun et al 2010). Interestingly, infusion of EGF into the uninjured mouse brain also increases NSPC number (Cooke et al 2011) but promotes significantly more glial differentiation than neuronal differentiation (Craig et al 1996, Kuhn et al 1997). However, EGF induces a different extent of NSPC glial differentiation and especially oligodendrocyte generation and remyelination upon brain damage (Aguirre et al 2007, Cantarella et al 2008, Gonzalez-Perez et al 2009), although there are reports that disagree with this opinion (Craig et al 1996, Gonzalez-Perez et al 2009, Kuhn et al 1997).

The precise cellular and molecular basis for the EGF effect on NSPC maintenance and differentiation is not clear (Christie & Turnley 2012, Nieto-Estevez et al 2013). However, EGF is suggested to promote NSPC proliferation and possibly to induce glial differentiation through activation of EGFR (Kuhn et al 1997, Lindberg et al 2012). Several other ligands of EGFR such as β -cellulin (Gomez-Gaviro et al 2012) and transforming growth factor alpha (TGF- α) (Fallon et al 2000, Tropepe et al 1997) have also been proposed to increase NSPC growth through activation of the EGFR (Schneider et al 2008). In addition, inhibition of EGFR activity is reported to promote neuronal differentiation in a spinal cord injury model (Ayuso-Sacido et al 2010, Ju et al 2012, Li et al 2013b). Therefore, EGF may bind to and activate EGFR associated signaling pathway to affect NSPC proliferation and differentiation.

1.7.2.4.1.2 Fibroblast growth factor-2 (FGF-2)

FGF-2 is another mitogen that is usually combined with EGF to stimulate NSPC proliferation in culture (Reynolds & Weiss 1992). FGF-2 has a molecular weight of 18 kDa. Nonetheless a 24 kDa isoform has also been reported (Powers et al 2000). FGF-2 is mainly produced and secreted by glial cells, particularly by mature astrocytes in the CNS (Reuss et al 1998). FGF-2 regulates NSPC activity in the neurogenic niche by acting on FGF receptors (FGFR) (Galvez-Contreras et al 2012). Generally, four types of FGFRs are expressed differently in different neurogenic locations in the adult brain, with FGFR-1 and 2 occurring in dividing V-SVZ NSPCs, and FGFR-3 only in non-proliferative progenitors in the V-SVZ (Frinchi et al 2008). Notably, FGFR-1 is likely only expressed by NSPC in the SVZ of lateral ventricle (Frinchi et al 2008).

Exogenous infusion of FGF-2 into the lateral ventricle stimulates NSPC proliferation in V-SVZ under normal physiological conditions (Kuhn et al 1997), while inhibition of the release of FGF-2 or pharmacological block of the FGF receptor (FGFR) does not affect NSPC neuronal differentiation. This suggests an important role for FGF-2 in mitotic activity of NSPC in the V-SVZ, but not in the induction of neuronal fate (Agasse et al 2007). FGF-2 knockout mice do not show a significant reduction in NSPC numbers in hippocampus, but display decreased hippocampal NSPC neuronal differentiation after brain injury (Werner et al 2011), which implies that the endogenous production of FGF-2, or activation of FGFR-1, is important for stimulation of NSPC growth and development in the hippocampus following brain damage (Werner et al 2011, Yoshimura et al 2001). However, brain insult promotes up-regulation and accumulation of FGF-2 in NSPCs, which efficiently migrate to the

injured cortex and produce an enhanced pool of immature neurons available for brain repair (Dayer et al 2007, Ganat et al 2002, Wagner et al 1999). Furthermore, FGF-2 supplement in cell culture medium seems to indirectly induce NSPCs to differentiate into astrocytes (Song & Ghosh 2004), and EGF-2 also appears to promote astrocyte activation after brain injury (Reuss et al 1998). Moreover, FGF2 regulates oligodendrocyte progenitors (OPCs) repair process in response to demyelination (Dayer et al 2007, Dehghan et al 2012, Frost et al 2003), as up-regulation of FGF2 ligands and FGFR expression are found in remyelinating lesions (Messersmith et al 2000). Several studies reported similar results i.e. that infusion of EGF in combination with FGF-2, generally improves neural repair through promotion of V-SVZ - derived NSPC proliferation and survival, and repopulation of neurons at the injured areas, leading to functional recovery (Nakatomi et al 2002, Oya et al 2008, Tureyen et al 2005, Winner et al 2008). However, these results do not agree with all published studies (Baldauf & Reymann 2005). Thus the function of FGF-2 in NSPC behaviour has not been fully elucidated. Nevertheless, FGF-2 gene transfer rescue of hippocampal functions in AD brain (Kiyota et al 2011) may be a therapeutic approach for neurodegenerative diseases via manipulation of NSPC growth factors.

1.7.2.4.1.3 Insulin and insulin-like growth factors (IGFs)

EGF and FGF-2 probably are the most potent mitogens for NSPC proliferation, but other growth factors like insulin are also found to stimulate cell proliferation (Rafalski & Brunet 2011), as can IGFs (IGF-1 and IGF-2) (Arsenijevic et al 2001). As a hormone that assists cell store glucose and fats as energy sources (Kadowaki et al 2003, Terauchi & Kadowaki 2002), insulin is also essential for brain function and maintenance (McNay 2007, Rafalski & Brunet 2011, Zhu et al 1990). While it is not

clear how insulin affects NSPC fate, withdrawal of insulin from the cell culture medium leads to autophagic death of NSPCs (Yu et al 2008), whereas high concentration of insulin induce neuronal differentiation of postnatal NSPCs (Han et al 2008). In general, the standard culture media for mouse, rodent and human NSPCs contain insulin and are required for NSPC survival in vitro (Rafalski & Brunet 2011, Yu et al 2008).

Insulin is produced from β cells of the pancreas and enters the brain across the blood brain barrier (BBB) in a receptor-mediated manner (Banks et al 1997a, Banks et al 1997b, Schwartz et al 1991). IGFs are synthesized and released by the liver into blood circulation. Like insulin, IGFs can be transported across the BBB (Reinhardt & Bondy 1994). Furthermore, IGF-1 is also produced in the adult brain, especially in regions of cells undergoing division such as the hippocampus, the V-SVZ and the OB (Bartlett et al 1991). This suggests that IGFs have an important role in NSPC proliferation (D'Ercole et al 1996).

Insulin and IGFs receptors are expressed in adult NSPC, and a number of studies have raised the possibility that insulin, IGF-1, and IGF-2 supply instructive signals to regulate the fate of NSPCs through activation of their corresponding receptors (Bracko et al 2012, Hsieh et al 2004, Supeno et al 2013). For example, insulin is suggested to promote proliferation and prolong survival of NSPCs by binding to insulin receptors which are expressed by NSPCs in culture (Erickson et al 2008). In addition, IGF-1 has been implicated in the control of long - term EGF/FGF2-responsive NSPC proliferation, and in the neuronal differentiation of NSPCs

(Kouroupi et al 2010, Supeno et al 2013, Zhang et al 2014). Furthermore, IGF-2 has also been proposed to regulate postnatal neurogenesis (Bracko et al 2012), as IGF-2 exerts its effects on stem cell differentiation through binding to insulin (Ziegler et al 2014).

1.7.2.4.2 Other factors affecting NSPC proliferation and differentiation

Apart from growth factors, NSPC proliferation or differentiation is regulated by a series of factors such as neurotrophins (Ahmed et al 1995, Scardigli et al 2014), morphogens (Ahn & Joyner 2005, Lie et al 2005, Lim et al 2000), cytokines and chemokines (Bajetto et al 2002, Barbero et al 2002) and transcription factors (Ahmed et al 2009). Notably, a few studies indicate that cysteine protease inhibitors may also participate in regulation of NSPC behavior (Hong et al 2002, Santos et al 2012). FGF induced NSPC proliferation was reported to require a cysteine protease inhibitor, CysC (Taupin et al 2000).

The role of neurotrophins is described briefly in this literature review because neurotrophins have also been reported to affect NSPC proliferation and differentiation. In the course of my studies, it was necessary to consider all potential factors that could mediate the effect of APP on NSPC proliferation and differentiation. Binding of nerve growth factors (NGF) to TrkA receptors and to a much lesser extent the binding of brain - derived neurotrophic factor (BDNF) to TrkB receptors have been suggested to exert effects on differentiation and proliferation of NSPCs (Ahmed et al 1995, Frielingsdorf et al 2007, Islam et al 2009, Scardigli et al 2014). However, the opposite effect of BDNF on NSPC behaviour has also been reported. BDNF did not stimulate neurogenesis in SVZ (Galvao et al 2008).

The morphogens, Wnt (Adachi et al 2007, Lie et al 2005), Shh (Ahn & Joyner 2005, Huangfu & Anderson 2005) and BMPs (Lim et al 2000) are also suggested to regulate NSPC proliferation and maintenance. For example, Wnt signalling increases NSPC proliferation in the neurogenic areas (Adachi et al 2007, Lie et al 2005). Shh, a soluble protein that is normally produced by astrocytes in the neurogenic niche of the adult brain is suggested to regulate NSPC proliferation both in vitro and vivo (Lai et al 2003, Ruiz i Altaba et al 2002). Finally, BMP signalling has been reported to suppresses neuronal and oligodendroglial differentiation of NSPC, but induce astroglialogenesis of NSPC during adult life (Lim et al 2000). Nevertheless, BMP inhibitor expression in the V-SVZ or SGZ can block the BMP signalling and thereby increases neurogenesis (Bonaguidi et al 2008, Lim et al 2000).

Chemokines also have been suggested to play a role in the regulation of NSPC proliferation, differentiation migration and survival (Bajetto et al 2002, Barbero et al 2002, Cartier et al 2005). Adult NSPCs can express chemokine receptors (CCRs) and chemokines that are basally produced in the CNS, especially in the neurogenic regions (Tran et al 2004, Turbic et al 2011). Specific chemokines have been shown to stimulate NSPC proliferation (Filippo et al 2013, Tran et al 2004, Turbic et al 2011). For instance, the pro-inflammatory cytokines interferon gamma ($IF\gamma$) and $TNF\alpha$ are major regulators of CCRs and chemokines in the CNS and in other tissues (Hiroi & Ohmori 2003, Suyama et al 2005). Although $IF\gamma$ and $TNF\alpha$ inhibit NSPC proliferation (Ben-Hur et al 2003, Iosif et al 2006), they can induce neuronal differentiation as well (Ben-Hur et al 2003, Kajiwarra et al 2005, Lum et al 2009, Wong et al 2004). There are additionally several other cytokines that are suggested to regulate NSPC proliferation and differentiation (Ernst & Jenkins 2004, Heinrich et al 2003, Kamimura et al 2003, Lan et al 2012, Turnley & Bartlett 2000).

A number of transcriptional factors (TFs) are suggested to play a role in maintaining the multipotent network of NSPCs based on loss of function or gain of function studies (Ahmed et al 2009, Favaro et al 2009, Kim et al 2007b, Lefebvre et al 2007, Lugert et al 2010, Marei & Ahmed 2013), although the role of transcriptional cascades in neuron production in the adult neurogenic regions is relatively unknown compared to that occurring in embryonic neurogenesis (Hsieh 2012). However, a series of TFs are known to be expressed in neurogenic niches. TFs include Sox2, which is found to be essential for NSPC proliferation and maintenance (Favaro et al 2009). Ascl1, a basic helix–loop–helix (bHLH) transcription factor that has been implicated in the induction of neuronal fate of NSPCs but not in the stimulation of glial differentiation (Kim et al 2011, Kim et al 2007b), and Hes5, a bHLH transcription factor that has been proposed to be a direct target of Notch signalling in regulation of cell growth (Ables et al 2011, Lathia et al 2008, Lugert et al 2010).

1.7.2.4.2.1 Cystatin C

As an endogenous cysteine protease inhibitor, cystatin C (CysC) (Bobek & Levine 1992) is ubiquitously produced and secreted by various tissues. It is found in most mammalian body fluids, and at particularly high concentrations in the CSF (Abrahamson et al 1986, Turk et al 2008). In the CNS, CysC is abundantly expressed (Hakansson et al 1996), and is produced by neurons, astrocytes and microglial cells (Miyake et al 1996, Palm et al 1995, Yasuhara et al 1993). CysC is a hydrophilic protein containing 120 amino - acid residues with two intramolecular disulfide bridges that are mainly involved in modulating the activity of cysteine proteases (Otto & Schirmeister 1997). CysC inhibits lysosomal cathepsins especially cathepsin B, H,

K, L (Abrahamson 1994), caspases (Thornberry 1997) as well as cytosolic calpains (Turk et al 1997). Cysteine proteases play important roles in various biological processes including protein degradation, cell proliferation, differentiation and apoptosis (Berdowska & Siewinski 2000, Chapman et al 1997, Grzelakowska-Sztabert 1998). Thus precise control of cysteine protease activities by inhibitors such as CysC is essential for normal cell function. CysC, like other cystatins, is an exosite binding inhibitor that binds close to the active site of the cysteine protease, but does not interact with the active site directly (Bode & Huber 2000). CysC is suggested to participate in a wide range of biological processes, such as the regulation of inflammatory responses (Kopitar-Jerala 2006, Magister & Kos 2013, Warfel et al 1987), cell proliferation and growth (Sun 1989, Tavera et al 1992), and neural repair (Suzuki et al 2014a, Watanabe et al 2014, Zhong et al 2013).

CysC null mice exhibit decreased basal levels of neurogenesis and impaired proliferation and migration of newly generated granule cells in the dentate gyrus following seizures (Pirttila et al 2005, Taupin et al 2000). This suggests that there is a possible relationship between CysC and adult neurogenesis (Kaur & Levy 2012). Dividing NSPCs in vitro and in vivo were both found to be immunoreactive for CysC antibody, while combined delivery of CysC and FGF-2 stimulated increased neurogenesis in the dentate gyrus (Taupin et al 2000). These results raise a possibility the CysC may act as autocrine factor for NSPC growth (Taupin et al 2000). Proteome analysis indicates that CysC is secreted into the conditioned medium by cells during NSPC proliferation (Dahl et al 2003, Taupin et al 2000). CysC purified from NSPC conditioned medium was shown to promote NSPC proliferation in vitro (Dahl et al 2004, Taupin et al 2000) although CysC released by ESCs into the conditioned

medium does not display a stimulatory effect on NSPC proliferation. Nevertheless, the concentration of exogenous CysC that was used in these experiments was 20 ng/ml which may not have been adequate to increase NSPC proliferation (Tham et al 2010). Moreover, CysC purified from adult NSPC was reported to induce ES cells to differentiate into NSPCs which retain the ability to self-renew and exhibit multipotency in vitro (Kato et al 2006). Furthermore, a higher proportion of β III tubulin (neuron marker), GFAP (astrocyte marker) and MBP (oligodendrocyte marker) positive cells was observed in ES-NSPC cultures treated with purified CysC compared to a control group (Kato et al 2006). Besides, CysC was shown to activate GFAP promoter and to be expressed in astrocyte progenitors during development (Kumada et al 2004). Therefore, CysC may contribute to both embryonic and adult NSPC fate and behaviour.

Up-regulation of CysC protein and mRNA levels in neurogenic regions of the dentate gyrus of the hippocampus were reported following brain injury and seizure (Aronica et al 2001, Hendriksen et al 2001, Lukasiuk et al 2002). Interestingly, increased CysC expression occurs at the time when neurogenesis is maximal (Kumada et al 2004, Nairismagi et al 2004, Parent et al 1997). As CysC is involved in neurogenesis during development, it may also be involved in neurogenesis after injury. Indeed, proliferation of adult NSPCs in SGZ was decreased in CysC knockout mice compared to litter-mate controls after brain injury (Pirttila et al 2005).

However, the mechanism underlying CysC - induced NSPC proliferation as well as ESC – NSPC differentiation is still uncertain (de Azevedo-Pereira et al 2011). CysC reportedly may regulate cell proliferation independent of its effect on cysteine

proteases (Sun 1989, Tavera et al 1992), possibly by acting as a growth factor to stimulate NSPC proliferation (Taupin et al 2000). However, cysteine proteases are also supposed to be an effector of cell proliferation and differentiation (Egberts et al 2004, Joyce et al 2004, Watkinson 1999, Yadaiah et al 2013b), which raises the possibility that regulation of cysteine protease activities by cysteine protease inhibitors may also affect NSPC proliferation and differentiation. Mouse ESCs have been shown to differentiate into more NSPCs by addition of cysteine protease inhibitors such as CysC and E-64 (de Azevedo-Pereira et al 2011), and higher β III tubulin – positive cells (neurons or neuronal progenitors) were found in ESC cultures which were treated with E-64 (de Azevedo-Pereira et al 2011). However, the mechanism of induced differentiation of mouse ESC into NSPC and neurons by inhibition of cysteine proteases may be distinct from CysC - induced neural differentiation (de Azevedo-Pereira et al 2011). Interestingly, CysC increases the number of NSPCs as well as GFAP-positive and nestin-positive cells from brain ESC culture. Nevertheless, the increased number of GFAP cells reportedly does not seem to be due to inhibition of cysteine proteases by CysC (Hasegawa et al 2007). Therefore, the basis of CysC induced proliferation and differentiation of NSPC needs further investigation.

1.7.2.4.2.2 Neurogenin 2

Neurogenin 2 (Ngn2) is a proneural factor that was originally found to improve neuronal differentiation in brain and spinal cord (Li et al 2012b, Ma et al 1996, Sommer et al 1996). Ngn2 is expression in both CNS and PNS during embryonic neural development (Bertrand et al 2002, Chapouton et al 2001, Parras et al 2002, Ross et al 2003, Sommer et al 1996). Ngn2 induces neuronal determination of NSPC

(Scardigli et al 2001, Simmons et al 2001, Thoma et al 2012), and has been suggested to participate in neuronal-type and morphological specification (Florio et al 2012, Hand et al 2005, Hand & Polleux 2011, Kele et al 2006, Ma et al 2008c, Park et al 2008a) as well as in the control neuron migration (Heng et al 2008). However, Ngn2 suppresses progenitor cell cycle progression during neural development and maturation while promoting neuronal differentiation (Florio et al 2012).

Ngn2 null mutants survive a few days postnatally (Andersson et al 2006), although the vast majority of Ngn2 null animals die around birth (Fode et al 1998, Fode et al 2000, Hand et al 2005). Nonetheless, in the absence of Ngn2, animals exhibit abnormalities including decreased cerebellum volume, a lobulation defect in the cerebral hemispheres and reduced numbers of NSPCs (Florio et al 2012). This suggests a crucial role of Ngn2 in embryonic neurogenesis. Ngn2 activity declines at a very late stage of embryonic development (Li et al 2012b). Nevertheless Ngn2 is also expressed by NSPCs in adult neurogenic niches (Ozen et al 2007, Roybon et al 2009b). This suggests that Ngn2 also plays an important role in adult neurogenesis (Galichet et al 2008, Ozen et al 2007, Roybon et al 2009b). Interestingly, Ngn2 also promotes the genesis of neurons from non-neural cell types such as fibroblasts (Liu et al 2013, Yan et al 2001a). Moreover, NSPCs transduced with Ngn2 displayed improved survival and differentiation after transplantation (Yi et al 2008) and endogenous NSPCs transfected with Ngn2 carrying retroviruses showed neuroblast-like morphology (Rogelius et al 2008). Therefore, investigation into function of Ngn2 in NSPC proliferation and differentiation may be helpful for understanding the mechanism of induction of neurons for stem cell therapies.

1.7.2.4.3 An in - vitro model for studying NSPCs: neurospheres

Embryonic stem cells (ESCs) (Reynolds & Weiss 1992) and neural stem cells as well as their progeny (NSPCs), may be derived from the CNS of animals of any age and induced to proliferate in response to growth factors (Reynolds & Weiss 1992, Vescovi et al 1993) or to differentiate when they are exposed to specific culture conditions (Helgason & Miller 2013, Reynolds & Weiss 1992) in vitro. NSPCs have the capacity for self-renewal through their selective response to EGF as well as to FGF in vitro. NSPC cultures can be established from the V-SVZ and SGZ-DG or other regions of the fetus or adult brain (Bordiuk et al 2014, Liu & Martin 2003, Lois & Alvarez-Buylla 1993, Palmer et al 1997, Reynolds & Weiss 1992, Richards et al 1992, Weiss et al 1996). The culture medium is usually serum free, but supplemented with B27 (containing retinoic acid and insulin) or N2 (containing insulin and transferrin) in order to select for expanding NSPCs (Gritti et al 1996, Reynolds & Weiss 1996, Vescovi et al 1993). NSPCs grow in vitro as spherical floating cluster known as neurospheres, but they can also be cultured as adherent cells (Palmer et al 1997, Potten & Loeffler 1990, Ray et al 1993, Weiss et al 1996). Upon removal of the growth factors, the neurospheres can generate three major types of CNS cells (Reynolds & Weiss 1992). Neurospheres can also be derived from ESCs (Louis & Reynolds 2005, Tropepe et al 2001, Ying et al 2002). A single neurosphere is generally thought to clonally expand from a single stem cell. Nonetheless the evidence indicates that both stem and progenitor cells can proliferate to form neurospheres (Reynolds & Rietze 2005).

The neurosphere culture system was the primarily in vitro NSPC culture system to illustrate the existence of a group of NSPCs in the adult brain with the properties of

self-renew and multipotency (Reynolds & Weiss 1992, Reynolds & Weiss 1996). The neurosphere culture model is a useful technique that allowed evaluation of NSPC proliferation, self-renewal and multipotency via testing on neurosphere forming or differentiation potential (Gritti et al 1996, Rietze et al 2001, Tropepe et al 1999, Uchida et al 2000). In addition, the neurosphere culture system allows the study of neural development and adult neurogenesis in vitro (Alexson et al 2006, Jensen & Parmar 2006), and especially to investigate intrinsic characteristics of NSPCs at different developmental stages (Falk et al 2002, Jensen & Parmar 2006, Klein et al 2005). In addition, this in vitro NSPC culture model has been used to study the differentiation potential of NSPCs upon exposure to specific extrinsic factors (Ben-Hur et al 2003, Jensen & Parmar 2006). Importantly, neurospheres in biomedical research can be applied to drug test screening, or to identify factors affecting NSPC fate. This is due to the possibility of culturing a pool of specialized neural cells (Galli 2013, Singec et al 2006), and the ability to efficiently manipulate intrinsic properties of NSPC by transduction (Falk et al 2002, Hack et al 2004, Heins et al 2002). Furthermore, neurosphere-derived cells have potential applications in stem cell therapy for neurological disorders such as neurodegenerative diseases because they may provide a sustainable cell source for neural regeneration (Gil-Perotin et al 2013).

1.8 Neurogenesis in neurodegenerative disease

A number of studies have reported that adult neurogenesis is affected by pathological situations including stroke (Greenberg 2007), seizure (Parent et al 1997), acute trauma (Gao et al 2009, Miles & Kernie 2008) and neurodegenerative disease (Jin et al 2004c, Winner et al 2011). Neurodegenerative diseases are a broad range of disorders that have the common features including the progressive structural and functional loss of

neurons and glial cells in the brain and spinal cord (Winner et al 2011). Chronic neurodegeneration may exert various effects on NSPC maintenance (Kandasamy et al 2010), proliferation (Hoglinger et al 2004, Jin et al 2004c), survival and functional integration (Winner et al 2011). Furthermore, altered or even impaired adult neurogenesis has been described in models of neurodegenerative diseases such as Alzheimer's disease (AD) (Jin et al 2004c, Mirochnic et al 2009, Rodriguez et al 2008), Parkinson's disease (PD) (Crews et al 2008, Nuber et al 2008, Winner et al 2004) and Huntington's disease (HD) (Kandasamy et al 2010, Kohl et al 2007). However, whether deficits in neurogenesis contribute to neurodegenerative diseases still remains unknown, although impaired olfaction and hippocampus associated cognitive and emotional deficits are commonly identified in variety of many neurodegenerative diseases (Winner et al 2011).

1.8.1 Neurogenesis in Alzheimer's disease

Adult neurogenesis was suggested to contribute to learning and memory (Deng et al 2010). The rate of neurogenesis decreases with age, and seems to be further affected by AD pathogenesis (Jin et al 2004c, Kuhn et al 1996). A number of studies have used AD transgenic model animals to investigate this effect (Jin et al 2004a, Perry et al 2012, Yu et al 2009), although the mechanism underlying the neurogenesis response to AD pathology is still unknown (Jin et al 2004a, Martinez-Canabal 2014, Perry et al 2012, Yu et al 2009). Altered adult neurogenesis has been reported in AD transgenic animal models in the neurogenic niches such as the V-SVZ and DG-SGZ depending on post-mortem analysis (Marlatt & Lucassen 2010). Post-mortem studies of neurogenesis found enhanced expression of immature neuronal markers in the brain of AD patients (Jin et al 2004c). Furthermore, increased neurogenesis has been

identified in several AD transgenic mouse models (Jin et al 2004a, Lopez-Toledano & Shelanski 2007, Yu et al 2009). Nonetheless, some other studies challenge these observations and suggest that there are no changes or even decreased NSPC proliferation, in patients with AD (Boekhoorn et al 2006, Donovan et al 2006, Moon et al 2014, Rodriguez et al 2008).

However, the conflicting results may be due to combination of factors. Adult neurogenesis may change in response to AD pathology (Perry et al 2012, Yu et al 2009) such as A β deposition and synaptic loss (Kanemoto et al 2014, Perry et al 2012). Therefore, the increased NSPC proliferation found in AD cases may be a compensatory response to the pathologic changes of AD. Different types of AD pathology such as cholinergic degeneration may have detrimental effects on adult neurogenesis (Perry et al 2012). In addition, NSPC proliferation was reported to increase prior to the appearance of A β pathology, although the viability of NSPCs declines over the time course of AD pathogenesis (Chishti et al 2001, Kanemoto et al 2014). Studies of neurogenesis in AD are dependent on the use of AD transgenic mouse models involving knock-in of various FAD-related human APP mutations, often in combination with PS (PS1 or PS2) or tau mutations (Chishti et al 2001, Citron et al 1998, Mullan et al 1992, Oddo et al 2003). Indeed, these APP variants, or different PS species or other critical molecules may account, at least to some degree, for the observed divergence in neurogenic fate in the different AD transgenic models from these studies (Lazarov & Marr 2010). In addition, different APP or PS mutations or other gene insertions may exert different effects on adult neurogenesis, which also may be a cause of the inconsistent results obtained among these AD animal studies.

1.8.1.1 Role of PS on adult neurogenesis

PS is the catalytic domain of the γ - secretase complex (De Strooper et al 1998), which is needed for proteolysis of a number of transmembrane proteins including Notch and β - catenin as well as APP (De Strooper 2003). PS mutants that are especially common in PS1 contribute to A β pathology by increasing the production of A β cleaved at position 42 (De Strooper et al 1998). PSs play a role in neural development or adult neurogenesis (Chen et al 2008, Shen et al 1997, Wong et al 1997). However, the precise mechanism by which PSs influence development is still unknown (Lazarov & Marr 2010). However, PS1 null mice display severe abnormalities in somitogenesis as well as neurogenesis in the brain (Shen et al 1997, Wong et al 1997). PS1 null mice do not survive postnatally, which has complicated studies of PS1 function in postnatal development (Shen et al 1997, Wong et al 1997). For this reason, investigations into the role of PS in neurogenesis in vivo have been carried out in PS1 transgenic mice (Lazarov & Marr 2010) or mice with a conditional deletion of the PS1 gene (Chen et al 2008, Feng et al 2001). In contrast, PS2 null mice show no obvious deficit, although inactivation of both PS2 and PS1 is embryonic lethal, suggesting functional redundancy between PS1 and PS2 (Donoviel et al 1999).

PS1 is expressed in NSPCs from neurogenic regions in the adult brain (Wen et al 2002a) and wild-type PS1 overexpression has been found to increase neurogenesis (Wen et al 2002b). Mice carrying a partial deletion of PS1 show no obvious difference in NSPC proliferation, which may possibly be explained by the compensatory effect exerted by PS2 (Chen et al 2008, Donovan et al 1999). Nevertheless, mice harbouring a conditional PS1 deletion, but a conventional PS2 deletion, exhibit both enhanced NSPC proliferation and neuronal differentiation

(Chen et al 2008). This increased NSPC proliferation and differentiation may be due to the pathological stimulation, such as cortical neuron loss in mice with partial deletion of PS1 and completely null of PS2 (Chen et al 2008).

Surprisingly, mice with a single mutation in PS1 have only a minor reduction in the number of neuroblasts. Nevertheless, a dramatic decline in neuroblast number was observed in APP/PS1 double knock-in mice bearing a FAD mutation (Zhang et al 2007a). Most murine studies link FAD-associated PS1 mutations with impaired neurogenesis. For example, a more recent study reported that endogenous expression of a FAD-linked PS1 variants may be sufficient to impair NSPC proliferation, differentiation and survival (Veeraraghavalu et al 2010, Veeraraghavalu et al 2013), and another study suggested that expression of several PS1 mutants in NSPCs derived from the V-SVZ leads to a deficit in self-renewal and premature differentiation towards a neuronal fate in vitro (Gadadhar et al 2011, Lee et al 1997, Thinakaran et al 1996). In addition, PS may act as regulators of adult neurogenesis possibly via Notch signaling (De Strooper et al 1999, Wong et al 1997), therefore the FAD-related PS1 variants may cause a loss of PS1 function in neurogenesis regulation (De Strooper 2007).

1.8.1.2 Role of tau in neurogenesis

The pathological tau isoforms found in AD also have been suggested to have an impact on neurogenesis (Pristera et al 2013) although comparatively fewer studies have focused on that idea. A transgenic mouse model conditionally expressing a fragment of hyper - phosphorylated tau had reduced NSPC proliferation and survival and increased cell death in the dentate gyrus of hippocampus (Pristera et al 2013).

However, another study using animals containing a human tau isoform exhibited increased neurogenesis and cell cycle events (Schindowski et al 2008). This opposite result may be due to different pathological forms of tau being employed in the studies, therefore the impact of tau in neurogenesis warrants further study.

1.9 Stem cell therapy for neurodegenerative disease

Neuron and glial cell loss occurs in many neurodegenerative diseases (Lindvall & Kokaia 2010, Lunn et al 2011). Acute neurodegenerative disorders, caused for example by insults such as stroke or traumatic brain injury, result in a loss of neurons at the specific injury sites (Lindvall & Kokaia 2010). Chronic neurodegeneration involving a slower progressive time course is often characterized by gradual loss of a particular subtypes of neurons (such as dopamine neurons in PD) or by a continuous degeneration of a wide range of neuronal subtypes as occurs in AD (Lindvall & Kokaia 2010).

Stem cell - based therapies have been proposed for the treatment of neurodegenerative diseases as a means of replacing impaired or dysfunctional neurons. Stem cells exhibit either pluripotency or multipotency, and have the capacity to self-renew and proliferate (Reynolds & Weiss 1992, Thomson et al 1998). Stem cell based – approaches are dependent on two fundamental concepts: 1) manipulation of endogenous neurogenesis to generate new neurons at the neurogenic niches. Neurons can migrate toward the injured sites to improve self-repair; and 2) transplantation of stem cells into the sites of damage either to replace loss or defective neurons or other types of cells, or to deliver growth factors to the damage site to promote regenerations (Lindvall et al 2012).

Stem and progenitor cells derived from different sources can be used to treat neurodegenerative disorders. They may be stem cells or neural stem cells or progenitor cells (Lunn et al 2011). Stem cells are a group of cells which have the ability to undergo self-renewal consistently and possess pluripotent behaviour and are able to differentiate into a broad range of cell types which give rise to cells found in various tissues and organs (Kim & de Vellis 2009). Two types of mammalian pluripotent stem cells have been identified, embryonic stem cells (ESCs) isolated from the inner cell mass of blastocysts (Thomson et al 1998) and embryonic germ cells (EGCs) obtained from post-implantation embryos (Donovan 2001, Shamblo et al 1998). However, multipotent NSPCs can be derived from fetal or adult neural tissues, or differentiate from ESCs or EGCs (Gaspard & Vanderhaeghen 2010, Wichterle et al 2002). Alternatively, mesenchymal stem cells (MSCs) that are isolated from adult bone marrow have been thought of as a source to produce the cells with multipotency and ability to self-renew (Satija et al 2009). MSCs differentiate into osteoblasts, chondrocytes as well as adipocytes in natural conditions. They are also able to trans-differentiate into neural specific lineages (Satija et al 2009). MSCs are an autologous cell source that may tolerate immune rejection after transplantation, but MSCs derived from patients with a genetic disease might not be expected to be a source of functional cells for transplantation (Cho et al 2010a, Choi et al 2010).

Recently, a new class of pluripotent stem cell has been developed by converting adult somatic cells such as skin fibroblasts or peripheral blood into ESC-like cells via the introduction of embryogenesis-related genes (Gianotti-Sommer et al 2008, Park et al 2008c, Takahashi et al 2007, Yu et al 2007). These cells are known as induced pluripotent stem cells (iPSCs). iPSCs are created by reprogramming somatic cells to

convert them into ESC-like cells through the introduction of selected transcription factors, such as Oct 3/4, Klf, Sox2 and c-Myc (Yamanaka 2008). Various combinations of factors are used to reprogram of fibroblasts. The factors are delivered through a vector, virus, protein or RNA-based approaches (Cho et al 2010b, Hanley et al 2010, Judson et al 2009, Yakubov et al 2010). iPSC technology allows cells to be induced to an ESC - like state to generate any cell-type, such as major cell types, that are then potentially useful for clinically therapeutic applications (Park & Eve 2009, Srivastava et al 2008, Yamanaka 2008). In addition, this technology provides an endless supply of stem cells from differentiated somatic cells of patients who are affected by neurological diseases (Kiskinis & Eggan 2010, Park et al 2008b, Salewski et al 2010). iPSCs are therefore patient-specific and thereby minimize the clearance of host rejection (Araki et al 2013) compare to traditional ESCs which are derived from other source of donors. Importantly, the programmed differentiation of patient iPSCs can possibly be developed to model the pathogenesis of human disease for understanding the underlying mechanism as well as for development of potential therapeutic strategies (Lunn et al 2011). However, there are still major obstacles to overcome before iPSCs become clinically acceptable for the treatment of human neural disorders. Firstly, cell reprogramming by either transcription factors or expressed proteins may introduce immunogenicity (Araki et al 2013, Zhao et al 2011) and tumorigenicity (Duinsbergen et al 2009, Lee et al 2013a, Miura et al 2009, Nakagawa et al 2008, Yu et al 2013) in recipients. Secondly, and importantly, the efficiency of generating specific neural cell type for the treatment of disease, and integration of iPSCs- derived neural cells into existing neural circuits, should be taken into consideration (Cai et al 2010, Tsuji et al 2010, Yu et al 2013).

In summary, stem cell based therapies are potentially a promising treatment for neurological disorders, as various preclinical studies have illustrated functional recovery in animal models after transplantation of neurons derived from ESCs or iPSCs (Hargus et al 2010, Ma et al 2012b, Roy et al 2006, Wernig et al 2008). However, the basis of improvement, whether it is due to the repair and reorganization of the damaged neural network and the neurotrophic supports, or through other mechanisms, requires further clarification. At least four issues have to be considered regarding safety and efficacy when stem cell therapy is applied to humans: 1) long-term survival and phenotypic and functional integration of the graft (either neurons or glial cells); 2) high purity of the graft to avoid unexpected interaction with circuitry; 3) tumours which may arise from the grafts; 4) and compatibility with the native immune system (Kim & de Vellis 2009).

1.10 APP and neurogenesis

Several studies showed that APP expression is increased during period of neuronal differentiation in embryonic CNS development (Clarris et al 1995, Hung et al 1992, Salbaum & Ruddle 1994). In addition, abundant APP expression in radial glial cells of developing fetus mouse brain can be detected by immunocytochemistry (Trapp & Hauer 1994). Thus APP has been suggested to play a role in embryonic CNS development (Clarris et al 1997, Hung et al 1992). There are many studies on the role of APP or its homologues, APLP1 and APLP2 in CNS development and neurogenesis (Bergmans et al 2010, Chen & Tang 2006, Heber et al 2000, Kanemoto et al 2014, Ma et al 2008b, Shariati et al 2013, Young-Pearse et al 2007). However, the present roles of these proteins remain unclear.

APP null mice do not exhibit extremely severe brain deficiencies, although they have reduced brain weight and other neuronal function impairments such as hypersensitivity to seizure-inducing stimuli (Dawson et al 1999, Muller et al 1994, Steinbach et al 1998, Zheng et al 1996, Zheng et al 1995). Interestingly, NSPCs isolated from APP KO mice at embryonic 14 days showed increased cell proliferation (Zheng et al 1996, Zheng et al 1995). However, another study indicates an opposite result, with decreased proliferation of NSPCs obtained from postnatal mice lacking APP, compare to the corresponding wild-type controls (Hu et al 2013). The differences may be due to the investigations being carried out on NSPCs cultured from different developmental time points, i.e. one from an embryonic stage, and the other from a postnatal stage (Hu et al 2013, Zheng et al 1996, Zheng et al 1995). The reason for the discrepancy in the results is unknown. However, the in vitro neurosphere culture system is very sensitive to culture methods and environment, which may change the sensitivity or likelihood of NSPC proliferation and differentiation (Jensen & Parmar 2006). In addition, a recent in vivo study reported increased NSPC proliferation in the adult hippocampus of APP null mice compared to the WT control mice (Wang et al 2014). A physiological deficiency in APP null mice (Dawson et al 1999, Steinbach et al 1998, Zheng et al 1996, Zheng et al 1995) may have been the cause of the increased neurogenesis (Hayon et al 2013, Jin et al 2004b, Jin et al 2004c, Parent et al 1997). For example, mice lacking APP are hypersensitive to seizures (Steinbach et al 1998), and this increased seizure activity may induce enhanced neurogenesis on cell proliferation at neurogenic niches (Parent et al 1997, Shetty et al 2012). Thus, the definitive role of APP in NSPC proliferation remains unclear.

One study has found a reduced number of spines on dendrites of cortical and hippocampal neurons in APP-KO mice (Lee et al 2010). In addition, APLP2 is reported to control differentiation of neural stem cells during cortical development (Shariati et al 2013). Furthermore, early postnatal death and serious neuroanatomical abnormalities are observed in APP/APLP2 and APLP1/APLP2 double knockout mice (Chen & Tang 2006, Heber et al 2000, von Koch et al 1997). This suggests that APP and its homologue probably play a critical role in embryonic and adult neurogenesis, and it also suggests a possible compensatory relationship between APP and APLP2 (Heber et al 2000, von Koch et al 1997). APP/APLP1/APLP2 triple knockout (APP tKO) mice have been created for deeper investigations of the function of the APP family in CNS development. Unfortunately, APP tKO mice are perinatally lethal (Herms et al 2004). However, NSPCs derived from APP tKO ESCs display normal neuronal differentiation and generation of active excitatory synapses in vitro, and the APP tKO neurons are morphologically intact and exhibit normal migration in vivo (Bergmans et al 2010). The reason for the death of the embryo around birth is unknown (Bergmans et al 2010). As the NSPC culture system is sensitive to culture conditions, the properties of cells to differentiate may change in response to the method and the environmental conditions applied in culture (Jensen & Parmar 2006). The reason why NSPCs response differently in vitro to in vivo may be that cells in vivo are regulated by a more complicated system of cellular interaction than cells in vitro (Jensen & Parmar 2006). To identify the role of APP in NSPC differentiation, more studies are required.

Numerous reports suggest that APP metabolites may be involved in the regulation of adult neurogenesis, especially in the modulation of NSPC proliferation or

differentiation. For example, a soluble form of APP, sAPP α has been reported to have a trophic effect on NSPC behaviour (Baratchi et al 2012, Caille et al 2004, Demars et al 2011, Ohsawa et al 1999). However, little is known about how sAPP α achieves the observed effects on NSPC division.

A β is another proposed regulator of NSPC proliferation and differentiation. However, the studies reporting an effect of A β are controversial or even conflicting (Chen & Dong 2009, Haughey et al 2002, Kanemoto et al 2014, Lee et al 2013b, Sotthibundhu et al 2009, Zheng et al 2013). These contradictions may be due to the different forms of A β used in the studies. Furthermore, various ways that A β is reconstituted in each study may alter the A β configuration, and subsequently affect NSPC behaviour. Notably, synthetic A β reportedly exerts different effects on NSPC biology than the endogenous A β (Nicolas & Hassan 2014).

The role of the intracellular domain of APP (AICD) in NSPC regulation remains unclear as well. By analogy with the NICD, AICD is suggested to act as a transcriptional factor that negatively regulates the gene encoding the EGFR, which is known to stimulate NSPC proliferation (Ayuso-Sacido et al 2010, Zhang et al 2007b). Besides, a study on APP KO mice that express AICD exhibit reduced proliferation as well as reduced survival of NSPC in the DG-SGZ (Ghosal et al 2010). In addition, one study suggests that transient axonal glycoprotein 1, a neural cell adhesion protein, interacts with APP to initiate AICD release to negatively modulate neurogenesis (Ma et al 2008b). However, another study does not agree with the conclusion that AICD participates in the nuclear signaling pathway (Hebert et al 2006), highlighting further studies in this area.

1.11 Hypothesis and aims of the study

APP has been well studied for its role in AD pathogenesis. Nevertheless, the functions of APP still remain a mystery, despite intensive research in this area. Understanding the biological function of APP may contribute to our understanding of the role of APP in AD and the normal biology of brain.

APP is likely to participate in a variety of neuronal and non-neuronal processes including synaptogenesis, neurite outgrowth and the regulation of a number of genes vital for cell viability (Allinquant et al 1995, Bush et al 1990, Lee et al 2009, Murayama et al 1996). APP expression occurs coordinately with stem cell maintenance in the developing rat olfactory system, and in neural stem cells (radial glial) during embryonic development (Clarris et al 1995, Trapp & Hauer 1994). Furthermore, APP is expressed in both neuroblasts as well as in neurons when NSPCs proliferate and differentiate (Clarris et al 1995, Fukuchi et al 1992, Masliah et al 1992). As adult neurogenesis is suggested to play a role in normal brain function (Akers et al 2014, Dupret et al 2007, Snyder et al 2011), APP could potentially regulate neural stem cell development. AD treatments that alter APP processing could potentially affect normal brain function. For this reason, it is vital to understand whether APP or APP metabolites exert effects on adult neurogenesis.

1.11.1 Hypothesis

The central hypothesis presented in this study is that APP regulates NSPC proliferation and differentiation. It is further hypothesized that APP – induced NSPC proliferation is due to the production of a secreted growth factor.

1.11.2 Aims

To test this hypothesis, the following specific aims of this study have been covered in each chapter:

1. To examine the effect of APP on NSPC proliferation. The proliferation of NSPCs obtained from APP overexpressing mice model (Tg2576 mice) and APP null mice (APP KO) grown in culture was compared with that of the corresponding background strain control.
2. To identify mechanisms by which APP may affect NSPC proliferation. Conditioned medium collected from Tg2576 NSPC culture and APP KO culture was analyzed to determine whether APP expression drives production of any secreted factors that could promote NSPC proliferation.
3. To examine the effect of APP on NSPC differentiation. NSPCs were derived from APP overexpressing Tg2576 mice and APP null mice and from background strain control mice. Upon differentiation of NSPC, the production of neurons, astrocytes as well as oligodendrocyte progenitors was examined.
4. To identify mechanisms by which APP may affect NSPC differentiation. The NSPC cultures were treated with the secreted fragments of APP ($A\beta$ and sAPP α). The expression of neuronal, astrocytic and oligodendrocyte progenitor markers in the cultures was examined.

Chapter 2

Materials and Methods

2.1 Materials

All materials and suppliers of materials used in this study are shown in Table 2.1.

2.1.1 Buffers, solutions and cell culture media

All solutions and buffers used in this study are listed in Table 2.2.

2.1.2 Antibodies used in this study

All the primary antibodies and all the secondary antibodies, and supplies of antibodies are shown in Table 2.3.

2.2 Methods

2.2.1 Neural stem/progenitor cell (NSPC) culture

NSPC cultures were prepared from the cerebral cortices of new born (postnatal day 0, P₀) human APP overexpressing mice (APPSW Tg2576, Taconic Farms Inc., Hudson, NY) and from the corresponding wild type littermate controls (C57Bl6 × SJL), as well as from APP knockout (APP KO, Jackson laboratory, Bar Harbor, ME) mice and the corresponding background wild type controls (C57Bl6). Brain cortices were dissected away from the cerebellum, and then the meninges and hippocampus also removed. Cortical tissue was then incubated in 1×TrypLE Express for 10 minutes at 37°C. The cortices were mechanically dissociated with 1000 µL fine tips, and then passed through a 40 µm cell strainer (BD Bioscience, North Ryde, Australia) to remove non-separated cells. NSPCs were plated at a density of 20,000 cells/mL in T75 cell culture flasks containing proliferation medium (Table 2.2). NSPCs were grown as neurospheres in suspension and incubated in a humidified incubator at 37°C with 5% CO₂. After culturing for 7 days, the neurospheres were collected by

centrifugation at $500 \times g$ for 5 min at room temperature. Conditioned medium was collected and stored at -80°C for further analysis. Neurospheres pellets were passaged by mechanical dissociation with 200 μL fine tips. Cells were stained with Trypan Blue and counted under a Bright-LineTM haemocytometer (Sigma-Aldrich Pty. Ltd. Castle Hill, Australia). The dissociated cells were passaged and either reseeded as suspension cultures or replated as adherent cultures for further analysis. All cultures were incubated in a humidified incubator at 37°C with 5% CO_2 .

For proliferation and differentiation assays, the NSPCs were cultured adherently on poly-L-lysine coated plates in either proliferation or differentiation medium (Table 2.2).

2.2.2 AlamarBlue Assay

The cell viability (expressed as an index of the number of living cells) was measured by an alamarBlue assay. The isolated cells were plated adherently in 100 μL of fresh proliferation medium on poly-L-lysine coated 96-well-plates at a density of 2000 cells/well, and to each well was added 100 μL of medium containing the treatment. The cells were then incubated for various time periods, after which 20 μL of alamarBlue reagent (Hayashi et al 1994) was added into each well and the plates incubated for another 4 hours. The fluorescence intensity was measured using a FLUOstar Optima microplate fluorescence plate reader (BMG Labtech GmbH Ortenberg, Germany) at an excitation wavelength of 540 nm and an emission wavelength of 590 nm. Cell number was presumed to be proportional to the relative fluorescence intensity.

2.2.3 EdU incorporation assay of cell proliferation

Cells were cultured adherently on a coverslip in proliferation media at a density of 20,000 cells per well in a 24 well-plate in which each well contained one coverslip. Plates were followed by incubated for up to 4 days, after which cells were incubated with 10 μ M EdU for 8 hours before fixing in 4% (w/v) paraformaldehyde in PBS for 15 min. Cells were permeabilized with 0.03% (v/v) Triton X-100 in PBS for 5 min, and then and then blocked in 4% goat serum in PBS for 20 min. The fixed cells were immunostained with polyclonal anti-Nestin antibody 1:5000 diluted in 2% goat serum in PBS and then incubated with a goat anti-mouse IgG conjugated to Alexa Fluor 488 (1:1000 diluted in 2% goat serum in PBS). EdU was visualized using the AlexaFluor-594 Click-iT® EdU Cell Proliferation Assay Kit immediately after immunocytochemistry. Cells on the coverslip were incubated at 20°C-23°C for 15 min in PBS/0.5% (v/v) Triton X-100 and then transferred to the EdU developing cocktail (10% v/v Click-iT® reaction buffer, 10% v/v Click-iT® reaction buffer additive, 80% v/v CuSO₄), incubated in the dark at 20°C–23°C for 30 min, and washed in PBS. Cell nuclei were counterstained by poststaining with 4',6-diamidino-2-phenylindole (DAPI) at 1:7000 dilution before coverslips were mounted in fluorescence mounting medium (DAKO, Cytomation). An estimate of cell proliferation was given by the proportion of cells positive for EdU staining in the population of cells positive for both of DAPI and Nestin (Alexa Fluor 488).

2.2.4 NSPC differentiation and cell fixation

Neurospheres were mechanically dissociated, and then isolated cells were plated at a density of 10⁵ cells/well on poly-L-lysine coated coverslips in differentiation medium

(Table 2.2) in 24-well plates. After 5-14 days in culture, the cells were washed once with pre-warmed (37°C) phosphate-buffered saline (PBS) and then by fixed with 4% (w/v) paraformaldehyde in PBS for 10 minutes. Fixed cells were washed with PBS three times and kept in PBS containing 0.02% (w/v) sodium azide at 4°C for immunocytochemistry (Chapter 4).

2.2.5 Immunoblotting

Samples had a one fifth volume of $5 \times$ Laemmli sample buffer (Table 2.2) added and heated to 95°C for 5 minutes. Mixtures were centrifuged at 10,000 rpm (microcentrifuge) for 5 seconds. Samples were loaded onto 8 or 12% sodium dodecyl sulfate Tris-glycine polyacrylamide (SDS-PAGE) gels to separate protein according to their size by electrophoresis. Proteins were then transferred electrophoretically onto polyvinylidene difluoride (PVDF) membranes. The PVDF membrane was blocked in 5% (w/v) skim milk powder in 50 mM Tris-buffered saline (pH 8) containing 0.05% (v/v) Tween 20 (TBS-T) for one hour. Specific proteins were detected by incubation with primary antibody at the appropriate dilution in 5% (w/v) skim milk TBS-T buffer at 4°C overnight. Membranes were washed for three or four times in TBS-T before incubation with HRP-conjugated secondary antibody of the appropriate dilution in 5% (w/v) skim milk TBS-T at room temperature for one hour. Chemiluminescence was measured using a CHEMI-SMART 5000 image acquisition system and images were captured using Chemi-Capt version 5001 (Vilber-Lourmat, Eberhardzell, Germany). For quantification of immunoreactivity, the integrated density of luminescence for each band was analyzed using ImageJ version 1.46.

Table 2.1 Materials and supplies of materials used in this study

Material	Supplier
30% Acrylamide/Bis solution	Bio Rad Laboratories Pty. Ltd., Gladesville, Australia
4',6-Diamidino-2-Phenylindo, Dihydrochloride (DAPI)	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Acetic acid (glacial)	Merck Serono Australia Pty. Ltd., Frenchs Forest, Australia
AlexaFluor-594 Click-iT EdU cell proliferation Kit	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Antipain ($C_{27}H_{44}N_{10}O_6$)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
sAPP α recombinant protein	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
B27 supplement	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Bio Rad DC protein assay kit	Bio Rad Laboratories Pty. Ltd., Gladesville, Australia
Bio Rad Silver Stain Plus Kit	Bio Rad Laboratories Pty. Ltd., Gladesville, Australia
cOmplete ultra protease inhibitor tablets	Roche Diagnostics Australia Pty. Ltd., Castle Hill, Australia
CNBr-activated Sepharose 4B	GE Healthcare Bioscience AB, Uppsala, Sweden
Chymostatin ($C_{31}H_{41}O_6N_7$)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
CysC primer (Cst 3) primer	Qiagen Pty. Ltd., Chadstone, Australia
CysC (Recombinant mouse protein)	R&D Systems, Inc. Minneapolis, MN
CysC (Recombinant human protein)	My Biosource San Diego, CA, USA
DAKO mounting medium	DAKO Australia Pty. Ltd. Campellfield, Australia
Deoxyribonuclease-1	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Dimethyl sulfoxide (DMSO)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Dulbecco's modified Eagle's medium (DMED)	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
E-64 ($C_{15}H_{27}N_5O_5$)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Fatty – acid free bovine serum albumin (BSA)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia

Table 2.1 (Cont'd)

Material	Supplier
Fetal bovine serum (FBS)	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Glyceraldehyde-3-phosphate dehydrogenase GAPDH primer	Geneworks Pty. Ltd., Hindmarsh, Australia
Glycerol	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Glycine	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Human recombinant epidermal growth factor (EGF)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Human recombinant basic fibroblast growth factor (bFGF)	PeproTech. Rocky Hill, USA
Hydrochloric Acid (37% w/v)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Igepal CA-630	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Immobilon chemiluminescent substrate	Millipore Australia Pty. Ltd, Kilsyth, Australia
β -Mercaptoethanol	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Non-fat dry milk powder	Woolworths Ltd., Bella Vista, Australia
Normal goat immunoglobulin G	R&D Systems Inc. Minneapolis, MN
Normal mouse immunoglobulin G	Dako Australia Pty. Ltd. Campbellfield, Australia
Paraformaldehyde	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Penicillin /streptomycin	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Pepstatin A (C ₃₄ H ₆₃ N ₅ O ₉)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Poly-L-lysine	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Polyvinylidene fluoride membrane (PVDF)	Bio Rad Laboratories Pty. Ltd., Gladesville, Australia
Potassium Chloride (KCl)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Potassium dihydrogen phosphate (KH ₂ PO ₄)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Protein G agarose gel	Roche Products Pty. Ltd., Dee Why, Australia
RNeasy mini Kit	Qiagen Pty. Ltd., Chadstone, Australia

Table 2.1 (Cont'd)

Material	Supplier
RT2 First Strand Kit	Qiagen Pty. Ltd., Chadstone, Australia
Silver Stain Plus	Bio Rad Laboratories Pty. Ltd., Gladesville, Australia
Sodium azide (NaN ₃)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Sodium bicarbonate (NaHCO ₃)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Sodium chloride (NaCl)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Sodium deoxycholate	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Sodium dodecyl sulfate (SDS)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Sodium phosphate monobasic dehydrate (Na ₂ HPO ₄)	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
SYBR master mix	Qiagen Pty. Ltd., Chadstone, Australia
Synthetic human sequence A β peptides (>95% pure)	Keck Biotechnology Resource Laboratory, New Haven, CT
Tris-base	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Triton X-100	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia
Trypan Blue	Sigma-Aldrich Pty. Ltd. Castle Hill, Australia

Table 2.2 Cell culture media, buffers and solutions used in this study

Cell culture media		
NSPC medium	Proliferation	Dulbecco's modified Eagle's medium (DMEM) contain 2% v/v B27 supplement, 20 ng/mL Human recombinant EGF and Human recombinant bFGF, 100 U/mL penicillin and 100 µg/mL streptomycin.
NSPC medium	Differentiation	Dulbecco's modified Eagle's medium (DMEM) contain 2% v/v B27 supplement, 1% v/v heat inactive fetal bovine serum (FBS), 100 U/mL penicillin and 100 µg/mL streptomycin.
Buffers and solutions		
Phosphate buffered saline (PBS)		137 mM NaCl, 2.7mM KCl, 1.8 mM KH ₂ PO ₄ , 10 mM Na ₂ HPO ₄ , pH 7.4
TBS-T (Immuno blotting)		150 mM NaCl, 25 mM Tris-base, pH 8.0, 0.05% Tween-20
SDS-PAGE running buffer		25 mM Tris-base, 192 mM Glycine, 0.1% SDS
Laemmli sample buffer		10% β-Mercaptoethanol, 2% SDS, 50 mM Tris pH 6.8, 10% glycerol
Cell lysis buffer		150 mM NaCl, 50 mM Tris, 0.5% w/v sodium-deoxycholate, 1% v/v Igepal-CA630, 0.1% SDS, pH 7.4
Electroblotting buffer		25 mM Tris, 192 mM Glycine, 20% v/v Ethanol pH 8.4
Coupling buffer		0.1 M NaHCO ₃ , 0.5 M NaCl, pH 8.3
Acetate buffer		0.1 M Acetate acid, 0.5 M NaCl, pH 4.0

Table 2.3 Primary and secondary antibodies used in this study

Primary antibodies			Supplier
Mouse anti human - A β monoclonal antibody (IgG) 6E10			Covance Pty. Ltd. North Ryde, New South Wales, Australia
Goat anti mouse IgG CysC antibody			R&D Systems Inc. Minneapolis, MN
Mouse anti IgG β III tubulin monoclonal antibody			Promega Australia (Alexandria, Australia)
Rabbit anti IgM NG2 chondroitin sulfate proteoglycan antibody			Millipore Australia Pty. Ltd, Kilsyth, Australia
Mouse anti IgG GFAP monoclonal antibody			BD Bioscience North Ryde, New South Wales, Australia
Rabbit anti IgG Nestin polyclonal antibody			Millipore Australia Pty. Ltd, Kilsyth, Australia
Secondary antibody	Conjugate	Application	Supplier
Goat - anti mouse IgG	Alexa 594	ICC	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Goat - anti mouse IgG	Alexa 488	ICC	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Goat - anti rabbit IgM	Alexa 594	ICC	Life Technologies Australia Pty. Ltd., Mulgrave, Australia
Rabbit anti- Goat	HRP	WB	DAKO Australia Pty. Ltd. Campbellfield, Australia
Goat anti - mouse	HRP	WB	DAKO Australia Pty. Ltd. Campbellfield, Australia

Chapter 3

Role of CysC in amyloid precursor protein-induced proliferation of neural stem/progenitor cells

3.1 Introduction

NSPCs are self-renewing, multipotent cells that can produce all of the major cellular phenotypes in the nervous system (Reynolds & Weiss 1992) (Richards et al 1992). NSPCs are important, not only because they produce the entire complement of neuronal and glial cells of the mature nervous system and continue to generate new neurons throughout life, but also because they may be useful for the therapeutic replacement of cells in neurodegenerative diseases. The mechanisms that promote NSPC proliferation are only partially understood (Small et al 2001). Both EGF and bFGF are well studied for their roles in stimulating NSPC proliferation *in vitro* (Moyse et al 2008, Richards et al 1992). However, other autocrine growth factors produced by the NSPCs themselves may also be necessary for optimum growth.

APP is a 110–130 kDa integral type I transmembrane glycoprotein that has been extensively studied for its role in Alzheimer's disease (AD) (Small et al 2001). Despite the very large number of published studies on APP, the normal function of APP remains a mystery. APP is encoded by a single gene located on chromosome 21 (Patterson et al 1988). APP is post-translationally glycosylated and phosphorylated and can be cleaved by two major proteolytic pathways. In one pathway, sequential cleavage of APP by α - and γ -secretase generates a large ectodomain fragment (sAPP α), that is secreted into the extracellular milieu, and a small C-terminal fragment (the APP intracellular domain, or AICD), that may have a role in regulating gene expression. In the other pathway, cleavage of APP by β - and γ -secretase generates a different ectodomain fragment (sAPP β), as well as the AICD peptide. Cleavage of APP via this second pathway also generates the A β that accumulates in AD (Nunan & Small 2000).

Although the normal function of APP is poorly understood, the pattern of APP expression suggests that it may be important for neuronal growth and differentiation, not only in the developing brain but also in the mature or aging nervous system. The expression of APP has been shown to increase as the nervous system matures (Small et al 1992). APP expression increases as NSPCs mature into neurons, and soluble APP has been reported to promote neural differentiation (Freude et al 2011, Lee & Cole 2007). APP may also play a role in later stages of neuronal development. For example, soluble APP is reported to stimulate neurite outgrowth in a variety of cell systems (Chasseigneaux et al 2011, Gakhar-Koppole et al 2008, Mattson et al 1992, Milward et al 1992, Small et al 1994). Studies have shown that APP expression is increased in the olfactory neuroepithelium at the developmental stage when neurogenesis and neurite outgrowth begin (Clarris et al 1995). Similarly, APP has been reported to regulate a number of developmental functions including neuronal migration (Young-Pearse et al 2007) and cell growth (Hornsten et al 2007, Joshi et al 2009). A role for APP in cell growth is supported by the rapid up-regulation of APP that occurs in response to axonal injury (Blumbergs et al 1995, Gentleman et al 1993, Itoh et al 2009). Dystrophic neurites found around amyloid plaques are highly immunoreactive for APP (Cochran et al 1991, Cras et al 1991, Tabaton et al 1992), consistent with the possibility that APP may play a role in neural repair.

Neurogenesis is reported to be increased in transgenic mice that overexpress APP. For example, Jin *et al.* (2004) reported a 2-fold increase in BrdU-labelled cells in PDGF-APP_{sw,Ind} mice at 3 months of age. More recently, López-Toledano and Shelanski (2007) reported similar findings. The increase in neural precursor proliferation was attributed either to a compensatory mechanism resulting from disease pathology in the mice (Jin et al 2004a) or to a direct effect of A β (Lopez-Toledano & Shelanski 2007).

To address the role of APP in NSPC proliferation and neurogenesis, the growth and proliferation in culture of NSPCs derived from APP transgenic mice (Tg2576) and from APP knockout (APP-KO) mice was examined. This study showed that the proliferation rate of NSPCs from APP-overexpressing cells is increased and that the proliferation of NSPCs from APP-KO cells is decreased when compared with the corresponding background strain NSPCs. Furthermore, these studies showed that this effect is mediated by a secreted factor. Despite previous suggestions that sAPP α can influence the growth of neural stem cells (Freude et al 2011, Hayashi et al 1994, Lazarov & Demars 2012), no evidence was found that the effect on NSPCs is mediated by sAPP α . Instead, the studies showed that APP-induced NSPC proliferation is mediated, at least in part, by secreted CysC.

3.2 Materials and Methods:

3.2.1 All materials used in this chapter are shown in Table 2.1.

3.2.2 In vitro measurement of cell number and cell proliferation

Dissociated cells cultured adherently on poly-L-lysine-coated 96-well plates were incubated for various periods. Then the viable cell number was measured using the alamarBlue Assay (Chapter 2). Cell proliferation was monitored by an EdU incorporation assay (Chapter 2).

3.2.3 Effect of conditioned medium on NSPC proliferation

Conditioned medium was collected from neurosphere cultures that had been grown over a period of 7 days. To examine the effect of conditioned medium on NSPC proliferation, dissociated cells from neurospheres were cultured adherently on poly-L-

lysine-coated 96-well plates in 100 μ l/well of proliferation medium. Sixteen hours after plating, 100 μ l of conditioned medium or normal proliferation medium (control) was added, and the cells were incubated for 3 or 5 days, after which cell number was measured using the alamarBlue assay.

3.2.4 Determination of the levels of sAPP α and CysC

The level of sAPP α and CysC in conditioned medium was determined by immunoblotting. The volume per well of conditioned medium that was analyzed was adjusted so that it represented the same number of viable cells, as determined using the alamarBlue assay. Routinely, \sim 10–30 μ l was loaded into each gel lane for analysis. For the analysis of intracellular CysC, cells were washed with PBS and then lysed in cell lysis buffer prior to analysis by immunoblotting (Chapter 2).

3.2.5 Immunodepletion of sAPP α , A β or CysC

For depletion of sAPP α , A β or CysC from conditioned medium, mAb 6E10, anti-mouse CysC antibody, or goat immunoglobulin G (35 μ g) was incubated with 500 μ l of protein G agarose gel (Roche Products Pty. Ltd., Dee Why, Australia) overnight at 4°C in 5 ml of PBS. The gel was then washed three times with 5 ml PBS, after which the gel was incubated with 5.5 ml of conditioned medium for 3 h at 4°C. Finally, the gel slurry containing conditioned medium was centrifuged (10,000 \times g), and the resulting supernatant fraction was assayed by immunoblotting or used for cell proliferation experiments.

3.2.6 Real-time PCR

RNA was extracted from the neurospheres of $n = 6$ independent mouse cohorts using an RNeasy mini kit as described by the manufacturer. Each preparation of neurospheres contained \sim 10⁶ cells. Six independent RNA extracts were obtained from

each neurosphere preparation. cDNA was obtained from 400 ng of RNA with an RT2 First Strand kit as described by the manufacturer. The CysC primers (Cst3) were from Qiagen, and the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) primers were from Geneworks. Glyceraldehyde-3-phosphate dehydrogenase GAPDH was used as an internal control. CysC (Cst3) and GAPDH primers were used at a concentration of 10 μ M. All samples were diluted 1:10 and analyzed in triplicate. Standard curves for Cst3 and GAPDH with concentrations 1, 0.5, 0.25, and 0.125 μ g were used to quantify Cst3 mRNA using SYBR master mix. The results were analyzed using a LightCycler 480 (Roche Diagnostics Australia Pty. Ltd., Castle Hill, Australia).

3.3 Results

3.3.1 Analysis of APP levels and proliferation of NSPCs derived from Tg2576 mice and background strain C57Bl/6 \times SJL mice

The amount of APP immunoreactivity in neurosphere cultures was examined by western blotting to confirm that the level of APP were higher in the Tg2576 cultures than in the background strain (C57Bl/6 \times SJL) littermate control cultures. At 2, 4, and 6 days after plating, conditioned medium was analyzed for APP by immunoblotting with mAb 22C11, which recognizes both mouse and human APP and the APP homolog amyloid protein-like protein-2, and with mAb 6E10, which recognizes human sAPP α . The results confirmed that APP levels were much higher in the medium of Tg2576 neurosphere cultures than in the background strain cultures (Fig. 3.1, A). A major band of 100 – 110 kDa was detected in the medium, corresponding to the molecular mass of sAPP α .




Figure 3.1 Analysis of APP levels and proliferation of NSPCs derived from Tg2576 mice and background strain C57Bl/6xSJL mice. A. Western blot analysis of the level of APP in the conditioned medium of neurosphere cultures. Blots were stained with mAb 22C11, which recognizes both human and mouse APP, and mAb 6E10, which is specific for human-sequence APP. B. Analysis of the proliferation of NSPCs. Neurospheres were dissociated and the isolated cells seeded on poly-L-lysine coated 96-well plates at a density of 2000 cells per well. After 2, 4 or 6 days in culture, the relative number of cells was estimated using the alamarBlue assay. Fluorescence intensity was taken as an index of cell number. Values are means \pm SEM (n=3 wells). C. Phase-contrast microscopy of neurospheres after 7 days in culture. Scale bar = 200 μ m. D. Quantitation of the area of neurospheres from experiment shown in C. Values are means \pm SEM (n=200 neurospheres). E. EdU incorporation assay of cell proliferation. NSPCs were cultured in proliferation medium for 4 days prior to incubation for 8 h with EdU (30). The result shows that a greater proportion of Tg2576 NSPCs were actively proliferating (incorporating EdU) than the corresponding background strain cells. The graph shows the mean values (\pm SEM) of % cells incorporating EdU (n=30). * = significantly different from corresponding values for the background strain C57BL/6xSJL cells (P<0.05, experiment in panel B, one-way ANOVA with post-hoc Tukey's test; experiment in panel D, Student's t test).

Next, to examine the role of APP in NSPC proliferation, the proliferation of cells derived from Tg2576 mice was compared with that of the background strain controls. Neurosphere cultures were dissociated into a single cell suspension on day 7, and then cells were cultured adherently on poly-L-lysine-coated 96-well plates. The proliferation of the cells was measured using the alamarBlue assay. Fluorescence intensity in an alamarBlue assay was taken as an index of the number of viable cells. Overall, the growth rate of the cells derived from the Tg2576 mice was significantly greater ($p < 0.05$, one-way ANOVA with post hoc Tukey's test) than that of the cells derived from the background strain mice (Fig. 3.1, B).

The growth of neurospheres was also examined to determine whether the proliferation of NSPCs was increased in the Tg2576 cultures. Neurospheres were cultured for a period of 7 days, after which they were examined under phase-contrast microscopy (Fig. 3.1, C and D). Neurospheres derived from Tg2576 mice were on average greater in size than the neurospheres from the background strain mice, confirming that the growth of NSPCs in Tg2576 cultures was greater than that of the background strain cultures.

To confirm that the increased growth of the Tg2576 NSPC was due to a higher proliferation rate, proliferation was measured using an EdU uptake assay (Young et al 2013). The percentage of EdU-positive cells in the Tg2576 cultures was ~20% higher than the cultures derived from background strain mice (Fig. 3.1, E). These results clearly supported the view that the increased growth observed using the alamarBlue assay and in the neurosphere cultures was due to an increase in the amount of NSPC proliferation.

3.3.2 Analysis of proliferation of NSPCs derived from APP knockout mice

To examine whether expression of endogenous mouse APP influences NSPC proliferation and to rule out the possibility that the increase in proliferation observed in Tg2576 cultures was due to a factor unrelated to APP overexpression, the proliferation rate of NSPCs from APP KO mice was also examined. NSPC proliferation was decreased in APP-KO cultures when compared with the corresponding background strain (C57Bl/6) cultures (Fig. 3.2). The growth rate of the APP-KO cells, as assessed by an alamarBlue assay, was ~60% of that of the cells derived from background strain mice (Fig. 3.2, A). Furthermore, neurospheres from APP-KO mice were smaller and less numerous than the C57Bl/6 neurospheres (Fig. 3.2, B and C). In addition, the percentage of proliferating EdU-positive cells was also significantly decreased (Fig. 3.2, D). These experiments clearly demonstrated that endogenous APP was also involved in the regulation of NSPC proliferation.

3.3.3 Effect of neurosphere conditioned medium on NSPC proliferation

To determine whether the effect of APP on proliferation to determine whether the effect on proliferation was mediated by a factor that was secreted into the culture medium, conditioned medium was prepared over 7 days from Tg2576 and the corresponding background strain (C57Bl/6xSJL) neurosphere cultures. In parallel, dissociated C57Bl/6xSJL NSPCs were plated and incubated for 16 h. Conditioned medium from the neurosphere cultures, or unconditioned proliferation medium was then added to the dissociated cell cultures. The amount of proliferation was measured after 5 days of incubation. There was a higher rate of proliferation in cultures containing conditioned medium (whether from C57Bl/6xSJL or Tg2576 cells) than in cultures containing unconditioned medium (Fig. 3.3, A).




Figure 3.2 Analysis of NSPC proliferation derived from APP knockout and from C57Bl/6 (corresponding background strain) mice. **A.** Neurospheres were dissociated and the isolated cells seeded on poly-L-lysine-coated 96-well plates at a density of 2000 cells per well. After 2d, 4d or 6d in culture, the relative number of cells was estimated using the alamarBlue assay. Fluorescence intensity was taken as an index of cell number. Values are means \pm SEM (n=3 wells). * = significantly different from corresponding values for the background strain C57Bl/6 cells ($P < 0.05$, one-way ANOVA) **B.** Phase-contrast microscopy of neurospheres after 7d in culture. Scale bar = 200 μm. **C.** Quantitation of the area of neurospheres from the experiment shown in **B.** Values are means \pm SEM (n=200 neurospheres). **D.** EdU incorporation assay of cell proliferation. The experiment was performed as described in Fig. 1. In panels **C** and **D**, the asterisk shows values that are significantly different from C57Bl/6 cultures ($P < 0.05$, Student's t test).

Importantly, the conditioned medium from the Tg2576 cell cultures stimulated cell proliferation more than the conditioned medium from the C57Bl/6xSJL cultures, supporting the view that the increased proliferation in the Tg2576 cultures was due to the secretion of a factor into the conditioned medium.

To test this hypothesis further, the effect of conditioned medium from APP-KO cell cultures on NSPC proliferation was examined. This time, isolated NSPCs grown adherently were incubated with unconditioned medium (control), conditioned medium from APP-KO neurosphere cultures, or conditioned medium from the corresponding background strain C57Bl/6 neurosphere cultures (Fig. 3.3, B). In contrast to the results with Tg2576 cultures, the APP-KO conditioned medium was significantly less potent in stimulating proliferation than the background strain conditioned medium. This result again supported the view that there was a factor secreted into the medium of APP-expressing cells that increased cell proliferation.

3.3.4 Effect of sAPP α and A β peptides on NSPC proliferation

As it has been reported that sAPP α can stimulate neural stem cell proliferation or differentiation (Freude et al 2011, Hayashi et al 1994, Lazarov & Demars 2012), the effect of recombinant human sAPP α on proliferation in our cultures was examined. However, despite repeated experiments aimed at determining whether sAPP α can stimulate NSPC proliferation, it was not possible to demonstrate any effect of sAPP α over a range of different concentrations (50–2000 ng/ml) (Fig. 3.4, A). Despite suggestions that A β amyloid may stimulate stem cell proliferation (Lopez-Toledano & Shelanski 2007), it was not possible to find any effect of A β 1–40 or A β 1–42 on proliferation (Fig. 3.4, A).




Figure 3.3 Effect of neurosphere conditioned medium on NSPC proliferation. A. Effect of unconditioned medium, conditioned medium from Tg2576 neurosphere cultures (CM-Tg2576) and conditioned medium from C57Bl/6xSJL neurosphere cultures (CM-C57Bl/6xSJL) on the growth of C57Bl/6xSJL NSPCs. Medium was conditioned for 7 d prior to adding to cultures to test for the effect on proliferation over a period of 5 d. Cell number is shown as a % of the fluorescence intensity measured in an alamarBlue assay compared to control (no conditioned medium). Values are means \pm SEM. * = significantly different from unconditioned medium (control) ($P < 0.05$). ** = significantly different from unconditioned (control) medium and from CM-C57Bl/6xSJL ($P < 0.05$). B. Effect of unconditioned medium, conditioned medium from APP knockout neurosphere cultures (CM-APP-KO) and conditioned medium from C57Bl/6 neurosphere cultures (CM-C57Bl/6) on the growth of wild-type (C57Bl/6) NSPCs. Medium was conditioned for 7 d prior to adding to cultures to test for the effect on proliferation over a period of 5 d. Cell number is represented by the % of the fluorescence intensity measured in an alamarBlue assay compared with control (no conditioned medium). Values are means \pm SEM. * = significantly different from unconditioned (control) medium ($P < 0.05$). *** = significantly different from unconditioned (control) medium and from CM-C57Bl/6 ($P < 0.05$).

To eliminate the possibility that the lack of effect of recombinant sAPP α might be due to the fact that the protein not being in a native conformation, the effect of removing endogenous sAPP α from the conditioned medium was tested (Fig. 3.4, B). More than 98% of the endogenously expressed sAPP α in the conditioned medium was removed by immunoabsorption with mAb 6E10 when compared with immunoabsorption with an immunoglobulin (Ig) control (Fig. 3.4, B *inset*). However, despite removal of most of the sAPP α , there was no decrease in the ability of the conditioned medium to stimulate proliferation. On the basis of these experiments, it was concluded that neither sAPP α nor A β (both of which are fragments of full-length APP that can be released into the culture medium) was the secreted factor that mediated the effect of APP overexpression on NSPC proliferation.

3.3.5 Levels of CysC and expression of CysC mRNA

As several studies have suggested that CysC is an important autocrine regulator of neural stem cell proliferation (Kato et al 2006, Taupin et al 2000), the possibility that CysC was the mediator of APP-induced NSPC proliferation was examined. Analysis of the conditioned medium from both Tg2576 cultures and APP-KO cultures by immunoblotting showed that CysC levels correlated with the effect on proliferation (Fig. 3.5). In addition to a major 14 kDa CysC band, a 16 kDa immunoreactive band was also present in the conditioned medium. This higher molecular mass band may represent a post-translationally modified form of CysC (Taupin et al 2000). Consistent with the view that the effect on proliferation was due to CysC, Immunoreactivity for CysC was found to be higher in the Tg2576 conditioned




Figure 3.4 Effect of sAPP α and A β peptides on NSPC proliferation. Isolated NSPCs were prepared from C57Bl/6xSJL mice and cultured adherently on poly-L-lysine. Graphs shows cell number as represented by the % of the fluorescence intensity measured in an alamarBlue assay. Values are means \pm SEM. A. Effect of recombinant sAPP α , A β_{1-40} and A β_{1-42} on NSPC proliferation. B. Immunoprecipitation of >98% of the sAPP α from Tg2576 conditioned medium did not decrease the ability of the conditioned medium to increase NSPC proliferation. The graph shows the effect of Tg2576 conditioned medium after immunoprecipitation with an immunoglobulin fraction (IP: Ig) and after immunoprecipitation with mAb 6E10 (IP: 6E10). Inset shows western blot analysis of the conditioned medium from both fractions. * = significantly different from control incubation. ** = significantly different from control incubation, but not significantly different from IP: Ig incubation.




Figure 3.5 Levels of CysC and expression of *CysC* mRNA. Figure shows representative western blots and quantification of CysC immunoreactivity by image-capture analysis (panels A-D). Values are means \pm SEM (n=4). A. Western blotting analysis of CysC in conditioned medium of Tg2576 mouse NSPC cultures and the corresponding background strain (C57Bl/6xSJL) cultures. Each lane represents a sample from a different line of NSPCs. B. Western blotting analysis of CysC in conditioned medium of APP-KO mouse NSPC cultures and the corresponding background strain (C57Bl/6) cultures. Each lane represents a sample from a different line of NSPCs. All statistical comparisons were made comparing staining intensities of images derived from the same blot. E & F. *CysC* mRNA expression determined by real-time PCR. Values are means \pm SEM (n=6). * = significantly different (P<0.05) from corresponding background strain control cultures (Student's t test).

medium (Fig. 3.5, A), and to be lower in the APP-KO cell conditioned medium (Fig. 3.5, B).

To determine whether the changes in secreted extracellular CysC reflected changes in the intracellular pool, the level of cell-associated CysC was examined (Fig. 3.5, C and D). Surprisingly, the level of CysC in the cell lysates was lower in both Tg2576 cultures and APP-KO cultures when compared with the corresponding background strain cells for each group. The level of CysC mRNA was also analyzed by real-time PCR (Fig. 3.5, E and F). These experiments showed that although the level of CysC mRNA expression was lower in the APP-KO cells, there was no significant difference in expression in the Tg2576 cells.

3.3.6 Role of secreted CysC in stimulating NSPC proliferation.

Similar to previous studies (Kato et al 2006, Taupin et al 2000), CysC increased NSPC proliferation in a concentration-dependent manner (50–100 ng/ml CysC) (Fig. 3.6, A). As the level of CysC in the conditioned medium of background strain cells was higher than this concentration range (Fig. 3.6, B panel i), this indicated that the concentration of CysC in the conditioned medium in both wild-type and background strain cells was sufficiently high to influence NSPC proliferation. Therefore, the possibility that the endogenous secreted factor in the conditioned medium was identical to CysC was tested. CysC was immunodepleted from the conditioned medium of both C57Bl/6xSJL and Tg2576 cultures (Fig. 3.6, B panel ii) and the effect of the immunodepleted medium on cell proliferation was measured.







Figure 3.6 Role of secreted CysC in stimulating NSPC proliferation. The number of viable cells was calculated from the fluorescence intensity in an alamarBlue assay.

A. Effect of CysC on NSPC proliferation. Dissociated neurosphere-derived cells were plated on poly-L-lysine-coated plates and then incubated for 5 days in proliferation medium containing various concentrations of CysC. Values are means \pm SEM, (n=4) and shown as % of the control value (no added CysC). * = significantly different from the control group (no CysC) ($P < 0.05$, ANOVA with post-hoc Tukey's test).

B. Western blots analysis of CysC immunoreactivity in cell culture medium. (i) Comparison of the level of CysC in 48 ml of C57Bl/6xSJL cell medium compared with 100 ng of recombinant CysC. (ii) Western blot stained for CysC showing efficiency of immunodepletion of CysC from the C57Bl/6xSJL and Tg2576 conditioned medium. Ig = medium after immunodepletion with control immunoglobulin; CC = medium after immunodepletion with CysC antibody. (iii) Western blot stained for Cys C showing efficiency of immunoprecipitation of CysC from the C57Bl/6 and APP-KO cell medium. C & D. Immunoprecipitation of CysC removes the APP-induced growth factor that stimulates proliferation. C. C57Bl/6xSJL neurosphere-derived cells were incubated with unconditioned medium or conditioned medium from C57Bl/6 x SJL cultures or Tg2576 cultures that was previously immunoabsorbed with a non-specific Ig fraction (IP: Ig) or with an anti-CysC antibody (IP: CC). D. APP-KO neurosphere-derived cells were incubated with unconditioned medium or conditioned medium from C57Bl/6 cultures or APP-KO cultures that was previously immunoabsorbed with a non-specific Ig fraction (IP: Ig) or with an anti-CysC antibody (IP: CC). Values in panels C and D are means \pm SEM (n=4) and shown as % of the mean of the values for unconditioned medium. * = significantly different ($P < 0.05$, ANOVA with post-hoc Tukey's test).

Immunodepletion of CysC from the conditioned medium completely removed the APP-associated increase in NSPC proliferation (Fig. 3.6, C). Proliferation was not significantly decreased when background strain NSPC conditioned medium was immunodepleted of CysC. However, after immunodepletion of CysC from the Tg2576 conditioned medium, the level of NSPC proliferation was significantly lower than the corresponding incubation in which the conditioned medium was preabsorbed with Ig.

Similar results were obtained in separate experiments using conditioned medium from C57Bl/6 and APP-KO cultures. In these experiments, the immunodepleted conditioned medium was tested on cultures of APP-KO cells (Fig. 3.6, D). Immunodepletion of CysC (Fig. 3.6, B panel iii) resulted in a significant decrease in proliferation of NSPCs when compared with the corresponding incubations in which the conditioned medium was preabsorbed with Ig. Taken together, these results clearly indicated that CysC was a major contributor to the ability of APP to increase NSPC proliferation.

3.4 Discussion

The result presented in this chapter demonstrate that APP expression regulates the proliferation of NSPCs and that this effect is mediated, at least in part, by an APP-stimulated increase in CysC secretion. The results increase our understanding of both the normal function of APP and the mechanisms involved in neural stem cell proliferation and differentiation.

The findings also provide an explanation for previous studies demonstrating that neural stem cell proliferation is increased in transgenic mice that overexpress APP (Jin et al 2004a, Lopez-Toledano & Shelanski 2007). Studies by Jin *et al.* (2004)

reported that neural stem cell proliferation was increased in PDGF-APP^{sw},Ind mice. López-Toledano and Shelanski (2006) confirmed and extended this observation. In both studies, no clear demonstration of the mechanism of increased proliferation was provided, although it was suggested that the stimulation of proliferation may have been due to A β accumulation or some aspect of associated A β pathology (Jin et al 2004a, Lopez-Toledano & Shelanski 2007). The present studies clearly support the view that NSPC proliferation is directly influenced by the expression of APP, although it is possible that a product of APP metabolism may be responsible for this effect. No direct evidence that this effect was influenced by sAPP α or A β peptides was found.

Interestingly, although neural stem cell proliferation is increased in APP-overexpressing mice, there is no obvious phenotype resulting from this increase. For example, prior to the onset of AD-type pathology, the brains of young APP transgenic mice appear relatively normal. Nevertheless, subtle abnormalities may be present in these mice. For example, overexpression of APP in transgenic mice is reported to lead to changes in synaptic density (Mucke et al 1994), and another study reports that APP transgenic mice exhibit persistent locomotor hyperactivity (Rodgers et al 2012). Whether these abnormalities are due to an increase in the number of specific neuronal populations is unclear. In the case of APP knockout mice, there are clear abnormalities, most notably agenesis of the corpus callosum (Muller et al 1994). However, whether this phenotype is due to a neural stem cell proliferation deficit is also unclear.

A surprising finding to emerge from these studies was that the effect of APP overexpression was not mediated through sAPP α . A large number of studies have

suggested that sAPP α has trophic properties on neural stem cells (Freude et al 2011, Hayashi et al 1994, Kwak et al 2006a, Lazarov & Demars 2012, Zhou et al 2011b). In the present study, recombinant human sAPP α had no effect on proliferation, nor did immunoprecipitation of endogenous secreted mouse sAPP α inhibit proliferation. Instead, immunoprecipitation of CysC from the culture medium was found to decrease the APP-induced stimulation, clearly demonstrating that some of the effect on proliferation was due to CysC. Moreover, CysC stimulated NSPC proliferation, and levels in the culture medium were correlated with APP expression, clearly supporting this idea. Although there may also be inhibitory factors that are secreted by NSPC that can affect cell proliferation, the study showed that almost all, if not all, of the APP – induced NSPC proliferation is mediated through secretion of CysC. Thus, if there were a secreted inhibitory factor, it would have to contribute only a very minor role.

Interestingly, in experiments with APP-KO cells, there was a significant residual effect of APP, even after removal of most of the CysC. CysC-immunodepleted C57Bl/6 conditioned medium was still significantly more potent in stimulating the proliferation of APP-KO cells than the corresponding immunodepleted APP-KO conditioned medium (Fig. 3.6, D). This suggests the possibility that there is an additional, as yet unidentified, secreted molecule that also contributes to APP-induced NSPC proliferation. This effect was not seen in experiments where the effect of conditioned medium on C57Bl6xSJL cells was tested (Fig. 3.6, C). The reason for this difference is unclear, but it could relate to the fact that the APP-KO cells may be more sensitive to this unidentified factor.

The mechanism of the APP-stimulated increase in CysC and NSPC proliferation is also not yet known. The experiments suggest that there may be two mechanisms involved. Real-time PCR experiments showed that *CysC* expression was decreased in APP-KO cells when compared with the corresponding background strain cells. As the extracellular domain of APP (*i.e.* sAPP α) was not found to stimulate proliferation, this suggests that the APP intracellular domain (AICD) may be involved in mediating this effect. Indeed, based on an analogy with the Notch intracellular domain (NICD), which is also released by γ -secretase (Guruharsha et al 2012), a number of studies suggest that the AICD may regulate gene expression (Pardossi-Piquard et al 2005). Whether AICD regulates the expression of CysC is not yet known and will require further studies.

However, CysC expression was not increased in Tg2576 cells when compared with the corresponding background strain cells. Indeed, a surprising finding was that although CysC secretion was higher in the Tg2576 conditioned medium (Fig. 3.5, A), levels were lower in the cell lysate (Fig. 3.5, C). This suggests the possibility that the higher levels of CysC in the Tg2576 culture medium may be due to an increased rate of CysC secretion with a concomitant decrease in intracellular CysC. The idea that increased APP may result in increased secretion is consistent with published studies. For example, another group reported that APP overexpression can increase vesicle exocytosis in PC12 cells (Lee et al 2008a). Furthermore, the cytoplasmic domain of APP can interact with proteins associated with synaptic vesicle release such as synaptotagmin-1 (Kohli et al 2012). Thus, it is possible that the cytoplasmic domain of APP may possess two different functions: 1) it may be translocated to the nucleus

to alter gene expression or 2) it may interact with proteins on the cytoplasmic leaflet of the plasma membrane to alter events such as exocytosis. The balance of these two functions could conceivably be regulated by the level of APP expression and the degree of saturation of binding to different adaptor proteins.

Finally, the study may have implications for understanding the role of CysC in the pathogenesis of AD. It is interesting to note that a polymorphism (G73A) in CysC has been linked to AD (Hua et al 2012). Furthermore, CysC is increased in regions around A β deposits in the AD brain (Kaur & Levy 2012, Steinhoff et al 2001), suggesting that it may play a role in pathogenesis or in response to A β pathology. Indeed, two groups found that CysC may have a protective effect as APP transgenic mice that had higher CysC expression were found to have diminished A β deposition (Kaeser et al 2007, Mi et al 2007). As APP is also increased in dystrophic neurites around amyloid plaques, it is tempting to speculate that increased CysC may be due to an increase in local APP expression. Further studies on the role of APP in regulating CysC expression in the AD brain may help to identify new targets for drug development in AD.

Chapter 4

Effect of cysteine protease inhibitors on NSPC proliferation

4.1 Introduction

CysC has been identified as an endogenous cysteine protease inhibitor (Bobek & Levine 1992). CysC is produced and released abundantly by a majority of tissues in mammals, and is found in high concentrations in CSF (Abrahamson et al 1986, Turk et al 2008). Recently, a potential role for cysteine proteases in NSPC proliferation has been proposed (Hu et al 2013, Kato et al 2006, Taupin et al 2000).

Cysteine proteases belong to one of five classes of proteases. These five groups include the aspartic, cysteine, metallo-, serine and threonine protease (Puente et al 2003). At present, three major families of cysteine proteases have been well studied in mammalian cells: caspases, calpains, and cathepsins. Caspases are important for programmed cell death (Fernando et al 2005, Nicholson & Thornberry 2003). However, caspase-3 activity has also been found to participate in cell development (Fernando et al 2005, Yoneyama et al 2014). Calpains are nonlysosomal proteases that, along with their endogenous inhibitor, calpastatin, participate in modulation of cell proliferation in the SVZ (Machado et al 2015, Santos et al 2012). Inhibition of calpains is also reported to increase NSPC proliferation (Machado et al 2015, Santos et al 2012). Cysteine cathepsins are secreted lysosomal peptidases that are suggested to be involved in NSPC development and differentiation (de Azevedo-Pereira et al 2011, Salvioli et al 2008). Activities of lysosomal cathepsins (Abrahamson 1994), caspases (Thornberry 1997) and cytosolic calpains (Turk et al 1997) may all be inhibited by members of the cystatin superfamily, including CysC.

Immunodepletion of CysC from APP overexpressing conditioned medium was found to prevent-APP induced NSPC proliferation (Chapter 3). Additionally, CysC has been

suggested to be an autocrine factor secreted by cells to promote NSPC proliferation in vitro (Dahl et al 2004, Taupin et al 2000). However, it still remains to be determined whether CysC acts like a growth factor (i.e binds to a growth factor receptor) or whether the effect on proliferation is mediated through inhibition of cysteine proteases.

To address this issue, NSPCs were treated with selective irreversible cysteine protease inhibitors (E-64, E-64C and E-64D), a reversible serine/cysteine protease inhibitor (antipain) an aspartic protease inhibitor (pepstatin A) as well as a broad spectrum protease inhibitor (chymostatin) to see whether other protease inhibitors could mimic the effect of CysC. The results showed that low concentrations of E-64 and antipain significantly promote NSPC proliferation in vitro, although NSPC growth was suppressed at high concentrations. An attempt was made to identify candidate cysteine proteases that might mediate the effect of CysC using affinity chromatography. However, no cysteine proteases were identified by the method. Nevertheless, the results still provide strong evidence that regulation of cysteine protease activities may affect NSPC proliferation.

4.2 Materials and methods

4.21 Materials

Stock solutions of protease inhibitors were prepared as follows. E-64 and antipain were dissolved in sterilized deionized water at a concentration of 30 mM and 50 mg/ml respectively, while E-64C, E-64D, pepstatin A and chymostatin were prepared in dimethyl sulfoxide (DMSO) at a concentration of 5 mg/ml, 10 mg/ml, 36 mM and 20 mM respectively. All the protease inhibitors were stored at -20°C prior to use.

4.2.2 NSPC proliferation

NSPCs were prepared as described in Chapter 2. Dissociated NSPCs were cultured adherently on poly-L-lysine coated 96-well plates at a density of 2000 cells per well in proliferation medium containing various concentrations of protease inhibitors (E-64, E-64C, E-64D, antipain, pepstain A and chymostatin). Cells were then incubated for 3 and 5 days. Cell number was measured using the alamarBlue assay (Chapter 2).

4.2.3 Preparation of CysC - Sepharose

To prepare a CysC - Sepharose affinity resin, recombinant CysC (1 mg) was dissolved in 3.3 ml of coupling buffer (0.1 M NaHCO₃, 0.5 M NaCl, pH 8.3) and then incubated with 170 mg of cyanogen bromide (CNBr)-activated Sepharose (prepared in 700 µl of coupling buffer) at room temperature for 2 hours. A control CNBr-activated Sepharose 4B resin was prepared by a similar method but was incubated with coupling buffer alone, lacking CysC. After formation of the covalent bond between CysC and the CNBr-activated Sepharose, the resin was centrifuged at 200g and washed with 4 ml coupling buffer 3 times to remove the excess unbound CysC. The resin was incubated with 0.2 M glycine-HCl solution (pH 8.0) for 2 hours to block any remaining active groups. The CysC-Sepharose was then washed four times with 0.1 M acetic acid solution containing 0.5 M sodium chloride (pH 4.0). Then the CysC-Sepharose resin was resuspended in 1ml of PBS and stored at 4°C for use. The control resin lacking CysC was treated by the same procedure as the CysC resin.

4.2.4 Affinity purification and analysis of fractions

For affinity purification, 500 µl of either CysC-Sepharose or control-Sepharose was incubated separately with 5 ml of NSPC-conditioned medium and unconditioned

medium respectively at 4°C overnight. The resins were washed three times with 5 ml of 1×PBS. An aliquot (200 µl) of the resin, for silver stain analysis, was resuspended in an equal volume of Laemmli sample buffer (10% (v/v) β-mercaptoethanol, 2% (w/v) SDS, 50 mM Tris-HCl pH 6.8, 10% (v/v) glycerol) and heated at 95°C for 5 minutes. The supernatant fraction was collected and 60 µl/lane loaded onto a 12% glycine SDS-polyacrylamide gel for electrophoresis. After electrophoresis, the protein was then visualized in the gel using a Bio-Rad silver stain kit (Table 2.1). The gel was fixed in a solution containing a 40% methanol/10% acetic acid (v/v) solution for 30 min. After that, the gel was immersed in Bio-Rad oxidizer for 5 min and then washed with deionized water for 15 min. The gel was then stained with the Bio-Rad silver reagent for 10 min, followed by a quick rinse with deionized water. The gel was developed with the Bio-Rad developer solution for 30 seconds until a brown precipitate appeared. Finally, the gel was immersed in 5% acetic acid (v/v) for 15 min to stop the staining reaction.

4.3 Results

4.3.1 Effect of protease inhibitors on NSPC proliferation

In order to examine whether cysteine protease inhibitors can promote NSPC proliferation, NSPCs were incubated with various protease inhibitors. The irreversible cysteine protease inhibitor E-64, the cysteine and serine protease inhibitor, antipain, the aspartic protease inhibitor, pepstatin A, and the broad-spectrum protease inhibitor chymostatin were tested over a range of concentrations, as was cysC (Fig. 4.1, A). The concentration of CysC was chosen based on two previous publications (Kato et al 2006, Taupin et al 2000), while the concentrations of Antipain, E-64, Chymostatin, Pepstatin A E-64D and E-64 C were chosen based on those recommended by the

manufacturer (see Chapter 2). Cell proliferation was estimated by cell viability, determined using an alamarBlue assay.

CysC promoted NSPC proliferation more than 50% at a concentration of 6.6 nM although it had little effect at concentrations of 3.3 nM or 1.32 nM (Fig. 4.1, A). E-64 (0.1 μ M) stimulated NSPC proliferation by nearly 70%, but inhibited NSPC proliferation at concentrations of 10 and 100 μ M (Fig. 4.1, B). Antipain promoted NSPC proliferation by approximately 30% at concentrations of 0.15 μ M and 1.5 μ M, but it suppressed NSPC proliferation at a concentration of 150 μ M (Fig. 4.1, B). The aspartic protease inhibitor, pepstatin A, and the broad-spectrum protease inhibitor, chymostatin did not stimulate NSPC proliferation at any concentration. However, both inhibitors decreased proliferation at a concentration of 10 and 100 μ M (Fig. 4.1, B). The data demonstrate that only the cysteine protease inhibitors, E-64 and antipain promoted NSPC proliferation. However, this occurred only at lower concentrations.

To investigate the mechanism of E-64's effect on NSPC proliferation further, two synthetic analogues of E-64, E-64C and E-64D, were tested for their effects on NSPC culture. E-64C is a membrane-impermeable cysteine protease inhibitor and binds to cysteine proteases in a similar manner to E-64 (Matsumoto et al 1999). E-64D, is an ethyl ester of E-64C, and is a membrane permeable inhibitor that inhibits intracellular proteases such as calpains (McGowan et al 1989).




Figure 4.1 Effect of protease inhibitors on NSPC proliferation. Dissociated neurosphere-derived cells were plated on poly-L-lysine-coated plates at the density of 2000 cells per well and then incubated for 5 days in proliferation medium containing various concentrations of inhibitors. The relative number of cells was measured using the alamarBlue assay and fluorescence intensity shown in the figure was taken as an index of cell number. Values are mean \pm SEM and expressed as % of the corresponding control (no inhibitors), ANOVA with post-hoc Tukey's test. Panel A: Effect of CysC on NSPC proliferation. * = significantly different from the control (no CysC) ($P < 0.05$, $n = 12$). Panel B: Effect of antipain, E-64, chymostatin and pepstatin A on NSPC proliferation. For antipain and E-64, * = significantly different from the control (no antipain or E-64) ($P < 0.05$, $n = 6$); for chymostatin and pepstatin A, $P > 0.05$, $n = 6$. Panel C: Effect of E-64C and E-64D on NSPC proliferation. E-64C $P > 0.05$, $n = 10$, E-64D, * = significantly different from the corresponding vehicle control ($P < 0.05$, $n = 9$).

The results indicated that E-64D slightly promoted NSPC proliferation at a concentration of 0.3 μM , whereas E-64C did not affect NSPC proliferation at the same concentration (Fig. 4.1, C). Neither E-64C nor E-64D exerted an effect on NSPC proliferation at concentrations of 3, 30 and 300 μM (Fig. 4.1, C).

4.3.2 CysC affinity purification of proteases in conditioned medium

As CysC is cell membrane impermeable and yet it stimulated cell proliferation, it was logical to look for secreted proteases that might inhibit proliferation. Cell surface might also have been a possibility. However, a number of studies show that cell – surface proteases are generally also secreted (Lozzo 1998). In order to identify proteases that might bind to CysC, CysC affinity chromatography was performed on the NSPC conditioned medium. CysC-Sepharose was incubated with NSPC conditioned medium. The affinity resin was then collected and cysteine proteases bound to the resin were eluted and analyzed by protein electrophoresis. The bands of the proteins in the gel were visualized by a Bio-Rad silver stain kit. A resin lacking CysC was used as a control for comparison (Fig. 4.2). The study was aimed at identifying cysteine protease inhibitors that bind to proteases (Fig. 4.2). However, it was still uncertain whether the protease was a cathepsin (a secreted cysteine protease) or some other kind of protease. For this reason, cathepsin might not have been an ideal positive control.

A series of proteins were identified in both the unconditioned medium (lane 1) and conditioned medium (lane 2). Although a number of proteins were present in the unconditioned medium (presumably components of original medium), no specific




Figure 4.2 Analysis of fractions from cystatin C affinity chromatography by SDS - polyacrylamide electrophoresis. Proteins were separated by gel electrophoresis, and the proteins were visualized by silver staining. Lanes 1 to 6 were loaded with different fractions collected from cystatin C affinity chromatography. 1: Unconditioned medium. 2: Conditioned medium. 3: Conditioned medium after incubation with the control CNBr Sepharose. 4: Conditioned medium after incubation with the cystatin C Sepharose. 5: Eluate collected from conditioned medium eluted from control Sepharose. 6: Eluate collected from conditioned medium eluted from cystatin C Sepharose.

band was present in lane 2, that was not also present in lane 1. In addition, fractions containing conditioned medium that had been preadsorbed with control Sepharose (lane 3) and CysC Sepharose (lane 4) respectively also had a number of proteins. Nevertheless, no clear difference was seen between the two fractions. Lane 5 and 6 that were loaded with the bound fractions from the control CNBr Sepharose (lane 5) and CysC Sepharose (lane 6). No band was observed in the region around 21-30 kDa, which is the size most cysteine proteases are predicted to be. A broad band of approximately 13-15 kDa was observed in lane 6. This protein possesses a very similar molecular weight to that of CysC previously observed in NSPC culture (Hu et al 2013). Thus, this band was probably CysC that had leaked from the CysC Sepharose. This 13-15 kDa band was probably too small to be a cysteine protease, that normally have a molecular mass in the range of 21-30 kDa (Grzonka et al 2001).

In conclusion, no cysteine proteases were detected in the conditioned medium using silver staining. It is assumed that any cysteine proteases present in the conditioned medium are likely at too low a concentration for further analysis. As no specific proteins were detected by silver staining, this approach was not pursued.

4.4 Discussion

The results presented in this chapter showed that along with CysC, the specific cysteine protease inhibitors, E-64, E-64D and antipain can promote NSPC proliferation. However, neither the broad-spectrum protease inhibitor, chymostatin nor the aspartic protease inhibitor, pepstatin A, stimulated NSPC proliferation. Taken together, the results support the idea that NSPC proliferation is regulated by one or more cysteine proteases. Indeed, previous studies have shown that cysteine proteases

can ablate protein synthesis and thereby arrest cell proliferation (Chun et al 2002, Salmena et al 2003, Yadaiah et al 2013a, Yan et al 2001b), which also supports this view. On this basis, CysC may increase NSPC proliferation through acting as an inhibitor of cysteine protease activity.

Despite the plausibility of this hypothesis, no cysteine proteases were identified in the NSPC conditioned medium using the method of CysC-Sepharose affinity chromatography. Although CysC, E-64 and antipain stimulated NSPC proliferation at lower concentrations in this study (Fig. 4.1, A and B), they inhibit cysteine protease activities through different mechanisms. CysC binds adjacent to the catalytic center of cysteine proteases to block the substrate access, but without directly engaging with the active site of the cysteine protease (Rzychon et al 2004). E-64 and E-64D bind directly to the catalytic center of cysteine proteases (Feng et al 1996, Matsumoto et al 1999) by irreversibly modifying the thiol group of the cysteine protease to a thioether group. The inhibitory action of E-64 and E-64 D occurs through nucleophilic attack by the thiol group of the cysteine protease on the epoxide of the inhibitor (Hanada et al 1978a, Hanada et al 1978b, Matsumoto et al 1999, Tamai et al 1981). Importantly, E-64 does not interact with the functional thiol group of non-protease enzymes. The reversible cysteine/serine protease inhibitor, antipain also reversibly inhibits cysteine protease activities through the peptidyl aldehyde group binding covalently to cysteine proteases (Frommer et al 1979, Otto & Schirmeister 1997, Suda et al 1972). However, no matter whether these inhibitors affect the activity of cysteine protease directly or indirectly, inhibition of cysteine protease activities by CysC, E-64 and antipain, at least at low concentrations, led to increased NSPC proliferation (Fig. 4.1, A and B).

This argues that it is likely to be the cysteine protease activity itself that is important for the NSPC proliferation effect.

E-64, antipain and CysC are not able to cross the plasma membrane. Thus, it is likely that E-64 and CysC may affect NSPC proliferation by inhibiting an extracellular cysteine protease such as a cathepsin. Cathepsins are found to participate in protein degradation not only in the lysosome and secretory vesicles but also in the extracellular space (Green & Lund 2005, Hook et al 2012, Stoka et al 2005). Overproduction of cathepsins is usually accompanied by extracellular secretion of procathepsins that are processed or activated automatically once in acidic conditions or by extracellular proteases or polysaccharides (Beckman et al 2009, Mason & Massey 1992, Pungercar et al 2009, Vasiljeva et al 2005).

As cysteine cathepsins are proposed to have multiple biological roles including cell growth and death, as well as the production of biologically active peptides (Gopinath et al 2013, Goulet et al 2007, Hook et al 2012, Minokadeh et al 2010, Moles et al 2009, Nomura & Katunuma 2005, Veillard et al 2011), inhibitors that are able to block activation of procathepsin or to inhibit the activity of cysteine cathepsins may play a role in the regulation of cell proliferation. For example, a number of studies have indicated that cysteine cathepsins can degrade extracellular matrix (ECM) proteins to influence homeostasis of the ECM. Digestion of the ECM could exert an effect on NSPC proliferation (Gattazzo et al 2014, Hou et al 2003, Kurtz & Oh 2012, Maciewicz & Wotton 1991). For this reason, regulation of the activity of these extracellular cysteine cathepsins by inhibitors like E-64 or CysC may affect NSPC proliferation. As the serine protease inhibitor, antipain can increase NSPC

proliferation, and can also block activity of serine proteases, a role for serine proteases in NSPC proliferation cannot be ruled out (Xia et al 2013).

In this study, E-64 promoted NSPC proliferation, whereas two other synthetic analogues of E-64, namely E-64C and E-64D, exerted either a poor effect (E-64D) or no effect (E-64C) on NSPC proliferation (Fig. 4.1, C). Like E-64, E-64C is non-cell permeable, and preferentially affects activities of cysteine proteases in a similar way to E-64 (Matsumoto et al 1999, Matsumoto et al 1989). The possible explanation for the different effects of the E-64 analogues may be that alteration of the structure of E-64 may affect the ability of the compound to undertake nucleophilic attack on the epoxide (Matsumoto et al 1999, Matsumoto et al 1989). While E-64C and E-64D are known to inhibit a number of cysteine proteases in a manner similar to E-64, it is possible that the proteases responsible for NSPC proliferation may not be readily inhibited by the analogues.

E-64D is an ethyl ester form of E-64C and is cell membrane permeable. E-64D was found to slightly promote NSPC proliferation at a concentration of 0.3 μ M (Fig. 4.1, C). E-64D also contains the epoxide structure that is involved in the interaction within the active site of the cysteine protease in E-64 (Hanada et al 1978a, Hanada et al 1978b, Matsumoto et al 1989, Wilcox & Mason 1992). The reason why E-64D stimulated NSPC proliferation, but E-64C did not, is not clear. However, one possibility is that altering the molecular structure of E-64C to E-64D promotes an interaction between the inhibitor and the cysteine proteases. Therefore, E-64D may stimulate NSPC proliferation by acting on an extracellular cysteine protease (Hook et al 2011), as was observed with CysC, antipain and E64. Alternatively, addition of an

ethyl group onto E-64C increases the lipophilicity of the compound, which making E64D membrane permeable and capable of being transported into the cells. Thus it might be argued that this is an evidence for an action on an intracellular protease. If this was the case, it would be surprising to find that CysC, a membrane -impermeable inhibitor would also be active. It is more likely that the ethyl esterification of E-64C to E-64D affects the compound's affinity for the proteases. Or it may act by another undefined mechanism.

In this study, low concentrations of E-64, E-64D and antipain exhibited a stimulatory effect on NSPC proliferation while higher concentrations usually exerted an inhibitory effect on cell proliferation. CysC has also been reported to promote NSPC proliferation in a concentration-dependent manner, as high concentrations of CysC reportedly suppress NSPC proliferation (Taupin et al 2000). Therefore, low concentrations of cysteine protease inhibitors may block the activities of the cysteine protease that is responsible for NSPC proliferation. High concentrations of cysteine proteases inhibitors are more likely tend to excessively suppress cysteine proteases activities required for NSPC proliferation or maintenance (Kurtz & Oh 2012).

CysC affinity chromatography was used to further investigate which cysteine proteases might interact with CysC to regulate NSPC proliferation. However, no cysteine proteases were identified in this study. No proteins were identified in the region of 21-30 kDa in which many cysteine proteases are normally found. A band was clearly observed around 13-15 kDa in the lane 6 (Fig. 4.2) at a position corresponding to CysC. This band had the same molecular weight as CysC previously identified in NSPC conditioned medium (Chapter 3). This band was probably a

contaminant that had leaked from Sepharose resin and therefore it was not analyzed further. However, it cannot be concluded that CysC does not bind to a cysteine protease from this result. Firstly, the amount of extracellular cysteine proteases in the NSPC conditioned medium that interact with CysC may be too low to be characterized by silver staining. CysC may possibly bind to several cysteine proteases which are present at very low concentrations in the NSPC conditioned medium. Secondly, it is possible that the cysteine protease may bind to CysC but may not bind well to CysC-Sepharose. This could be due to Sepharose resin attaching to CysC close to the binding site for cysteine proteases. This possibility is quite likely given the fact that CysC is a relatively small protein. Unfortunately, there was not sufficient time to investigate these possibilities further.

Chapter 5

Role of APP in neural stem/progenitor cell differentiation

5.1 Introduction

Neural stem or progenitor cells (NSPCs) are able to self-renew and also retain the ability to give rise to the major cell types of the CNS including glial and neuronal lineages (Palmer et al 1999, Reynolds & Weiss 1992, Richards et al 1992). The investigation of NSPC biology is important not only because NSPC persist into adulthood, but also because of the possibility that NSPCs may be a therapeutic agent for the treatment of neurodegenerative diseases or other neurological disorders via replacement of lost, or injured cells (Hsieh 2012, Taylor et al 2013). Theoretically, NSPCs may provide an inexhaustible cell source for cell repair or tissue engineering in neurobiology (Gil-Perotin et al 2013). However, the process of NSPC differentiation is controlled and modulated by a variety of endogenous factors in vivo (Huang et al 2010, Katsimpardi et al 2014, Lopez-Juarez et al 2012, Roybon et al 2009a). Thus a good understanding of how NSPC behaviour is regulated is crucial if NSPCs are to be used as a source of neural cell types that can function properly integrate into the existing circuitry.

β -Amyloid protein ($A\beta$) plays a central role in AD pathogenesis (Nunan & Small 2000, Zheng & Koo 2011). $A\beta$ is produced by processing of the β -amyloid precursor protein (APP). However, the normal biological function of APP still remains elusive. Increased APP overexpression associated with neuronal differentiation has been observed during neuronal development (Clarris et al 1995, Hung & Selkoe 1994, Trapp & Hauer 1994). APP was reported to induce neural differentiation of pluripotent stem cells (Khandekar et al 2012). In addition, enhanced neuronal differentiation was also observed in APP overexpressing transgenic mice (Jin et al 2004a, Lopez-Toledano & Shelanski 2007, Yu et al 2009). However, studies on the

effect of APP deficiency found that embryonic stem cells from APP triple knockout mice still can differentiate into neurons both in vivo and vitro, indicating that APP expression does not play an essential role in the initiation of neuronal differentiation (Bergmans et al 2010). Nevertheless, delayed neuronal differentiation and development are also associated with decreased APP, APLP1 and APLP2 expression (Shariati et al 2013). Moreover, the soluble secreted fragments of APP, sAPP α and A β have been reported to drive NSPC differentiation, either towards an astrocytic or neuronal fate in vitro (Baratchi et al 2012, Chen & Dong 2009, Gakhar-Koppole et al 2008, Kwak et al 2006a, Lopez-Toledano & Shelanski 2007). Taken together, these findings suggest a role for APP in NSPC differentiation.

The study reported in Chapter 3 showed that NSPCs derived from APP transgenic mice (Tg2576) proliferate more rapidly than the corresponding wild type mice, while NSPCs obtained from APP KO mice proliferate less readily than their background strain of wild type mice (Hu et al 2013). The enhanced NSPC proliferation was induced by APP-mediated secretion of a protease inhibitor, CysC (Hu et al 2013). Neither A β nor sAPP α exerted an effect on proliferation.

To address the role of APP in NSPC differentiation, the differentiation of NSPCs derived from APP transgenic mice (Tg2576) and from APP knockout (APP KO) mice was examined. These studies showed that APP induces neuronal and astrocytic differentiation of NSPCs, but has no effect on oligodendrocyte differentiation. Although previous studies have suggested that sAPP α or A β induce NSPC differentiation (Heo et al 2007, Kwak et al 2006a, Lopez-Toledano & Shelanski 2007), in the present study, neuronal and astrocytic differentiation were not found to

be influenced by sAPP α or synthetic A β . Furthermore, CysC also did not induce NSPC differentiation.

5.2 Materials and methods

5.2.1 Materials used in this study are shown in Table 2.1.

5.2.2 NSPC differentiation

Neurospheres were prepared from Tg2576 mice and the corresponding background strain mice C57Bl/6 \times SJL, as well as APP KO mice and the corresponding background strain littermate mice C57Bl/6. NSPC cultures were prepared following the procedure described in chapter 3 section 3.2.1. After 7 days in culture, neurospheres were mechanically dissociated and the isolated cells were plated on poly- L- lysine coated coverslips in 24 -well plates at the density of 100,000 cells per well in proliferation medium (Table 2.2). The proliferation medium was removed after the cells were seeded on the coverslip, and then the cells were grown in a differentiation medium [(DMEM supplemented with 2% (v/v) B27, 100 units/ml penicillin, 100 units/ml streptomycin and 1% (v/v) heat - inactivated foetal calf serum (FCS)] and incubated for 5 or 14 days at 37°C in an atmosphere containing 5% CO₂. To test the effect of CysC, sAPP α , A β ₄₀ and A β ₄₂ on NSPC differentiation, cells were maintained in differentiation medium containing CysC, sAPP α , A β ₄₀ and A β ₄₂ for 5 days.

5.2.3 Immunocytochemistry

Cells were fixed with 4% (w/v) paraformaldehyde in phosphate-buffered saline (PBS) for 15 min, after which cells were permeabilized with 0.03% (v/v) Triton X-100 in PBS for 5 minutes, and then incubated in 10% (v/v) sheep serum in PBS for 1 hour to block non-specific binding sites. Fixed, permeabilized cells were double immunostained with a mouse IgG anti- β III tubulin antibody (1:1000 diluted in 10% (v/v) sheep serum in PBS) and a rabbit IgG anti-nestin antibody (1:5000 diluted in 10% (v/v) sheep serum in PBS), or a mouse IgG anti-GFAP antibody (1:3000 dilution in 10% (v/v) sheep serum in PBS) and a rabbit IgG anti-nestin antibody (1:5000 diluted in 10% (v/v) sheep serum in PBS) at 4°C overnight, respectively. After that the cells were incubated with a goat anti-mouse IgG conjugated to Alexa Fluor 488 or 594 (1:1000 diluted in 10% (v/v) sheep serum in PBS) or with goat anti-rabbit IgG conjugated to Alexa Fluor 488 or 594 (1:1000 diluted in 10% (v/v) sheep serum in PBS), and counterstained with 4', 6-diamidino-2-phenylindole DAPI at 1:10,000 dilution in PBS.

There was no need to permeabilize the fixed cells that were immunostained with a rabbit IgM anti-NG₂ antibody. In this case, the fixed cells were directly incubated in 10% (v/v) sheep serum in PBS for 1 hour to block non-specific binding sites and then incubated in rabbit anti-NG₂ in 10% (v/v) sheep serum in PBS at 4°C overnight. After that the cells were incubated with a rabbit goat anti-IgM conjugated to Alexa Fluor 594 (1:1000 diluted in 10% (v/v) sheep serum in PBS), then the cells were permeabilized with 0.03% (v/v) Triton X-100 in PBS for 5 minutes. The cells were blocked in 10% (v/v) sheep serum in PBS for 1 hour, and incubated with rabbit IgG anti-nestin antibody (1:5000 diluted in 10% (v/v) sheep serum in PBS) at 4°C

overnight. Finally, the cells were incubated with goat anti-rabbit IgG conjugated to Alexa Fluor 488 (1:1000 diluted in 10% (v/v) sheep serum in PBS) and counterstained with DAPI (at 1:10,000 dilution in PBS). The cells were mounted in DAKO mounting medium and allowed to dry overnight before imaging.

5.2.4 Image capture and data analysis

Images of fluorescently labelled cells were captured under the 20x objective using a Nikon Ti-E microscope (Nikon corporation, Tokyo, Japan). To quantify the amount of β III tubulin⁺, GFAP⁺, and NG2⁺ fluorescence relative to DAPI⁺ fluorescence, 9 random fields were selected from 1 coverslip and 3 coverslips were analyzed for each group in every experiment. There were approximately 100 cells in each field. Images were analysed using ImageJ version 1.46 software and statistical analysis was performed with GraphPad Prism version 5.04 software. Data were tested by Student's *t* test or one-way ANOVA. Post hoc comparisons were analyzed using Tukey's test. Differences were considered statistically significant when the probability (*p*) of the null hypothesis < 0.05. Data values are shown as the means \pm S.E. All results were derived from at least three independent experiments in which cells were derived from at least three different mice of the same strain.

5.3 Results

5.3.1 Differentiation of NSPC

Previous in vitro studies on NSPC proliferation found that NSPCs derived from APP KO mice proliferated more slowly than NSPCs obtained from wild type (WT) mice of the background strain (C57BL/6), while NSPCs from APP overexpressing (Tg2576)

mice proliferated more rapidly than NSPC from corresponding wild type (C57BL/6 × SJL) mice (Hu et al 2013). This result showed that APP expression exerts an effect on NSPC proliferation.

To examine the possibility that APP also influences NSPC differentiation, the neuronal differentiation of NSPCs prepared from postnatal day 0 APP KO and the corresponding WT (C57BL/6) mice, along with Tg2576 mice and the corresponding background strain WT (C57BL/6 × SJL) mice was examined. After 5 or 14 days in culture, the cells were fixed and labelled with neuronal, astrocytic and oligodendrocyte progenitor (OPC) markers, β III tubulin, GFAP and NG2, respectively.

The proportion of β III tubulin⁺, GFAP⁺ and NG2⁺ positive cells to total viable cells in the culture was determined. Three independent experiments were carried out and 21 fields were randomly selected from three separate cultures from each experiment with approximately 100 viable cells in each field.

The experiment showed that a relatively smaller percentage of APP KO NSPCs were stained with β III tubulin⁺ than WT NSPCs after both 5 and 14 days in culture (Fig. 5.1, A and D). In contrast, there was a higher proportion of β III tubulin⁺ neurons in the Tg2576 NSPC cultures than in the WT NSPC culture after 5 and 14 days (Fig. 5.1, A and D). The percentage of GFAP⁺ cells in the Tg2576 NSPC cultures was similar to the percentage of GFAP⁺ astrocytes in the WT culture after 5 days (Fig. 5.1, B and E). However, after 14 days, the proportion of GFAP⁺ cells in the Tg2576 NSPC cultures was significantly higher than the percentage of cells with GFAP positive staining in




Figure 5.1 Effect of APP on NSPC differentiation. Neuronal, astrocytic and oligodendrocyte progenitor (OPC) differentiation was detected by expression of their specific marker. NSPCs were grown on poly-L-lysine coated coverslips in differentiation medium, and maintained for 5 or 14 days respectively before immunostaining for β III tubulin (panel A, neuronal marker), GFAP (panel B, astrocytic marker), or NG2 (panel C, oligodendrocyte progenitor marker).




Figure 5.1 (Cont'd) The percentage of cells with β III tubulin, GFAP or NG2 expression was calculated by counting the number of cells with β III tubulin, GFAP and NG2 positive staining and dividing by the total number of cells stained with DAPI, respectively. Panels D - F show quantitation of immunofluorescence staining (A - C). The data were collected from three independent experiments, values are means. Two tailed t tests: panel D, C57BL/6 versus APP KO : $P < 0.001$, $n = 45$ (5 days) and $P < 0.001$, $n = 62$ (14 days); C57BL/6×SJL versus Tg2576, $P < 0.01$, $n = 61$ (5 days) and $P < 0.05$, $n = 58$ (14 days); Panel E: C57BL/6 versus APP KO, $P < 0.001$, $n = 40$ (5 days) and $P < 0.0001$, $n = 60$ (14 days); C57BL/6×SJL versus Tg2576, $P > 0.05$, $n = 63$ (5 days); $P < 0.05$, $n = 63$ (14 days); Two way ANOVA: Panel F: C57BL/6 versus APP KO, $P < 0.05$, $n = 60$ (5 days) and $P < 0.05$, $n = 59$ (14 days); C57BL/6×SJL versus Tg2576, $P > 0.05$, $n = 55$ (5 days) and $P > 0.05$, $n = 59$ (14 days).

the WT cultures (Fig. 5.1, B and E). A lower percentage of GFAP⁺ cells was observed in APP KO NSPC cultures compared to the WT culture both after 5 days and 14 days (Fig. 5.1, B and E).

The percentage of NG2⁺ cells in WT cultures was higher than in the APP KO cultures after 5 days, but after 14 days the proportion of NG2⁺ cells was lower in the WT culture than in the APP KO culture after 14 days (Fig. 5.1, C and F). However, there was no difference in the proportion of NG2⁺ cells between the Tg2576 and the WT cultures after 5 or 14 days.

5.3.2 CysC is not involved in APP induced differentiation

The studies reported in Chapter 2 showed that APP-induced NSPC proliferation was due to secretion of CysC. For this reason, the possibility that CysC is a mediator of APP-induced NSPC differentiation was examined. NSPCs from APP KO and the corresponding background strain WT (C57BL/6) mice were prepared and incubated in differentiation medium containing 0, 100 or 200 ng/ml of CysC. After 5 days, there was no difference in the proportion of β III tubulin⁺ neurons and GFAP⁺ astrocytes in any of the treatment groups (Fig. 5.2, A-F). This indicated that CysC did not influence NSPC differentiation into neurons or astrocytes and it suggested that the effects of APP on NSPC differentiation were not due to CysC secretion.

5.3.3 sAPP α , A β ₄₀ or A β ₄₂ do not influence neuronal and astrocytic differentiation

As NSPC differentiation was not influenced by CysC, the next question was whether differentiation was influenced by sAPP α , or by A β ₄₀ or A β ₄₂ which are the major secreted fragments of APP. Indeed, a few studies have reported sAPP α or A β may

exert a neurotrophic effect on neural stem cell differentiation (Chen & Dong 2009, Demars et al 2011, Fonseca et al 2013, Freude et al 2011). In order to explore the effect of sAPP α , A β ₄₀ and A β ₄₂ on NSPC differentiation, APP KO NSPCs were grown in differentiation medium containing sAPP α at a concentration of 10 nM which was previously reported to stimulate NSPC proliferation (Demars et al 2011), or A β ₄₀ or A β ₄₂ at a concentration of 1 μ M which has been suggested to promote neuronal or astrocytic differentiation (Chen & Dong 2009). Controls for the corresponding vehicle were also included. The proportion of cells with β III tubulin⁺ neurons, GFAP⁺ astrocytes and NG2⁺ OPCs (Fig. 5.3, A - C) was then calculated. However, after 5 days there was no difference observed in the percentage of cells positive for β III tubulin, GFAP and NG2 after treatment with sAPP α (Fig. 5.3, C and F) A β ₄₀ (Fig. 5.3, A and D) or A β ₄₂ (Fig. 5.3, B and E). It was concluded that the effect of APP on NSPC neuronal or astrocytic differentiation was probably not due to sAPP α , A β ₄₀ or A β ₄₂.

5.4 Discussion

The study reported in this chapter shows that APP can influence NSPC differentiation into neurons and astrocytes but plays little role in oligodendrocyte (OPC) differentiation. However, the data suggest that APP-induced neuronal and astrocytic differentiation is not due to secretion of CysC, nor due to the proteolytic products of APP processing, sAPP α , and A β ₄₀ or A β ₄₂.

Increased APP expression has been reported during neuronal differentiation (Hung et al 1992), which suggests a potential role for APP in neuronal fate development. Additionally, enhanced neuronal differentiation has been reported in young transgenic




Figure 5.2 Effect of cystatin C on neuronal and astrocytic differentiation. NSPCs were grown in differentiation medium containing cystatin C for 5 days and then the cells were fixed and labelled for specific markers of neurons (β III tubulin) and astrocytes (GFAP). Panel A and B show representative immunofluorescence images.




Figure 5.2 (Cont'd) Panel C - F show quantitation of immunofluorescence staining (A and B). The percentage of β III tubulin and GFAP expression in culture was determined as a ratio of the number of β III tubulin (panel A, green) or GFAP - positive cells (panel B, red) to the total number of cells stained with DAPI. In panels C, D, E and F, values are means \pm SEM and shown as % of the control (0 ng/ml cystatin C) ($P > 0.05$, ANOVA with post-hoc Tukey's test).




Figure 5.3 Effect of $A\beta_{40}$, $A\beta_{42}$ and sAPP α on NSPC differentiation. NSPCs from APP KO mice were differentiated for 5 days in culture and then treated with $A\beta_{40}$ (1 μ M), $A\beta_{42}$ (1 μ M) or sAPP α (100 nM). The cells were fixed and labelled for markers of neurons (β III tubulin), astrocytes (GFAP) and OPCs (NG2), respectively. Cells were counterstained with DAPI.




Figure 5.3 (Cont'd) Panel D - F show quantitation of immunofluorescence staining in panels A - C. The proportion of cells with β III tubulin (green, panel A), GFAP (red, panel B), or NG2 (green, panel C) staining was calculated as a ratio of the number of cells labelled with DAPI, respectively. Panels D, E and F show the effect of Aβ₄₀, Aβ₄₂ and sAPPα respectively versus their corresponding vehicle controls. Values are means ± SEM and shown as % of the corresponding vehicle control ($P > 0.05$, $n = 63$).

mice overexpressing APP (Lopez-Toledano & Shelanski 2007), a result consistent with the view that expression of APP may participate in neuronal differentiation (Fig. 5.1, A and D).

A few studies have reported that APP-induced multipotent NSPC differentiation is due to the major soluble form of APP, sAPP α (Demars et al 2011, Freude et al 2011, Kwak et al 2006a). However, this result was not observed in the current study, which found that recombinant sAPP α had no effect on NSPC differentiation (Fig. 5.2, C and F). Nonetheless, Kwak et al (2006) reported that the secreted form of APP (e.g sAPP α) can enhance glial differentiation but not neuronal differentiation in vivo. However, in the present study, NSPCs treated with sAPP α did not display increased astrocytic or OPC differentiation. The reasons for the difference are unclear. sAPP α may exert different effects in vitro and in vivo. Indeed, an in vivo situation is a much more complicated system than one in vitro. Therefore the observed phenomena, such as enhanced glial differentiation in vivo, could be attributed to a series of complex signal pathways, not just to the infusion of sAPP α . An alternative reason is that cells in the different studies did not come from the same sources, and sAPP α may exert different effects on different types of neural stem cells (Freude et al 2011). Only further studies will be able to address this issue.

While this study found that APP causes neural differentiation, this was not due to A β production, because the two major forms of A β did not affect NSPC differentiation. Several studies have reported that A β peptides have neurotrophic effects and that they can stimulate NSPC differentiation (Chen & Dong 2009, Fonseca et al 2013, Haughey et al 2002, Lopez-Toledano & Shelanski 2004, Lopez-Toledano & Shelanski 2007,

Sotthibundhu et al 2009). Nonetheless, other groups reported the opposite results, i.e. that A β has an inhibitory effect on neurogenesis, neural differentiation or NSPC fate determination (Haughey et al 2002, He et al 2013). Once again, the reason for the difference between these studies is unclear. However, there are various forms of A β_{40} and A β_{42} including different aggregated forms of A β . It is possible that the forms of A β used in the present study were not the same as the forms that induced differentiation in the literature (Chen & Tang 2006, He et al 2013). For instance, He et al (2013) found that A β_{42} oligomers induce NSPC senescence in adult hippocampus, while another two groups observed an increase in NSPC differentiation following A β_{40} or A β_{42} peptide treatment (Chen & Dong 2009, Fonseca et al 2013). Methods by which A β is prepared such as the solvent that is used to dissolve A β may lead to changes in A β conformation (Soto et al 1994, Zagorski & Barrow 1992). A β contains a completely hydrophobic C-terminal region that promotes β strand structure, and an N-terminal domain, that allows formation of different secondary structures (Soto et al 1994). However, the N-terminal domain of A β can exist either in an α helical or in β strand conformation based on the environmental conditions (pH, hydrophobicity of surrounding molecules). Therefore altered A β conformation may also influence NSPC behavior (Chen & Dong 2009, Soto et al 1994, Sotthibundhu et al 2009).

Another possibility that may explain why there are differences about the effect of A β on NSPC differentiation is that the studies reported here were carried out on NSPCs derived from a different model. The response of these NSPCs may be variable depending on the system used. One study based using animal models reported a stimulatory effect of A β on neuronal differentiation of NSPCs in the SVZ

(Sotthibundhu et al 2009), while another study examining the effect of A β on NSPCs from the SVZ showed an inhibitory effect on neuronal differentiation (Haughey et al 2002). The reason for the discrepancy is unclear. To conclude, further studies on the effect of A β on NSPCs behaviour are required.

CysC has been reported to promote NSPC proliferation (Dahl et al 2003, Taupin et al 2000) and NSPC proliferation is regulated by APP through CysC (Hu et al 2013). A few papers have been reported that CysC can influence NSPC differentiation (Kato et al 2006, Kumada et al 2004). In one, CysC induced ES cells to differentiate into NSPCs and β III tubulin⁺ cells in this study (Kato et al 2006). CysC can also activate the GFAP promoter and induce astro-glial differentiation during mouse brain development (Kumada et al 2004). However, the result of this study did not support the idea that CysC can influence adult NSPC differentiation. It is possible that CysC may affect the development of ES cells into neural stem cells, as reported previously (Kato et al 2006), but it may have no effect on adult NSPC development and differentiation.

The molecular basis by which APP induces differentiation remains unclear and needs for future investigation. Studies done by the author in collaboration with Dr Marta Bolos found that an increased expression of a proneural basic helix-loop-helix (bHLH) transcription factor, neurogenin 2 (Ngn 2) is likely to be a mediator of this effect (Bolos et al 2014). In that study, transfection of Ngn2 into APP KO NSPCs stimulated increased neuronal differentiation of NSPCs, whereas knocking down Ngn2 expression in WT NSPCs led to a decrease in neuronal differentiation (Bolos et al 2014). Therefore, APP overexpression may cause up-regulation of Ngn2 expression and thereby induce neuronal differentiation.

Ngn2 is unlikely to be involved in glial differentiation because neurogenins including Ngn2 are reported to promote neuronal differentiation, but inhibit glial differentiation (Sun et al 2001). APP may mediate glial differentiation via a BMP family signalling pathway (Kwak et al 2014), monocyte chemoattractant protein-1 expression (Vrotsos et al 2009) or interleukin-6 cytokine release (Kwak et al 2006b, Ringheim et al 1998). However, the mechanism by which APP modulates Ngn2 expression and NSPC differentiation remains unclear.

Chapter 6 Discussion

The neuropathology of AD was first described by Alois Alzheimer in 1907 (Alzheimer 1907). Despite intensive studies, the mechanism of AD development still remains unclear. There are no effective treatments to prevent, or even delay the progression of AD. AD is characterized by the deposition of amyloid plaques and neurofibrillary tangles in the brain (Kidd 1964, Terry 1963). A β is the major component of the amyloid plaques (Masters et al 1985), and is proposed to be the key causative agent of AD (Hardy & Selkoe 2002, Small et al 1992). A chronic imbalance of production and clearance of A β has been proposed to trigger progression of AD (Hardy & Selkoe 2002).

A β is produced by sequential proteolytic cleavage of the β - amyloid precursor protein (APP) (Kang et al 1987). According to the amyloid hypothesis of AD, several therapeutic approaches have proposed that are mainly aimed at reducing A β production via alteration of APP processing. Indeed, many compounds that target APP processing to lower A β production have already been studied in clinical trials (Small et al 2004). However, the biological function of APP is still unknown. A major motivation for this study was to understand the function of APP. APP has been intensively studied for its relationship with AD. Many second generation AD therapies are focusing on decreasing A β production through inhibition of APP proteolytic processing. Therefore, a better understanding of the function of APP is essential to develop approaches for AD treatment, as it is possible that altering APP metabolism may also alter its function and thereby induce toxicity.

Several studies have reported enhanced NSPC proliferation in APP overexpressing transgenic mice (Jin et al 2004a, Lopez-Toledano & Shelanski 2007, Yu et al 2009) as

well as in the brain of patients with AD (Jin et al 2004c). In addition, increased APP expression has been observed at early stages of neuronal differentiation, maturation and fate specification during embryonic development (Clarris et al 1995, Salbaum & Ruddle 1994). However, other studies have reported disrupted NSPC proliferation or differentiation in APP transgenic mice (Haughey et al 2002, Mu & Gage 2011). Several cytokines that are released during inflammation in the AD brain have been found to cause a deficit in NSPC proliferation or differentiation (Keohane et al 2010, Kohman & Rhodes 2013, Monje et al 2003). Thus, there is a possibility that the disease-related inflammation in mice that overexpress APP, and which produce A β , may be the cause of the altered neurogenesis in these studies.

It was hypothesized that APP is involved in the regulation of adult NSPC proliferation and differentiation. The studies presented in this thesis were aimed at exploring the potential role of APP in NSPC proliferation and differentiation by examining NPSC proliferation and differentiation in APP overexpressing and APP knockout NSPC culture and by further investigating the mechanisms involved in this action. Neurogenesis occurs not only in development but also persists into adulthood with active NSPCs continuously undergoing proliferation and differentiation in neurogenic niches (Alvarez-Buylla & Garcia-Verdugo 2002, Eriksson et al 1998). Furthermore, adult neurogenesis was reported to be associated with discrimination, learning and memory (Deng et al 2009, Gheusi et al 2000, Rochefort et al 2002). Therefore, it is essential to identify proper therapeutic targets for AD treatments without interfering with the normal function of APP.

The in vitro NSPC culture studies reported in chapter 3 and 5 indicated that APP may

participate in the regulation of NSPC proliferation and differentiation. In the study presented here, increased NSPC proliferation was observed in NSPCs derived from APP transgenic (Tg2576) mice while decreased NSPC proliferation was observed in cultures from APP KO mice (Chapter 3). However, enhanced NSPC proliferation was affected neither by the secreted soluble form of APP, sAPP α , nor by the β -amyloid protein, A β (Chapter 3). Instead, these studies demonstrated that APP-induced NSPC proliferation is mediated by secretion of the cysteine protease inhibitor, CysC. CysC was identified in the NSPC conditioned medium, and the secreted CysC also was found to also stimulate NSPC proliferation in vitro (Taupin et al 2000, Tavera et al 1992). This finding was supported by the observation that recombinant CysC could promote NSPC proliferation.

CysC is secreted and targeted extracellularly through the secretory system. APP overexpression has been reported to enhance vesicle exocytosis in PC12 cells (Lee et al 2008a). Thus, increased CysC secretion might be due to increased vesicle exocytosis of CysC into the cell culture media. In support of this idea, the studies presented in Chapter 3 showed that lower intracellular levels of CysC were detected in cell lysates of Tg2576 NSPCs compared to the corresponding WT NSPCs. In addition, the studies found higher CysC secretion levels in conditioned medium from APP overexpressing (Tg2576) cells, while lower levels of secreted CysC were present in APP KO conditioned medium. Surprisingly, there was no increase in CysC mRNA levels in Tg2576 cells compared to the corresponding wild type control cells. This result may suggest the possibility that APP overexpression in Tg2576 cells reaches a saturation point, so increased APP expression does not further increase CysC mRNA transcription. The overexpression of APP may play little role in CysC transcription,

but it may enhance CysC secretion in Tg2576 cells. Moreover, studies based on real-time PCR indicated that there were lower *CysC* mRNA levels in APP KO NSPCs compared to the corresponding wild type NSPCs (Chapter 3). This suggests that lower expression levels of APP, such as those that occur endogenously, may be able to regulate CysC transcription.

In agreement with this study, which showed increased CysC in APP NSPCs, APP overexpression has previously been reported to be associated with increased CysC levels in APP transgenic mice (Steinhoff et al 2001). In addition, CysC was found to colocalize with A β in vascular amyloid and in amyloid plaque cores of patients with AD and other neurodegenerative diseases (Haan et al 1994, Levy et al 2001, Maruyama et al 1990). Co-deposition of CysC and amyloid plaques (Fig. 6.1) was also observed in non-demented aged individuals (Levy et al 2001). Moreover, enhanced amyloid plaque degradation occurs in APP transgenic mice with CysC genetically deleted (Sun et al 2008), despite CysC having been shown to suppress amyloid deposition in AD mouse models (Kaeser et al 2007, Mi et al 2007). However, CysC deletion in human APP transgenic mice was reported to significantly lower A β in another study (Wang et al 2012). Therefore, a complex interplay between APP expression, A β plaque, CysC production and amyloid deposition may occur in AD. This needs further examination in future studies.

CysC null mice exhibited a decreased capability for neurogenesis (Pirttila et al 2005), which supports a role for CysC in NSPC proliferation. Thus, this suggests the possibility that manipulation of CysC expression or secretion may facilitate NSPC proliferation and thereby possibly enlarge the pool of stem cell treatments. Besides,

CysC is an endogenous inhibitor of cysteine proteases and it may regulate NSPC proliferation through inhibition of a protease. Indeed, some proteases have also been found to be associated with NSPC proliferation and differentiation (de Azevedo-Pereira et al 2011, Egberts et al 2004, Ekdahl et al 2001, Joyce et al 2004, Santos et al 2012, Watkinson 1999, Yadaiah et al 2013a). However, further studies of the mechanism by which cysteine protease activity is involved in NSPC proliferation or differentiation is required, and this may provide a new insights and understanding of stem cell biology.

The specific cysteine protease inhibitor, E-64, was also found to promote NSPC proliferation (Chapter 4). This result supports the idea that CysC may act as a protease inhibitor to stimulate cell growth, although CysC affinity chromatography did not purify any cysteine proteases that bind to CysC. E-64 is a non-cell membrane permeable cysteine protease, thus it may inhibit an extracellular cysteine protease to stimulate NSPC proliferation (Chapter 4). CysC was shown to inhibit several extracellular cathepsins, including cathepsins B, H, K, L, and S (Bernstein et al 1996), but it was inactivated by cathepsin D (Abrahamson et al 1991, Lenarcic et al 1991). Some cathepsins have been reported to induce neuronal apoptosis, therefore inhibition of these cathepsins may increase cell survival or perhaps cell growth (Boland & Campbell 2004, Kingham & Pocock 2001).

An imbalance between production and metabolism of cysteine proteases and their endogenous inhibitor may be involved in amyloid pathology in AD (Nakamura et al 1991, Sun et al 2008). Cathepsin B has been reported to effectively reduce A β , because inhibition of cathepsin B in mice overexpressing APP results in increased

plaque load (Mueller-Steiner et al 2006, Sun et al 2008, Wang et al 2012). In addition co-localization of cathepsins and senile plaques in the brain of AD patients has been reported (Cataldo & Nixon 1990). Furthermore, cathepsins B and L have also been implicated in APP processing (Klein et al 2009). These studies reveal a correlation between cysteine proteases and amyloid deposition. Therefore, further investigation into CysC or E-64 bound by which cysteine protease inhibitors (Chapter 4) may not only contribute to direct NSPC proliferation for stem cell therapy, but also contribute to modulate A β degradation for AD treatment.

APP also promoted enhanced neuronal and astrocytic differentiation of NSPCs. However, this effect was not caused by CysC, A β or sAPP α (Chapter 5). This result is consistent with another study from our group showing that APP overexpression can promote neuronal differentiation, and that APP-induced neuronal differentiation is regulated by the proneural basic helix-loop-helix (bHLH) transcription factor, neurogenin 2 (Ngn2) (Bolos et al 2014). Ngn2 plays an important role in driving neuronal differentiation during neurogenesis (Ali et al 2011, Nieto et al 2001), and Ngn2 alone is sufficient to improve differentiation of embryonic stem cells into mature neurons (Thoma et al 2012). Thus, this suggests that APP-induced neuronal differentiation of NSPCs is via modulation of Ngn2 expression (Bolos et al 2014). However, the basis of how APP activates Ngn2 transcription is still unclear.

As the extracellular fragments of APP-A β and sAPP α were not found to promote NSPC proliferation, this suggests that the intracellular domain of APP (AICD) liberated by γ - secretase cleavage of APP processing is involved in APP induced CysC gene expression (Fig 6.1). AICD has been reported to interact with nuclear




Figure 6.1 Model of the possible role of APP in NSPC proliferation and differentiation. Briefly, the APP intracellular domain (AICD) is released by APP processing (1) and then interacts with an adaptor protein to form a complex (2). The AICD- adaptor protein complex translocates into the nucleus and possibly regulates gene transcription (3) such that of *CysC* (4) or genes that relate to NSPC differentiation e.g Ngn2 (5). Secreted *CysC* (6) may either modulate cysteine protease activity to promote NSPC proliferation (8) or co-deposit with Aβ (9). AICD may also facilitate NSPC differentiation by activating the transcription of genes related to NSPC differentiation (7).

adaptor proteins to form complexes that translocate into the nucleus, where they regulate gene expression (Cao & Sudhof 2001, Cao & Sudhof 2004, Dawkins & Small 2014, Giliberto et al 2008, Kimberly et al 2005, von Rotz et al 2004, Xu et al 2007). However, AICD may negatively affect NSPC proliferation, as decreased NSPC proliferation has been observed in APP KO mice expressing AICD (Ghosal et al 2010). Nevertheless, NSPC proliferation remained normal at 6 weeks of age but was reduced at 3 months in APP KO mice expressing AICD (Ghosal et al 2010). Neuro inflammation has been suggested as a possible explanation for the effect of AICD on NSPC proliferation, as AICD has been reported to impair NSPC proliferation via inducing neuroinflammation (Ghosal et al 2010). The involvement of inflammation is supported by the finding that anti-inflammatory agents can rescue defective NSPC proliferation (Ghosal et al 2010). Thus, the decreased NSPC proliferation in AICD transgenic mice may be due to neuroinflammation (Ghosal et al 2009, Ryan & Pimplikar 2005, Vogt et al 2011). The impaired NSPC proliferation in APP null mice expressing AICD may be due to activation of glycogen synthase kinase-3 β (GSK-3 β), a protein involved in cell apoptosis (Ghosal et al 2010). However, GSK-3 β mRNA and protein levels were not altered in these AICD transgenic mice (Ghosal et al 2010, Hetman et al 2000, Ryan & Pimplikar 2005). Therefore, the mechanism of the effect on NSPC proliferation is unclear (Ghosal et al 2010). It is possible that CysC changes the cleavage of endogenous APP to AICD, which further affects NSPC proliferation and differentiation. Nonetheless, the current study provides evidence that CysC is downstream of APP (Chapter 3). However, whether AICD plays a role in modulation of CysC transcription is not yet confirmed and will need further study.

In addition, the extracellular domain fragments of APP, namely A β and sAPP α , were not found to promote neuronal or astrocytic differentiation in the present study. Thus this supports the idea that the intracellular domain of APP may be involved in mediating this effect, via activation of gene transcription (Fig 6.1). AICD may be a transcription regulating factor (Cao & Sudhof 2001, Cao & Sudhof 2004, Giliberto et al 2008, Kimberly et al 2005, von Rotz et al 2004, Xu et al 2007). For example, AICD has been reported to facilitate the nerve growth factor response in neuronal cell differentiation and promote neurite outgrowth as well as to improve axonal elongation and path finding (Ryan & Pimplikar 2005, Zhou et al 2012). Nevertheless, other studies indicate that AICD suppresses neuronal specification of NSPCs or induces neuronal specific apoptosis (Ohkawara et al 2011, Shu et al 2015). However, APP KO mice that overexpress AICD exhibited normal neuronal differentiation of NSPCs (Ghosal et al 2010). Therefore, whether AICD functions as a gene transcriptional factor to regulate expression of genes controlling NSPC differentiation needs further investigation.

In summary, the main finding of this study was that APP overexpression induced NSPC proliferation and differentiation (Chapter 3 and 5), and that APP may play a stimulatory role in adult NSPC neurogenesis. For this reason, AD therapies based on altering APP processing may have to be considered carefully in order to not interfere with any normal physiology related to APP function in the human brain. In addition, APP-induced neuronal differentiation may be mediated via *Ngn2* transcription (Bolos et al 2014), whereas the APP induced NSPC proliferation was found to be due to secretion of the cysteine protease inhibitor CysC. It is possible that CysC is released from the cells by non – secretory mechanism (i.e following cell death). However,

previous studies have shown that CysC is actively secreted (Taupin et al 2000). Thus, the presence of CysC in the culture medium is likely to be due to active secretion. This idea is further supported by the observation that most of the CysC was in the culture medium rather than in the cell lysate fraction. Furthermore, low concentrations of other cysteine protease inhibitors such as E-64 and antipain also stimulated NSPC proliferation in this study. As modulation of cysteine proteases activity is suggested to be related to amyloid pathology in AD (Klein et al 2009, Sun et al 2008, Wang et al 2012), further investigations of the molecular basis of APP's involvement with NSPC proliferation and differentiation, and the role of CysC, may not only contribute to the development of new stem therapies for neurological diseases such as AD, but may also provide an understanding of the underlying causes of neurodegeneration.

Reference:

- Ables JL, Breunig JJ, Eisch AJ, Rakic P. 2011. Not(ch) just development: Notch signalling in the adult brain. *Nat Rev Neurosci* 12: 269-83
- Ables JL, Decarolis NA, Johnson MA, Rivera PD, Gao Z, et al. 2010. Notch1 is required for maintenance of the reservoir of adult hippocampal stem cells. *J Neurosci* 30: 10484-92
- Abrahamson M. 1994. Cystatins. *Methods Enzymol* 244: 685-700
- Abrahamson M, Barrett AJ, Salvesen G, Grubb A. 1986. Isolation of six cysteine proteinase inhibitors from human urine. Their physicochemical and enzyme kinetic properties and concentrations in biological fluids. *J Biol Chem* 261: 11282-9
- Abrahamson M, Buttle DJ, Mason RW, Hansson H, Grubb A, et al. 1991. Regulation of cystatin C activity by serine proteinases. *Biomed Biochim Acta* 50: 587-93
- Abramov E, Dolev I, Fogel H, Ciccotosto GD, Ruff E, Slutsky I. 2009. Amyloid-beta as a positive endogenous regulator of release probability at hippocampal synapses. *Nature neuroscience* 12: 1567-76
- Access Economics. 2009. keeping Dementia front of mind: Incidence and prevalence 2009-2050. *Report for Alzheimer's Australia*
- Adachi K, Mirzadeh Z, Sakaguchi M, Yamashita T, Nikolcheva T, et al. 2007. Beta-catenin signaling promotes proliferation of progenitor cells in the adult mouse subventricular zone. *stem cells* 25: 2827-36
- Adlard PA, Cherny RA, Finkelstein DI, Gautier E, Robb E, et al. 2008. Rapid restoration of cognition in Alzheimer's transgenic mice with 8-hydroxy quinoline analogs is associated with decreased interstitial Abeta. *Neuron* 59: 43-55
- Adolfsson O, Pihlgren M, Toni N, Varisco Y, Buccarello AL, et al. 2012. An effector-reduced anti-beta-amyloid (Abeta) antibody with unique abeta binding properties promotes neuroprotection and glial engulfment of Abeta. *J Neurosci* 32: 9677-89
- Agasse F, Nicoleau C, Petit J, Jaber M, Roger M, et al. 2007. Evidence for a major role of endogenous fibroblast growth factor-2 in apoptotic cortex-induced subventricular zone cell proliferation. *Eur J Neurosci* 26: 3036-42
- Aguirre A, Dupree JL, Mangin JM, Gallo V. 2007. A functional role for EGFR signaling in myelination and remyelination. *Nature neuroscience* 10: 990-1002
- Aguirre A, Rubio ME, Gallo V. 2010. Notch and EGFR pathway interaction regulates neural stem cell number and self-renewal. *Nature* 467: 323-7
- Aguirre AA, Chittajallu R, Belachew S, Gallo V. 2004. NG2-expressing cells in the subventricular zone are type C-like cells and contribute to interneuron generation in the postnatal hippocampus. *J Cell Biol* 165: 575-89
- Aguzzi A, Barres BA, Bennett ML. 2013. Microglia: scapegoat, saboteur, or something else? *Science* 339: 156-61
- Ahmed S, Gan HT, Lam CS, Poonepalli A, Ramasamy S, et al. 2009. Transcription factors and neural stem cell self-renewal, growth and differentiation. *Cell Adh Migr* 3: 412-24

- Ahmed S, Reynolds BA, Weiss S. 1995. BDNF enhances the differentiation but not the survival of CNS stem cell-derived neuronal precursors. *J Neurosci* 15: 5765-78
- Ahn K, Shelton CC, Tian Y, Zhang X, Gilchrist ML, et al. 2010. Activation and intrinsic gamma-secretase activity of presenilin 1. *Proc Natl Acad Sci U S A* 107: 21435-40
- Ahn S, Joyner AL. 2005. In vivo analysis of quiescent adult neural stem cells responding to Sonic hedgehog. *Nature* 437: 894-7
- Aisen PS. 2002. The potential of anti-inflammatory drugs for the treatment of Alzheimer's disease. *Lancet Neurol* 1: 279-84
- Akers KG, Martinez-Canabal A, Restivo L, Yiu AP, De Cristofaro A, et al. 2014. Hippocampal neurogenesis regulates forgetting during adulthood and infancy. *Science* 344: 598-602
- Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, et al. 2000. Inflammation and Alzheimer's disease. *Neurobiol Aging* 21: 383-421
- Al-Bassam J, Ozer RS, Safer D, Halpain S, Milligan RA. 2002. MAP2 and tau bind longitudinally along the outer ridges of microtubule protofilaments. *J Cell Biol* 157: 1187-96
- Alexson TO, Hitoshi S, Coles BL, Bernstein A, van der Kooy D. 2006. Notch signaling is required to maintain all neural stem cell populations--irrespective of spatial or temporal niche. *Dev Neurosci* 28: 34-48
- Alfonso J, Le Magueresse C, Zuccotti A, Khodosevich K, Monyer H. 2012. Diazepam binding inhibitor promotes progenitor proliferation in the postnatal SVZ by reducing GABA signaling. *Cell Stem Cell* 10: 76-87
- Ali F, Hindley C, McDowell G, Deibler R, Jones A, et al. 2011. Cell cycle-regulated multi-site phosphorylation of Neurogenin 2 coordinates cell cycling with differentiation during neurogenesis. *Development* 138: 4267-77
- Alley GM, Bailey JA, Chen D, Ray B, Puli LK, et al. 2010. Memantine lowers amyloid-beta peptide levels in neuronal cultures and in APP/PS1 transgenic mice. *Journal of neuroscience research* 88: 143-54
- Allinquant B, Hantraye P, Mailleux P, Moya K, Bouillot C, Prochiantz A. 1995. Downregulation of amyloid precursor protein inhibits neurite outgrowth in vitro. *J Cell Biol* 128: 919-27
- Altman J, Bayer SA. 1990. Mosaic organization of the hippocampal neuroepithelium and the multiple germinal sources of dentate granule cells. *J Comp Neurol* 301: 325-42
- Altman J, Das GD. 1965. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol* 124: 319-35
- Alvarez-Buylla A, Garcia-Verdugo JM. 2002. Neurogenesis in adult subventricular zone. *J Neurosci* 22: 629-34
- Alves da Costa C, Sunyach C, Pardossi-Piquard R, Sévalle J, Vincent B, et al. 2006. Presenilin-dependent gamma-secretase-mediated control of p53-associated cell death in Alzheimer's disease. *J Neurosci* 26: 6377-85
- Alzheimer A. 1907. Über eine eigenartige Erkrankung der Hirnrinde. *Allgemeine Zeitschrift für Psychiatrie und phychisch-Gerichtliche Medizin, (Berlin)* 64: 146-48
- Alzheimer A, Stelzmann RA, Schnitzlein HN, Murtagh FR. 1995. An English translation of Alzheimer's 1907 paper, "Über eine eigenartige Erkrankung der Hirnrinde". *Clin Anat* 6: 429-31

- Alzheimer's Association. 2014. 2014 Alzheimer's Disease Facts and Figures.
- Alzheimer's Disease International. 2012. Overcoming the stigma of dementia. *World Alzheimer's Report*
- Alzheimer's Association. 2006. Early-Onset Dementia: A National Challenge, a Future Crisis. *Washington, D.C.: Alzheimer's Association*
- Ancolio K, Dumanchin C, Barelli H, Warter JM, Brice A, et al. 1999. Unusual phenotypic alteration of beta amyloid precursor protein (betaAPP) maturation by a new Val-715 --> Met betaAPP-770 mutation responsible for probable early-onset Alzheimer's disease. *Proc Natl Acad Sci U S A* 96: 4119-24
- Andersen K, Launer LJ, Ott A, Hoes AW, Breteler MM, Hofman A. 1995. Do nonsteroidal anti-inflammatory drugs decrease the risk for Alzheimer's disease? The Rotterdam Study. *Neurology*. 45: 1441-45
- Andersson E, Jensen JB, Parmar M, Guillemot F, Bjorklund A. 2006. Development of the mesencephalic dopaminergic neuron system is compromised in the absence of neurogenin 2. *Development* 133: 507-16
- Ando K, Iijima KI, Elliott JI, Kirino Y, Suzuki T. 2001. Phosphorylation-dependent regulation of the interaction of amyloid precursor protein with Fe65 affects the production of beta-amyloid. *J Biol Chem* 276: 40353-61
- Anthony TE, Klein C, Fishell G, Heintz N. 2004. Radial glia serve as neuronal progenitors in all regions of the central nervous system. *neuron* 25: 881-90
- Araki R, Uda M, Hoki Y, Sunayama M, Nakamura M, et al. 2013. Negligible immunogenicity of terminally differentiated cells derived from induced pluripotent or embryonic stem cells. *Nature* 494: 100-4
- Arancio O, Chao MV. 2007. Neurotrophins, synaptic plasticity and dementia. *Curr Opin Neurobiol* 17: 325-30
- Arendash GW, Gordon MN, Diamond DM, Austin LA, Hatcher JM, et al. 2001. Behavioral assessment of Alzheimer's transgenic mice following long-term Abeta vaccination: task specificity and correlations between Abeta deposition and spatial memory. *DNA Cell Biol* 20: 737-44
- Arispe N, Rojas E, Pollard HB. 1993. Alzheimer disease amyloid beta protein forms calcium channels in bilayer membranes: blockade by tromethamine and aluminum. *Proc Natl Acad Sci U S A* 90: 567-71
- Aronica E, van Vliet EA, Hendriksen E, Troost D, Lopes da Silva FH, Gorter JA. 2001. Cystatin C, a cysteine protease inhibitor, is persistently up-regulated in neurons and glia in a rat model for mesial temporal lobe epilepsy. *Eur J Neurosci* 14: 1485-91
- Arriagada PV, Growdon JH, Hedley-Whyte ET, Hyman BT. 1992. Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology* 42: 631-9
- Arruda-Carvalho M, Sakaguchi M, Akers KG, Josselyn SA, Frankland PW. 2011. Posttraining ablation of adult-generated neurons degrades previously acquired memories. *J Neurosci* 31: 15113-27
- Arsenijevic Y, Weiss S, Schneider B, Aebischer P. 2001. Insulin-like growth factor-I is necessary for neural stem cell proliferation and demonstrates distinct actions of epidermal growth factor and fibroblast growth factor-2. *J Neurosci* 21: 7194-202

- Arvanitakis Z, Leurgans SE, Wang Z, Wilson RS, Bennett DA, Schneider JA. 2011. Cerebral amyloid angiopathy pathology and cognitive domains in older persons. *Annals of neurology* 69: 320-7
- Ashe KH, Zahs KR. 2010. Probing the biology of Alzheimer's disease in mice. *Neuron* 66: 631-45
- Australian Institute of Health and Welfare. 2012. Dementia in Australia.
- Ayuso-Sacido A, Moliterno JA, Kratovac S, Kapoor GS, O'Rourke DM, et al. 2010. Activated EGFR signaling increases proliferation, survival, and migration and blocks neuronal differentiation in post-natal neural stem cells. *J Neurooncol* 97: 323-37
- Baitsch D, Bock HH, Engel T, Telgmann R, Muller-Tidow C, et al. 2011. Apolipoprotein E induces antiinflammatory phenotype in macrophages. *Arterioscler Thromb Vasc Biol* 31: 1160-8
- Bajetto A, Bonavia R, Barbero S, Schettini G. 2002. Characterization of chemokines and their receptors in the central nervous system: physiopathological implications. *Journal of neurochemistry* 82: 1311-29
- Baker J, Liu JP, Robertson EJ, Efstratiadis A. 1993. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell* 75: 73-82
- Baldauf K, Reymann KG. 2005. Influence of EGF/bFGF treatment on proliferation, early neurogenesis and infarct volume after transient focal ischemia. *Brain Res* 1056: 158-67
- Bales KR, Verina T, Dodel RC, Du Y, Altstiel L, et al. 1997. Lack of apolipoprotein E dramatically reduces amyloid beta-peptide deposition. *Nat Genet* 17: 263-4
- Bali J, Gheinani AH, Zurbriggen S, Rajendran L. 2012. Role of genes linked to sporadic Alzheimer's disease risk in the production of beta-amyloid peptides. *Proc Natl Acad Sci U S A* 109: 15307-11
- Balu R, Pressler RT, Strowbridge BW. 2007. Multiple modes of synaptic excitation of olfactory bulb granule cells. *J Neurosci* 27: 5621-32
- Banks WA, Jaspan JB, Huang W, Kastin AJ. 1997a. Transport of insulin across the blood-brain barrier: saturability at euglycemic doses of insulin. *Peptides* 18: 1423-9
- Banks WA, Jaspan JB, Kastin AJ. 1997b. Selective, physiological transport of insulin across the blood-brain barrier: novel demonstration by species-specific radioimmunoassays. *Peptides* 18: 1257-62
- Baratchi S, Evans J, Tate WP, Abraham WC, Connor B. 2012. Secreted amyloid precursor proteins promote proliferation and glial differentiation of adult hippocampal neural progenitor cells. *Hippocampus* 22: 1517-27
- Barbero S, Bajetto A, Bonavia R, Porcile C, Piccioli P, et al. 2002. Expression of the chemokine receptor CXCR4 and its ligand stromal cell-derived factor 1 in human brain tumors and their involvement in glial proliferation in vitro. *Ann N Y Acad Sci* 973: 60-9
- Bard F, Cannon C, Barbour R, Burke RL, Games D, et al. 2000. Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. *Nature medicine* 6: 916-9
- Bartlett WP, Li XS, Williams M, Benkovic S. 1991. Localization of insulin-like growth factor-1 mRNA in murine central nervous system during postnatal development. *Dev Biol* 147: 239-50

- Bartus RT, Dean RL, 3rd, Beer B, Lippa AS. 1982. The cholinergic hypothesis of geriatric memory dysfunction. *Science* 217: 408-14
- Basak JM, Verghese PB, Yoon H, Kim J, Holtzman DM. 2012. Low-density lipoprotein receptor represents an apolipoprotein E-independent pathway of Abeta uptake and degradation by astrocytes. *J Biol Chem* 287: 13959-71
- Bayer SA. 1983. 3H-thymidine-radiographic studies of neurogenesis in the rat olfactory bulb. *Exp Brain Res* 50: 329-40
- Bayley PJ, Salmon DP, Bondi M, W., Bui BK, Olichney J, et al. 2000. Comparison of the serial position effect in very mild Alzheimer's disease, mild Alzheimer's disease, and amnesia associated with electroconvulsive therapy. *J Int Neuropsychol Soc* 6: 290-8
- Beckman M, Freeman C, Parish CR, Small DH. 2009. Activation of cathepsin D by glycosaminoglycans. *Febs J* 276: 7343-52
- Bedard A, Parent A. 2004. Evidence of newly generated neurons in the human olfactory bulb. *Brain Res Dev Brain Res* 151: 159-68
- Behr D, Hesse L, Masters CL, Multhaup G. 1996. Regulation of amyloid protein precursor (APP) binding to collagen and mapping of the binding sites on APP and collagen type I. *J Biol Chem* 271: 1613-20
- Bell RD, Sagare AP, Friedman AE, Bedi GS, Holtzman DM, et al. 2007. Transport pathways for clearance of human Alzheimer's amyloid beta-peptide and apolipoproteins E and J in the mouse central nervous system. *J Cereb Blood Flow Metab* 27: 909-18
- Belluzzi O, Benedusi M, Ackman J, LoTurco JJ. 2003. Electrophysiological differentiation of new neurons in the olfactory bulb. *J Neurosci* 23: 10411-8
- Belnoue L, Grosjean N, Abrous DN, Koehl M. 2011. A critical time window for the recruitment of bulbar newborn neurons by olfactory discrimination learning. *J Neurosci* 31: 1010-6
- Belvindrah R, Hankel S, Walker J, Patton BL, Muller U. 2007. Beta1 integrins control the formation of cell chains in the adult rostral migratory stream. *J Neurosci* 27: 2704-17
- Ben-Hur T, Ben-Menachem O, Furer V, Einstein O, Mizrachi-Kol R, Grigoriadis N. 2003. Effects of proinflammatory cytokines on the growth, fate, and motility of multipotential neural precursor cells. *Mol Cell Neurosci* 24: 623-31
- Bentahir M, Nyabi O, Verhamme J, Tolia A, Horre K, et al. 2006. Presenilin clinical mutations can affect gamma-secretase activity by different mechanisms. *Journal of neurochemistry* 96: 732-42
- Bentham P, Gray R, Sellwood E, Hills R, Crome P, Raftery J. 2008. Aspirin in Alzheimer's disease (AD2000): a randomised open-label trial. *Lancet Neurol* 7: 41-9
- Berdowska I, Siewinski M. 2000. [The role of cysteine cathepsins and their inhibitors in physiological and neoplastic processes]. *Postepy Biochem* 46: 73-84
- Bergmans BA, De Strooper B. 2010. gamma-secretases: from cell biology to therapeutic strategies. *Lancet Neurol* 9: 215-26
- Bergmans BA, Shariati SA, Habets RL, Verstreken P, Schoonjans L, et al. 2010. Neurons generated from APP/APLP1/APLP2 triple knockout embryonic

- stem cells behave normally in vitro and in vivo: lack of evidence for a cell autonomous role of the amyloid precursor protein in neuronal differentiation. *stem cells* 28: 399-406
- Bernstein HG, Kirschke H, Wiederanders B, Pollak KH, Zipress A, Rinne A. 1996. The possible place of cathepsins and cystatins in the puzzle of Alzheimer disease: a review. *Mol Chem Neuropathol* 27: 225-47
- Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE. 2007. Systematic meta-analyses of Alzheimer disease genetic association studies: the AlzGene database. *Nat Genet* 39: 17-23
- Bertrand N, Castro DS, Guillemot F. 2002. Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* 3: 517-30
- Bhattacharyya BJ, Banisadr G, Jung H, Ren D, Cronshaw DG, et al. 2008. The chemokine stromal cell-derived factor-1 regulates GABAergic inputs to neural progenitors in the postnatal dentate gyrus. *J Neurosci* 28: 6720-30
- Bhattacharyya R, Barren C, Kovacs DM. 2013. Palmitoylation of amyloid precursor protein regulates amyloidogenic processing in lipid rafts. *J Neurosci* 33: 11169-83
- Bierer LM, Hof PR, Purohit DP, Carlin L, Schmeidler J, et al. 1995. Neocortical neurofibrillary tangles correlate with dementia severity in Alzheimer's disease. *Arch Neurol* 52: 81-8
- Bignami A, Dahl D. 1974. Astrocyte-specific protein and neuroglial differentiation. An immunofluorescence study with antibodies to the glial fibrillary acidic protein. *J Comp Neurol* 153: 27-38
- Billings LM, Oddo S, Green KN, McGaugh JL, LaFerla FM. 2005. Intraneuronal Aβ causes the onset of early Alzheimer's disease-related cognitive deficits in transgenic mice. *Neuron* 45: 675-88
- Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, et al. 1997. A metalloproteinase disintegrin that releases tumour-necrosis factor-α from cells. *nature* 385: 729-33
- Blair P, Flaumenhaft R. 2009. Platelet α-granules: basic biology and clinical correlates. *Blood Rev* 23: 177-89
- Blennow K, Hampel H, Zetterberg H. 2014. Biomarkers in amyloid-β immunotherapy trials in Alzheimer's disease. *Neuropsychopharmacology* 39: 189-201
- Blennow K, Zetterberg H, Rinne JO, Salloway S, Wei J, et al. 2012. Effect of immunotherapy with bapineuzumab on cerebrospinal fluid biomarker levels in patients with mild to moderate Alzheimer disease. *Arch Neurol* 69: 1002-10
- Blumbergs PC, Scott G, Manavis J, Wainwright H, Simpson DA, McLean AJ. 1995. Topography of axonal injury as defined by amyloid precursor protein and the sector scoring method in mild and severe closed head injury. *J Neurotrauma* 12: 565-72
- Blurton-Jones M, Kitazawa M, Martinez-Coria H, Castello NA, Muller FJ, et al. 2009. Neural stem cells improve cognition via BDNF in a transgenic model of Alzheimer disease. *Proc Natl Acad Sci U S A* 106: 13594-9
- Bobek LA, Levine MJ. 1992. Cystatins--inhibitors of cysteine proteinases. *Crit Rev Oral Biol Med* 3: 307-32
- Bode W, Huber R. 2000. Structural basis of the endoproteinase-protein inhibitor interaction. *Biochim Biophys Acta* 1477: 241-52

- Boekhoorn K, Joels M, Lucassen PJ. 2006. Increased proliferation reflects glial and vascular-associated changes, but not neurogenesis in the presenile Alzheimer hippocampus. *Neurobiol Dis* 24: 1-14
- Bohrmann B, Baumann K, Benz J, Gerber F, Huber W, et al. 2012. Gantenerumab: a novel human anti-Aβ antibody demonstrates sustained cerebral amyloid-beta binding and elicits cell-mediated removal of human amyloid-beta. *Journal of Alzheimer's disease : JAD* 28: 49-69
- Boku S, Nakagawa S, Takamura N, Kato A, Takebayashi M, et al. 2013. GDNF facilitates differentiation of the adult dentate gyrus-derived neural precursor cells into astrocytes via STAT3. *Biochem Biophys Res Commun* 434: 779-84
- Boland B, Campbell V. 2004. Amyloid-beta -mediated activation of the apoptotic cascade in cultured cortical neurones: a role for cathepsin-L. *Neurobiol Aging*. 25: 83-91
- Boldrini M, Underwood MD, Hen R, Rosoklija GB, Dwork AJ, et al. 2009. Antidepressants increase neural progenitor cells in the human hippocampus. *Neuropsychopharmacology* 34: 2376-89
- Bolos M, Hu Y, Young KM, Foa L, Small DH. 2014. Neurogenin 2 mediates amyloid-beta precursor protein-stimulated neurogenesis. *J Biol Chem* 289: 31253-61
- Bonaguidi MA, Peng CY, McGuire T, Falciglia G, Gobeske KT, et al. 2008. Noggin expands neural stem cells in the adult hippocampus. *J Neurosci* 28: 9194-204
- Bonifacino JS, Traub LM. 2003. Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annu Rev Biochem* 72: 395-447
- Bordiuk OL, Smith K, Morin PJ, Semenov MV. 2014. Cell proliferation and neurogenesis in adult mouse brain. *PLoS One* 9: e111453
- Borg JP, Ooi J, Levy E, Margolis B. 1996. The phosphotyrosine interaction domains of X11 and FE65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. *Mol Cell Biol* 16: 6229-41
- Braak H, Alafuzoff I, Arzberger T, Kretschmar H, Del Tredici K. 2006. Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* 4: 389-404
- Braak H, Braak E. 1991a. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol* 82: 239-59
- Braak H, Braak E. 1991b. Neuropathological staging of Alzheimer-related changes *Acta Neuropathol* 82: 239-59
- Braak H, Braak E. 1995. Staging of Alzheimer's disease-related neurofibrillary changes. *Neurobiol Aging* 16: 271-8; discussion 78-84
- Braak H, Thal DR, Ghebremedhin E, Del Tredici K. 2011. Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. *J Neuropathol Exp Neurol* 70: 960-9
- Bracko O, Singer T, Aigner S, Knobloch M, Winner B, et al. 2012. Gene expression profiling of neural stem cells and their neuronal progeny reveals IGF2 as a regulator of adult hippocampal neurogenesis. *J Neurosci* 32: 3376-87
- Braun SM, Jessberger S. 2014. Adult neurogenesis: mechanisms and functional significance. *Development* 141: 1983-6
- Bressler SL, Gray MD, Sopher BL, Hu Q, Hearn MG, et al. 1996. cDNA cloning and chromosome mapping of the human Fe65 gene: interaction of the

- conserved cytoplasmic domains of the human beta-amyloid precursor protein and its homologues with the mouse Fe65 protein. *Hum Mol Genet* 5: 1589-98
- Brezun JM, Daszuta A. 2000. Serotonin may stimulate granule cell proliferation in the adult hippocampus, as observed in rats grafted with foetal raphe neurons. *Eur J Neurosci* 12: 391-6
- Brown JP, Couillard-Despres S, Cooper-Kuhn CM, Winkler J, Aigner L, Kuhn HG. 2003. Transient expression of doublecortin during adult neurogenesis. *J Comp Neurol* 467: 1-10
- Brown MS, Ye J, Rawson RB, Goldstein JL. 2000. Regulated intramembrane proteolysis: a control mechanism conserved from bacteria to humans. *Cell* 100: 391-8
- Brunnström HR, Englund EM. 2009. Cause of death in patients with dementia disorders. *Eur J Neurol* 4: 488-92
- Burns JL, Hassan AB. 2001. Cell survival and proliferation are modified by insulin-like growth factor 2 between days 9 and 10 of mouse gestation. *Development* 128: 3819-30
- Burstein AH, Zhao Q, Ross J, Styren S, Landen JW, et al. 2013. Safety and pharmacology of ponezumab (PF-04360365) after a single 10-minute intravenous infusion in subjects with mild to moderate Alzheimer disease. *Clin Neuropharmacol* 36: 8-13
- Bush AI, Martins RN, Rumble B, Moir R, Fuller S, et al. 1990. The amyloid precursor protein of Alzheimer's disease is released by human platelets. *J Biol Chem* 265: 15977-83
- Bush AI, Multhaup G, Moir RD, Williamson TG, Small DH, et al. 1993. A novel zinc(II) binding site modulates the function of the beta A4 amyloid protein precursor of Alzheimer's disease. *J Biol Chem* 268: 16109-12
- Butterfield DA, Pocernich CB. 2003. The glutamatergic system and Alzheimer's disease: therapeutic implications. *CNS Drugs* 17: 641-52
- Buxbaum JD, Liu KN, Luo Y, Slack JL, Stocking KL, et al. 1998. Evidence that tumor necrosis factor alpha converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J Biol Chem* 273: 27765-7
- Buxbaum JD, Thinakaran G, Koliatsos V, O'Callahan J, Slunt HH, et al. 1988. Endoplasmic reticulum and trans-Golgi network generate distinct populations of Alzheimer beta-amyloid peptides. *J Neurosci* 18: 9629-37
- Caccamo A, Oddo S, Billings LM, Green KN, Martinez-Coria H, et al. 2006. M1 receptors play a central role in modulating AD-like pathology in transgenic mice. *Neuron* 49: 671-82
- Cady J, Koval ED, Benitez BA, Zaidman C, Jockel-Balsarotti J, et al. 2014. TREM2 variant p.R47H as a risk factor for sporadic amyotrophic lateral sclerosis. *JAMA Neurol* 71: 449-53
- Cai H, Wang Y, McCarthy D, Wen H, Borchelt DR, et al. 2001. BACE1 is the major beta-secretase for generation of Abeta peptides by neurons. *Nat Neurosci* 4: 233-4
- Cai J, Qi X, Kociok N, Skosyrski S, Emilio A, et al. 2012. beta-Secretase (BACE1) inhibition causes retinal pathology by vascular dysregulation and accumulation of age pigment. *EMBO Mol Med* 4: 980-91

- Cai J, Yang M, Poremsky E, Kidd S, Schneider JS, Iacovitti L. 2010. Dopaminergic neurons derived from human induced pluripotent stem cells survive and integrate into 6-OHDA-lesioned rats. *Stem cells and development* 19: 1017-23
- Caille I, Allinquant B, Dupont E, Bouillot C, Langer A, et al. 2004. Soluble form of amyloid precursor protein regulates proliferation of progenitors in the adult subventricular zone. *Development* 131: 2173-81
- Calero M, Rostagno A, Ghiso J. 2012. Search for amyloid-binding proteins by affinity chromatography. *Methods Mol Biol* 849: 213-23
- Calvo CF, Fontaine RH, Soueid J, Tammela T, Makinen T, et al. 2011. Vascular endothelial growth factor receptor 3 directly regulates murine neurogenesis. *Genes Dev* 25: 831-44
- Cameron HA, Tanapat P, Gould E. 1998. Adrenal steroids and N-methyl-D-aspartate receptor activation regulate neurogenesis in the dentate gyrus of adult rats through a common pathway. *Neuroscience* 82: 349-54
- Cameron HA, Woolley CS, McEwen BS, Gould E. 1993. Differentiation of newly born neurons and glia in the dentate gyrus of the adult rat. *Neuroscience* 56: 337-44
- Cantarella C, Cayre M, Magalon K, Durbec P. 2008. Intranasal HB-EGF administration favors adult SVZ cell mobilization to demyelinated lesions in mouse corpus callosum. *Dev Neurobiol* 68: 223-36
- Cao L, Wei D, Reid B, Zhao S, Pu J, et al. 2013. Endogenous electric currents might guide rostral migration of neuroblasts. *EMBO Rep* 14: 184-90
- Cao X, Sudhof TC. 2001. A transcriptionally [correction of transcriptively] active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* 293: 115-20
- Cao X, Sudhof TC. 2004. Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J Biol Chem* 279: 24601-11
- Cartier L, Hartley O, Dubois-Dauphin M, Krause KH. 2005. Chemokine receptors in the central nervous system: role in brain inflammation and neurodegenerative diseases. *Brain Res Brain Res Rev* 48: 16-42
- Cataldo AM, Nixon RA. 1990. Enzymatically active lysosomal proteases are associated with amyloid deposits in Alzheimer brain. *Proc Natl Acad Sci U S A* 87: 3861-5
- Chabrier MA, Blurton-Jones M, Agazaryan AA, Nerhus JL, Martinez-Coria H, LaFerla FM. 2012. Soluble abeta promotes wild-type tau pathology in vivo. *J Neurosci* 32: 17345-50
- Chapman HA, Riese RJ, Shi GP. 1997. Emerging roles for cysteine proteases in human biology. *Annu Rev Physiol* 59: 63-88
- Chapouton P, Schuurmans C, Guillemot F, Gotz M. 2001. The transcription factor neurogenin 2 restricts cell migration from the cortex to the striatum. *Development* 128: 5149-59
- Chasseigneaux S, Dinc L, Rose C, Chabret C, Couplier F, et al. 2011. Secreted amyloid precursor protein beta and secreted amyloid precursor protein alpha induce axon outgrowth in vitro through Egr1 signaling pathway. *PLoS One* 6: e16301
- Checler F, Sunyach C, Pardossi-Piquard R, Sevalle J, Vincent B, et al. 2007. The gamma/epsilon-secretase-derived APP intracellular domain fragments regulate p53. *Curr Alzheimer Res* 4: 423-6

- Chen Q, Nakajima A, Choi SH, Xiong X, Sisodia SS, Tang YP. 2008. Adult neurogenesis is functionally associated with AD-like neurodegeneration. *Neurobiol Dis* 29: 316-26
- Chen WW, Blurton-Jones M. 2012. Concise review: Can stem cells be used to treat or model Alzheimer's disease. *stem cells* 30: 2612-8
- Chen Y, Dong C. 2009. Abeta40 promotes neuronal cell fate in neural progenitor cells. *Cell Death Differ* 16: 386-94
- Chen Y, Tang BL. 2006. The amyloid precursor protein and postnatal neurogenesis/neuroregeneration. *Biochem Biophys Res Commun* 341: 1-5
- Cheng Y, Black IB, DiCicco-Bloom E. 2002. Hippocampal granule neuron production and population size are regulated by levels of bFGF. *Eur J Neurosci* 15: 3-12
- Chertoff M, Shrivastava K, Gonzalez B, Acarin L, Gimenez-Llort L. 2013. Differential modulation of TREM2 protein during postnatal brain development in mice. *PLoS One* 8: e72083
- Chishti MA, Yang DS, Janus C, Phinney AL, Horne P, et al. 2001. Early-onset amyloid deposition and cognitive deficits in transgenic mice expressing a double mutant form of amyloid precursor protein 695. *J Biol Chem* 276: 21562-70
- Cho GW, Noh MY, Kim HY, Koh SH, Kim KS, Kim SH. 2010a. Bone marrow-derived stromal cells from amyotrophic lateral sclerosis patients have diminished stem cell capacity. *Stem cells and development* 19: 1035-42
- Cho HJ, Lee CS, Kwon YW, Paek JS, Lee SH, et al. 2010b. Induction of pluripotent stem cells from adult somatic cells by protein-based reprogramming without genetic manipulation. *Blood* 116: 386-95
- Choi MR, Kim HY, Park JY, Lee TY, Baik CS, et al. 2010. Selection of optimal passage of bone marrow-derived mesenchymal stem cells for stem cell therapy in patients with amyotrophic lateral sclerosis. *Neurosci Lett* 472: 94-8
- Chouraki V, Seshadri S. 2014. Genetics of Alzheimer's disease. *Adv Genet* 87: 245-94
- Christie KJ, Turnley AM. 2012. Regulation of endogenous neural stem/progenitor cells for neural repair-factors that promote neurogenesis and gliogenesis in the normal and damaged brain. *Front Cell Neurosci* 6: 70
- Chun HJ, Zheng L, Ahmad M, Wang J, Speirs CK, et al. 2002. Pleiotropic defects in lymphocyte activation caused by caspase-8 mutations lead to human immunodeficiency. *Nature* 419: 395-9
- Citron M, Eckman CB, Diehl TS, Corcoran C, Ostaszewski BL, et al. 1998. Additive effects of PS1 and APP mutations on secretion of the 42-residue amyloid beta-protein. *Neurobiol Dis* 5: 107-16
- Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, et al. 1992. Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature* 360: 672-4
- Citron M, Westaway D, Xia W, Carlson G, Diehl T, et al. 1997. Mutant presenilins of Alzheimer's disease increase production of 42-residue amyloid beta-protein in both transfected cells and transgenic mice. *Nature medicine* 3: 67-72
- Clarris HJ, Cappai R, Heffernan D, Beyreuther K, Masters CL, Small DH. 1997. Identification of heparin-binding domains in the amyloid precursor

- protein of Alzheimer's disease by deletion mutagenesis and peptide mapping. *Journal of neurochemistry* 68: 1164-72
- Clarris HJ, Key B, Beyreuther K, Masters CL, Small DH. 1995. Expression of the amyloid protein precursor of Alzheimer's disease in the developing rat olfactory system. *Brain Res Dev Brain Res* 88: 87-95
- Cochran E, Bacci B, Chen Y, Patton A, Gambetti P, Autilio-Gambetti L. 1991. Amyloid precursor protein and ubiquitin immunoreactivity in dystrophic axons is not unique to Alzheimer's disease. *Am J Pathol* 139: 485-9
- Codega P, Silva-Vargas V, Paul A, Maldonado-Soto AR, Deleo AM, et al. 2014. Prospective identification and purification of quiescent adult neural stem cells from their in vivo niche. *Neuron* 82: 545-59
- Colditz MJ, Catts VS, Al-menhali N, Osborne GW, Bartlett PF, Coulson EJ. 2010. p75 neurotrophin receptor regulates basal and fluoxetine-stimulated hippocampal neurogenesis. *Exp Brain Res* 200: 161-7
- Colonna M. 2003. TREMs in the immune system and beyond. *Nature Reviews Immunology* 3: 445-53
- Cooke MJ, Wang Y, Morshead CM, Shoichet MS. 2011. Controlled epi-cortical delivery of epidermal growth factor for the stimulation of endogenous neural stem cell proliferation in stroke-injured brain. *Biomaterials* 32: 5688-97
- Cooper-Kuhn CM, Winkler J, Kuhn HG. 2004. Decreased neurogenesis after cholinergic forebrain lesion in the adult rat. *Journal of neuroscience research* 77: 155-65
- Corder EH, Saunders AM, Risch NJ, Strittmatter WJ, Schmechel DE, et al. 1994. Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer disease. *Nat Genet* 7: 180-4
- Côté S, Carmichael PH, Verreault R, Lindsay J, Lefebvre J, Laurin D. 2012. Nonsteroidal anti-inflammatory drug use and the risk of cognitive impairment and Alzheimer's disease. *Alzheimers Dement*. 8: 219 -26
- Cousins SL, Hoey SE, Anne Stephenson F, Perkinton MS. 2009. Amyloid precursor protein 695 associates with assembled NR2A- and NR2B-containing NMDA receptors to result in the enhancement of their cell surface delivery. *Journal of neurochemistry* 111: 1501-13
- Coyle JT, Price DL, DeLong MR. 1983. Alzheimer's disease: a disorder of cortical cholinergic innervation. *Science* 219: 1184-90
- Craig CG, Tropepe V, Morshead CM, Reynolds BA, Weiss S, van der Kooy D. 1996. In vivo growth factor expansion of endogenous subependymal neural precursor cell populations in the adult mouse brain. *J Neurosci* 16: 2649-58
- Cras P, Kawai M, Lowery D, Gonzalez-DeWhitt P, Greenberg B, Perry G. 1991. Senile plaque neurites in Alzheimer disease accumulate amyloid precursor protein. *Proc Natl Acad Sci U S A* 88: 7552-6
- Crawford FC, Freeman MJ, Schinka JA, Abdullah LI, Gold M, et al. 2000. A polymorphism in the cystatin C gene is a novel risk factor for late-onset Alzheimer's disease. *Neurology* 55: 763-8
- Crews L, Mizuno H, Desplats P, Rockenstein E, Adame A, et al. 2008. Alpha-synuclein alters Notch-1 expression and neurogenesis in mouse embryonic stem cells and in the hippocampus of transgenic mice. *J Neurosci* 28: 4250-60

- Cummings DM, Snyder JS, Brewer M, Cameron HA, Belluscio L. 2014. Adult neurogenesis is necessary to refine and maintain circuit specificity. *J Neurosci* 34: 13801-10
- Cummings JL, Vinters HV, Cole GM, Khachaturian ZS. 1998. Alzheimer's disease: etiologies, pathophysiology, cognitive reserve, and treatment opportunities. *Neurology* 51: S2-17; discussion S65-7
- Cupers P, Orlans I, Craessaerts K, Annaert W, De Strooper B. 2001. The amyloid precursor protein (APP)-cytoplasmic fragment generated by gamma-secretase is rapidly degraded but distributes partially in a nuclear fraction of neurones in culture. *Journal of neurochemistry* 78: 1168-78
- Curtis MA, Penney EB, Pearson AG, van Roon-Mom WM, Butterworth NJ, et al. 2003. Increased cell proliferation and neurogenesis in the adult human Huntington's disease brain. *Proc Natl Acad Sci U S A* 100: 9023-7
- Cuyvers E, Bettens K, Philtjens S, Van Langenhove T, Gijselinck I, et al. 2014. Investigating the role of rare heterozygous TREM2 variants in Alzheimer's disease and frontotemporal dementia. *Neurobiol Aging* 35: 726 e11-9
- D'Andrea MR, Nagele RG. 2002. MAP-2 immunolabeling can distinguish diffuse from dense-core amyloid plaques in brains with Alzheimer's disease. *Biotech Histochem* 77: 95-103
- D'Ercole AJ, Ye P, Calikoglu AS, Gutierrez-Ospina G. 1996. The role of the insulin-like growth factors in the central nervous system. *Mol Neurobiol* 13: 227-55
- Dahl A, Eriksson PS, Davidsson P, Persson AI, Ekman R, Westman-Brinkmalm A. 2004. Demonstration of multiple novel glycoforms of the stem cell survival factor Cc. *Journal of neuroscience research* 77: 9-14
- Dahl A, Eriksson PS, Persson AI, Karlsson G, Davidsson P, et al. 2003. Proteome analysis of conditioned medium from cultured adult hippocampal progenitors. *Rapid Commun Mass Spectrom* 17: 2195-202
- Dahlgren KN, Manelli AM, Stine WB, Jr., Baker LK, Krafft GA, LaDu MJ. 2002. Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 277: 32046-53
- Dahms SO, Hoefgen S, Roeser D, Schlott B, Guhrs KH, Than ME. 2010. Structure and biochemical analysis of the heparin-induced E1 dimer of the amyloid precursor protein. *Proc Natl Acad Sci U S A* 107: 5381-6
- Danysz W, Parsons CG. 2012. Alzheimer's disease, β -amyloid, glutamate, NMDA receptors and memantine--searching for the connections. *Br J Pharmacol* 167: 324-52
- Dawkins E, Small DH. 2014. Insights into the physiological function of the beta-amyloid precursor protein: beyond Alzheimer's disease. *Journal of neurochemistry* 129: 756-69
- Dawson GR, Seabrook GR, Zheng H, Smith DW, Graham S, et al. 1999. Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience* 90: 1-13
- Dayer AG, Jenny B, Sauvain MO, Potter G, Salmon P, et al. 2007. Expression of FGF-2 in neural progenitor cells enhances their potential for cellular brain repair in the rodent cortex. *Brain* 130: 2962-76

- de Azevedo-Pereira RL, Lima AP, Rodrigues Dde C, Rondinelli E, Medei EH, et al. 2011. Cysteine proteases in differentiation of embryonic stem cells into neural cells. *Stem Cells Dev* 20: 1859-72
- De Felice FG, Velasco PT, Lambert MP, Viola K, Fernandez SJ, et al. 2007. Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Biol Chem* 282: 11590-601
- De Marchis S, Temoney S, Erdelyi F, Bovetti S, Bovolin P, et al. 2004. GABAergic phenotypic differentiation of a subpopulation of subventricular derived migrating progenitors. *Eur J Neurosci* 20: 1307-17
- De Strooper B. 2003. Aph-1, Pen-2, and Nicastrin with Presenilin generate an active gamma-Secretase complex. *Neuron* 38: 9-12
- De Strooper B. 2007. Loss-of-function presenilin mutations in Alzheimer disease. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep* 8: 141-6
- De Strooper B, Annaert W, Cupers P, Saftig P, Craessaerts K, et al. 1999. A presenilin-1-dependent gamma-secretase-like protease mediates release of Notch intracellular domain. *Nature* 398: 518-22
- De Strooper B, Saftig P, Craessaerts K, Vanderstichele H, Guhde G, et al. 1998. Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* 391: 387-90
- Deane R, Sagare A, Hamm K, Parisi M, Lane S, et al. 2008. apoE isoform-specific disruption of amyloid beta peptide clearance from mouse brain. *The Journal of clinical investigation* 118: 4002-13
- Dehghan S, Javan M, Pourabdolhossein F, Mirnajafi-Zadeh J, Baharvand H. 2012. Basic fibroblast growth factor potentiates myelin repair following induction of experimental demyelination in adult mouse optic chiasm and nerves. *J Mol Neurosci* 48: 77-85
- Demars MP, Bartholomew A, Strakova Z, Lazarov O. 2011. Soluble amyloid precursor protein: a novel proliferation factor of adult progenitor cells of ectodermal and mesodermal origin. *Stem Cell Res Ther* 2: 36
- DeMattos RB, Brendza RP, Heuser JE, Kierson M, Cirrito JR, et al. 2001. Purification and characterization of astrocyte-secreted apolipoprotein E and J-containing lipoproteins from wild-type and human apoE transgenic mice. *Neurochem Int* 39: 415-25
- Demuro A, Mina E, Kaye R, Milton SC, Parker I, Glabe CG. 2005. Calcium dysregulation and membrane disruption as a ubiquitous neurotoxic mechanism of soluble amyloid oligomers. *J Biol Chem* 280: 17294-300
- Deng A, Irizarry MC, Nitsch RM, Growdon JH, Rebeck GW. 2001. Elevation of cystatin C in susceptible neurons in Alzheimer's disease. *Am J Pathol* 159: 1061-8
- Deng W, Aimone JB, Gage FH. 2010. New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 11: 339-50
- Deng W, Saxe MD, Gallina IS, Gage FH. 2009. Adult-born hippocampal dentate granule cells undergoing maturation modulate learning and memory in the brain. *J Neurosci* 29: 13532-42

- Desai M, Li T, Ross MG. 2011. Fetal hypothalamic neuroprogenitor cell culture: preferential differentiation paths induced by leptin and insulin. *Endocrinology* 152: 3192-201
- Devore S, Linster C. 2012. Noradrenergic and cholinergic modulation of olfactory bulb sensory processing. *Front Behav Neurosci* 6: 52
- Dinet V, An N, Ciccotosto GD, Bruban J, Maoui A, et al. 2011. APP involvement in retinogenesis of mice. *Acta Neuropathol* 121: 351-63
- Dodd PR. 2002. Excited to death: different ways to lose your neurones. *Biogerontology* 3: 51-6
- Dodel RC, Du Y, Depboylu C, Kurz A, Eastwood B, et al. 2002. A polymorphism in the cystatin C promoter region is not associated with an increased risk of AD. *Neurology* 58: 664
- Doetsch F, Alvarez-Buylla A. 1996. Network of tangential pathways for neuronal migration in adult mammalian brain. *Proc Natl Acad Sci U S A* 93: 14895-900
- Doetsch F, Caille I, Lim DA, Garcia-Verdugo JM, Alvarez-Buylla A. 1999. Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97: 703-16
- Doetsch F, Garcia-Verdugo JM, Alvarez-Buylla A. 1997. Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain. *J Neurosci* 17: 5046-61
- Doetsch F, Petreanu L, Caille I, Garcia-Verdugo JM, Alvarez-Buylla A. 2002. EGF converts transit-amplifying neurogenic precursors in the adult brain into multipotent stem cells. *Neuron* 36: 1021-34
- Dominguez D, Tournoy J, Hartmann D, Huth T, Cryns K, et al. 2005. Phenotypic and biochemical analyses of BACE1- and BACE2-deficient mice. *J Biol Chem* 280: 30797-806
- Donnelly RJ, Rasool CG, Bartus R, Vitek S, Blume AJ, Vitek M. 1988. Multiple forms of beta-amyloid peptide precursor RNAs in a single cell type. *Neurobiol Aging* 9: 333-8
- Donovan MH, Yazdani U, Norris RD, Games D, German DC, Eisch AJ. 2006. Decreased adult hippocampal neurogenesis in the PDAPP mouse model of Alzheimer's disease. *J Comp Neurol* 495: 70-83
- Donovan PJ. 2001. High Oct-ane fuel powers the stem cell. *Nat Genet* 29: 246-7
- Donoviel DB, Hadjantonakis AK, Ikeda M, Zheng H, Hyslop PS, Bernstein A. 1999. Mice lacking both presenilin genes exhibit early embryonic patterning defects. *Genes Dev* 13: 2801-10
- Doody RS, Thomas RG, Farlow M, Iwatsubo T, Vellas B, et al. 2014. Phase 3 trials of solanezumab for mild-to-moderate Alzheimer's disease. *N Engl J Med* 370: 311-21
- Doraiswamy PM. 2003. The role of the N-methyl-D-aspartate receptor in Alzheimer's disease: therapeutic potential. *Curr Neurol Neurosci Rep* 3: 373-8
- Dreses-Werringloer U, Lambert JC, Vingtdeux V, Zhao H, Vais H, et al. 2008. A polymorphism in CALHM1 influences Ca²⁺ homeostasis, Aβ levels, and Alzheimer's disease risk. *Cell* 133: 1149-61
- Duff K, Eckman C, Zehr C, Yu X, Prada CM, et al. 1996. Increased amyloid-β₄₂(43) in brains of mice expressing mutant presenilin 1. *Nature* 383: 710-3

- Duinsbergen D, Salvatori D, Eriksson M, Mikkers H. 2009. Tumors originating from induced pluripotent stem cells and methods for their prevention. *Ann N Y Acad Sci* 1176: 197-204
- Duman RS. 2004. Depression: a case of neuronal life and death? *Biol Psychiatry* 56: 140-5
- Dupret D, Fabre A, Dobrossy MD, Panatier A, Rodriguez JJ, et al. 2007. Spatial learning depends on both the addition and removal of new hippocampal neurons. *PLoS Biol* 5: e214
- Duyckaerts C, Potier MC, Delatour B. 2008. Alzheimer disease models and human neuropathology: similarities and differences. *Acta Neuropathol* 115: 5-38
- Dyrks T, Weidemann A, Multhaup G, Salbaum JM, Lemaire HG, et al. 1988. Identification, transmembrane orientation and biogenesis of the amyloid A4 precursor of Alzheimer's disease. *Embo J* 7: 949-57
- Easter SS, Jr., Ross LS, Frankfurter A. 1993. Initial tract formation in the mouse brain. *J Neurosci* 13: 285-99
- Ebinu JO, Yankner BA. 2002. A RIP tide in neuronal signal transduction. *Neuron* 34: 499-502
- Eckenhoff MF, Rakic P. 1984. Radial organization of the hippocampal dentate gyrus: a Golgi, ultrastructural, and immunocytochemical analysis in the developing rhesus monkey. *J Comp Neurol* 223: 1-21
- Eckman C, Mehta N, Crook R, Perez-Tur J, Prihar G, et al. 1997. A New Pathogenic Mutation in the APP Gene (I716V) Increases the Relative Proportion of A β 42(43). *Hum. Mol. Genet* 6: 2087-89
- Effenberger T, von der Heyde J, Bartsch K, Garbers C, Schulze-Osthoff K, et al. 2014. Senescence-associated release of transmembrane proteins involves proteolytic processing by ADAM17 and microvesicle shedding. *Faseb J* 28: 4847-56
- Egberts F, Heinrich M, Jensen JM, Winoto-Morbach S, Pfeiffer S, et al. 2004. Cathepsin D is involved in the regulation of transglutaminase 1 and epidermal differentiation. *J Cell Sci* 117: 2295-307
- Eggert S, Paliga K, Soba P, Evin G, Masters CL, et al. 2004. The proteolytic processing of the amyloid precursor protein gene family members APLP-1 and APLP-2 involves alpha-, beta-, gamma-, and epsilon-like cleavages: modulation of APLP-1 processing by n-glycosylation. *J Biol Chem* 279: 18146-56
- Ehm O, Goritz C, Covic M, Schaffner I, Schwarz TJ, et al. 2010. RBPJkappa-dependent signaling is essential for long-term maintenance of neural stem cells in the adult hippocampus. *J Neurosci* 30: 13794-807
- Ekdahl CT, Mohapel P, Elmer E, Lindvall O. 2001. Caspase inhibitors increase short-term survival of progenitor-cell progeny in the adult rat dentate gyrus following status epilepticus. *Eur J Neurosci* 14: 937-45
- Ellis P, Fagan BM, Magness ST, Hutton S, Taranova O, et al. 2004. SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Dev Neurosci* 26: 148-65
- Encinas JM, Michurina TV, Peunova N, Park JH, Tordo J, et al. 2011. Division-coupled astrocytic differentiation and age-related depletion of neural stem cells in the adult hippocampus. *Cell Stem Cell* 8: 566-79

- Endres K, Postina R, Schroeder A, Mueller U, Fahrenholz F. 2005. Shedding of the amyloid precursor protein-like protein APLP2 by disintegrin-metalloproteinases. *Febs J* 272: 5808-20
- Erickson RI, Paucar AA, Jackson RL, Visnyei K, Kornblum H. 2008. Roles of insulin and transferrin in neural progenitor survival and proliferation. *Journal of neuroscience research* 86: 1884-94
- Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, et al. 1998. Neurogenesis in the adult human hippocampus. *Nature medicine* 4: 1313-7
- Ernst M, Jenkins BJ. 2004. Acquiring signalling specificity from the cytokine receptor gp130. *Trends Genet* 20: 23-32
- Esch FS, Keim PS, Beattie EC, Blacher RW, Culwell AR, et al. 1990. Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science* 248: 1122-4
- Esler WP, Stimson ER, Jennings JM, Vinters HV, Ghilardi JR, et al. 2000. Alzheimer's disease amyloid propagation by a template-dependent dock-lock mechanism. *Biochemistry* 39: 6288-95
- Esposito MS, Piatti VC, Laplagne DA, Morgenstern NA, Ferrari CC, et al. 2005. Neuronal differentiation in the adult hippocampus recapitulates embryonic development. *J Neurosci* 25: 10074-86
- Evans KC, Berger EP, Cho CG, Weisgraber KH, Lansbury PT, Jr. 1995. Apolipoprotein E is a kinetic but not a thermodynamic inhibitor of amyloid formation: implications for the pathogenesis and treatment of Alzheimer disease. *Proc Natl Acad Sci U S A* 92: 763-7
- Eyre MD, Antal M, Nusser Z. 2008. Distinct deep short-axon cell subtypes of the main olfactory bulb provide novel intrabulbar and extrabulbar GABAergic connections. *J Neurosci* 28: 8217-29
- Fagan AM, Watson M, Parsadanian M, Bales KR, Paul SM, Holtzman DM. 2002. Human and murine ApoE markedly alters A beta metabolism before and after plaque formation in a mouse model of Alzheimer's disease. *Neurobiol Dis* 9: 305-18
- Falk A, Holmstrom N, Carlen M, Cassidy R, Lundberg C, Frisen J. 2002. Gene delivery to adult neural stem cells. *Exp Cell Res* 279: 34-9
- Fallon J, Reid S, Kinyamu R, Opole I, Opole R, et al. 2000. In vivo induction of massive proliferation, directed migration, and differentiation of neural cells in the adult mammalian brain. *Proc Natl Acad Sci U S A* 97: 14686-91
- Faulkner RL, Jang MH, Liu XB, Duan X, Sailor KA, et al. 2008. Development of hippocampal mossy fiber synaptic outputs by new neurons in the adult brain. *Proc Natl Acad Sci U S A* 105: 14157-62
- Favaro R, Valotta M, Ferri AL, Latorre E, Mariani J, et al. 2009. Hippocampal development and neural stem cell maintenance require Sox2-dependent regulation of Shh. *Nature neuroscience* 12: 1248-56
- Feng MH, Chan SL, Xiang Y, Huber CP, Lim C. 1996. The binding mode of an E-64 analog to the active site of cathepsin B. *Protein Eng* 9: 977-86
- Feng R, Rampon C, Tang YP, Shrom D, Jin J, et al. 2001. Deficient neurogenesis in forebrain-specific presenilin-1 knockout mice is associated with reduced clearance of hippocampal memory traces. *Neuron* 32: 911-26

- Feng SJ, Nie K, Gan R, Huang J, Zhang YW, et al. 2014. Triggering receptor expressed on myeloid cells 2 variants are rare in Parkinson's disease in a Han Chinese cohort. *Neurobiol Aging* 35: 1780 e11-2
- Fernando P, Brunette S, Megeney LA. 2005. Neural stem cell differentiation is dependent upon endogenous caspase 3 activity. *FASEB J.* 19: 1671-73
- Figueroa DJ, Morris JA, Ma L, Kandpal G, Chen E, et al. 2002. Presenilin-dependent gamma-secretase activity modulates neurite outgrowth. *Neurobiol Dis* 9: 49-60
- Filippo TR, Galindo LT, Barnabe GF, Ariza CB, Mello LE, et al. 2013. CXCL12 N-terminal end is sufficient to induce chemotaxis and proliferation of neural stem/progenitor cells. *Stem Cell Res* 11: 913-25
- Filippov V, Kronenberg G, Pivneva T, Reuter K, Steiner B, et al. 2003. Subpopulation of nestin-expressing progenitor cells in the adult murine hippocampus shows electrophysiological and morphological characteristics of astrocytes. *Mol Cell Neurosci* 23: 373-82
- Finckh U, von der Kammer H, Velden J, Michel T, Andresen B, et al. 2000. Genetic association of a cystatin C gene polymorphism with late-onset Alzheimer disease. *Arch Neurol* 57: 1579-83
- Finelli D, Rollinson S, Harris J, Jones M, Richardson A, et al. 2014. TREM2 analysis and increased risk of Alzheimer's disease. *Neurobiol Aging*
- Fiore F, Zambrano N, Minopoli G, Donini V, Duilio A, Russo T. 1995. The regions of the Fe65 protein homologous to the phosphotyrosine interaction/phosphotyrosine binding domain of Shc bind the intracellular domain of the Alzheimer's amyloid precursor protein. *J Biol Chem* 270: 30853-6
- Fiske BK, Brunjes PC. 2001. Cell death in the developing and sensory-deprived rat olfactory bulb. *J Comp Neurol* 431: 311-9
- Florio M, Leto K, Muzio L, Tinterri A, Badaloni A, et al. 2012. Neurogenin 2 regulates progenitor cell-cycle progression and Purkinje cell dendritogenesis in cerebellar development. *Development* 139: 2308-20
- Fluhrer R, Capell A, Westmeyer G, Willem M, Hartung B, et al. 2002. A non-amyloidogenic function of BACE-2 in the secretory pathway. *Journal of neurochemistry* 81: 1011-20
- Fode C, Gradwohl G, Morin X, Dierich A, LeMeur M, et al. 1998. The bHLH protein NEUROGENIN 2 is a determination factor for epibranchial placode-derived sensory neurons. *Neuron* 20: 483-94
- Fode C, Ma Q, Casarosa S, Ang SL, Anderson DJ, Guillemot F. 2000. A role for neural determination genes in specifying the dorsoventral identity of telencephalic neurons. *Genes Dev* 14: 67-80
- Fonseca MB, Sola S, Xavier JM, Dionisio PA, Rodrigues CM. 2013. Amyloid beta peptides promote autophagy-dependent differentiation of mouse neural stem cells: Abeta-mediated neural differentiation. *Mol Neurobiol* 48: 829-40
- Forstl H, Kurz A. 1999. Clinical features of Alzheimer's disease. *Eur Arch Psychiatry Clin Neurosci* 6
- Francis R, McGrath G, Zhang J, Ruddy DA, Sym M, et al. 2002. aph-1 and pen-2 are required for Notch pathway signaling, gamma-secretase cleavage of betaAPP, and presenilin protein accumulation. *Dev Cell* 3: 85-97

- Frank S, Burbach GJ, Bonin M, Walter M, Streit W, et al. 2008. TREM2 is upregulated in amyloid plaque-associated microglia in aged APP23 transgenic mice. *Glia* 56: 1438-47
- Frautschy SA, Yang F, Irrizarry M, Hyman B, Saido TC, et al. 1998. Microglial response to amyloid plaques in APPsw transgenic mice. *Am J Pathol* 152: 307-17
- Frenkel D, Solomon B, Benhar I. 2000. Modulation of Alzheimer's beta-amyloid neurotoxicity by site-directed single-chain antibody. *J Neuroimmunol* 106: 23-31
- Freude KK, Penjwini M, Davis JL, LaFerla FM, Blurton-Jones M. 2011. Soluble amyloid precursor protein induces rapid neural differentiation of human embryonic stem cells. *J Biol Chem* 286: 24264-74
- Freund C, Ward-van Oostwaard D, Monshouwer-Kloots J, van den Brink S, van Rooijen M, et al. 2008. Insulin redirects differentiation from cardiogenic mesoderm and endoderm to neuroectoderm in differentiating human embryonic stem cells. *stem cells* 26: 724-33
- Frielingsdorf H, Simpson DR, Thal LJ, Pizzo DP. 2007. Nerve growth factor promotes survival of new neurons in the adult hippocampus. *Neurobiol Dis* 26: 47-55
- Frinchi M, Bonomo A, Trovato-Salinaro A, Condorelli DF, Fuxe K, et al. 2008. Fibroblast growth factor-2 and its receptor expression in proliferating precursor cells of the subventricular zone in the adult rat brain. *Neurosci Lett* 447: 20-5
- Frommer W, Junge B, Muller L, Schmidt D, Truscheit E. 1979. [New enzyme inhibitors from microorganisms (author's transl)]. *Planta Med* 35: 195-217
- Frost EE, Nielsen JA, Le TQ, Armstrong RC. 2003. PDGF and FGF2 regulate oligodendrocyte progenitor responses to demyelination. *J Neurobiol* 54: 457-72
- Fryer JD, Simmons K, Parsadanian M, Bales KR, Paul SM, et al. 2005. Human apolipoprotein E4 alters the amyloid-beta 40:42 ratio and promotes the formation of cerebral amyloid angiopathy in an amyloid precursor protein transgenic model. *J Neurosci* 25: 2803-10
- Fuentealba LC, Obernier K, Alvarez-Buylla A. 2012. Adult neural stem cells bridge their niche. *Cell Stem Cell* 10: 698-708
- Fukuchi K, Deeb SS, Kamino K, Ogburn CE, Snow AD, et al. 1992. Increased expression of beta-amyloid protein precursor and microtubule-associated protein tau during the differentiation of murine embryonal carcinoma cells. *Journal of neurochemistry* 58: 1863-73
- Fukuda S, Kato F, Tozuka Y, Yamaguchi M, Miyamoto Y, Hisatsune T. 2003. Two distinct subpopulations of nestin-positive cells in adult mouse dentate gyrus. *J Neurosci* 23: 9357-66
- Fukumoto H, Cheung BS, Hyman BT, Irizarry MC. 2002. Beta-secretase protein and activity are increased in the neocortex in Alzheimer disease. *Arch Neurol* 59: 1381-9
- Gadadhar A, Marr R, Lazarov O. 2011. Presenilin-1 regulates neural progenitor cell differentiation in the adult brain. *J Neurosci* 31: 2615-23
- Gakhar-Koppole N, Hundeshagen P, Mandl C, Weyer SW, Allinquant B, et al. 2008. Activity requires soluble amyloid precursor protein alpha to promote

- neurite outgrowth in neural stem cell-derived neurons via activation of the MAPK pathway. *Eur J Neurosci* 28: 871-82
- Galichet C, Guillemot F, Parras CM. 2008. Neurogenin 2 has an essential role in development of the dentate gyrus. *Development* 135: 2031-41
- Galli R. 2013. The neurosphere assay applied to neural stem cells and cancer stem cells. *Methods Mol Biol* 986: 267-77
- Galvao RP, Garcia-Verdugo JM, Alvarez-Buylla A. 2008. Brain-derived neurotrophic factor signaling does not stimulate subventricular zone neurogenesis in adult mice and rats. *J Neurosci* 28: 13368-83
- Galvez-Contreras AY, Gonzalez-Castaneda RE, Luquin S, Gonzalez-Perez O. 2012. Role of fibroblast growth factor receptors in astrocytic stem cells. *Curr Signal Transduct Ther* 7: 81-86
- Games D, Adams D, Alessandrini R, Barbour R, Berthelette P, et al. 1995. Alzheimer-type neuropathology in transgenic mice overexpressing V717F beta-amyloid precursor protein. *Nature* 373: 523-7
- Ganat Y, Soni S, Chacon M, Schwartz ML, Vaccarino FM. 2002. Chronic hypoxia up-regulates fibroblast growth factor ligands in the perinatal brain and induces fibroblast growth factor-responsive radial glial cells in the subependymal zone. *Neuroscience* 112: 977-91
- Gao X, Enikolopov G, Chen J. 2009. Moderate traumatic brain injury promotes proliferation of quiescent neural progenitors in the adult hippocampus. *Exp Neurol* 219: 516-23
- Garcia-Segura LM, Perez J, Pons S, Rejas MT, Torres-Aleman I. 1991. Localization of insulin-like growth factor I (IGF-I)-like immunoreactivity in the developing and adult rat brain. *Brain Res* 560: 167-74
- Gardella JE, Ghiso J, Gorgone GA, Marratta D, Kaplan AP, et al. 1990. Intact Alzheimer amyloid precursor protein (APP) is present in platelet membranes and is encoded by platelet mRNA. *Biochem Biophys Res Commun* 173: 1292-8
- Gaspard N, Vanderhaeghen P. 2010. Mechanisms of neural specification from embryonic stem cells. *Curr Opin Neurobiol* 20: 37-43
- Gattazzo F, Urciuolo A, Bonaldo P. 2014. Extracellular matrix: a dynamic microenvironment for stem cell niche. *Biochim Biophys Acta* 1840: 2506-19
- Gauthier S, Kaur G, Mi W, Tizon B, Levy E. 2011. Protective mechanisms by cystatin C in neurodegenerative diseases. *Front Biosci (Schol Ed)* 3: 541-54
- Gauthier S, Panisset M, Nalbantoglu J, Poirier J. 1997. Alzheimer's disease: current knowledge, management and research. *Cmaj* 157: 1047-52
- Ge S, Goh EL, Sailor KA, Kitabatake Y, Ming GL, Song H. 2006. GABA regulates synaptic integration of newly generated neurons in the adult brain. *Nature* 439: 589-93
- Ge S, Sailor KA, Ming GL, Song H. 2008. Synaptic integration and plasticity of new neurons in the adult hippocampus. *J Physiol* 586: 3759-65
- Ge S, Yang CH, Hsu KS, Ming GL, Song H. 2007. A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron* 54: 559-66

- Gentleman SM, Graham DI, Roberts GW. 1993. Molecular pathology of head trauma: altered beta APP metabolism and the aetiology of Alzheimer's disease. *Prog Brain Res* 96: 237-46
- Gheusi G, Cremer H, McLean H, Chazal G, Vincent JD, Lledo PM. 2000. Importance of newly generated neurons in the adult olfactory bulb for odor discrimination. *Proc Natl Acad Sci U S A* 97: 1823-8
- Ghosal K, Stathopoulos A, Pimplikar SW. 2010. APP intracellular domain impairs adult neurogenesis in transgenic mice by inducing neuroinflammation. *PLoS One* 5: e11866
- Ghosal K, Vogt DL, Liang M, Shen Y, Lamb BT, Pimplikar SW. 2009. Alzheimer's disease-like pathological features in transgenic mice expressing the APP intracellular domain. *Proc Natl Acad Sci U S A* 106: 18367-72
- Giacobini E. 1998. Invited review: Cholinesterase inhibitors for Alzheimer's disease therapy: from tacrine to future applications. *Neurochem Int* 32: 413-9
- Giannakopoulos P, Herrmann FR, Bussière T, Bouras C, Kövari E, et al. 2003. Tangle and neuron numbers, but not amyloid load, predict cognitive status in Alzheimer's disease. *Neurology* 60
- Gianotti-Sommer A, Rozelle SS, Sullivan S, Mills JA, Park SM, et al. 2008. Generation of human induced pluripotent stem cells from peripheral blood using the STEMCCA lentiviral vector In *StemBook*. Cambridge (MA)
- Gil-Perotin S, Duran-Moreno M, Cebrian-Silla A, Ramirez M, Garcia-Belda P, Garcia-Verdugo JM. 2013. Adult neural stem cells from the subventricular zone: a review of the neurosphere assay. *Anat Rec (Hoboken)* 296: 1435-52
- Gilberto L, Zhou D, Weldon R, Tamagno E, De Luca P, et al. 2008. Evidence that the Amyloid beta Precursor Protein-intracellular domain lowers the stress threshold of neurons and has a "regulated" transcriptional role. *Mol Neurodegener* 3: doi:10.1186/750-326-3-12
- Giuffrida ML, Caraci F, Pignataro B, Cataldo S, De Bona P, et al. 2009. Beta-amyloid monomers are neuroprotective. *J Neurosci* 29: 10582-7
- Glenner GG, Wong CW. 1984. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* 120: 885-90
- Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, et al. 1991. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349: 704-6
- Golde TE, Eckman CB, Younkin SG. 2000. Biochemical detection of Abeta isoforms: implications for pathogenesis, diagnosis, and treatment of Alzheimer's disease. *Biochim Biophys Acta* 1502: 172-87
- Gomez-Gaviro MV, Scott CE, Sesay AK, Matheu A, Booth S, et al. 2012. Betacellulin promotes cell proliferation in the neural stem cell niche and stimulates neurogenesis. *Proc Natl Acad Sci U S A* 109: 1317-22
- Gomez-Isla T, Hollister R, West H, Mui S, Growdon JH, et al. 1997. Neuronal loss correlates with but exceeds neurofibrillary tangles in Alzheimer's disease. *Annals of neurology* 41: 17-24
- Gomez-Lamarca MJ, Cobreros-Reguera L, Ibanez-Jimenez B, Palacios IM, Martin-Bermudo MD. 2014. Integrins regulate epithelial cell differentiation by modulating Notch activity. *J Cell Sci*

- Gonzalez-Perez O, Jauregui-Huerta F, Galvez-Contreras AY. 2010. Immune system modulates the function of adult neural stem cells. *Curr Immunol Rev* 6: 167-73
- Gonzalez-Perez O, Quinones-Hinojosa A. 2010. Dose-dependent effect of EGF on migration and differentiation of adult subventricular zone astrocytes. *Glia* 58: 975-83
- Gonzalez-Perez O, Romero-Rodriguez R, Soriano-Navarro M, Garcia-Verdugo JM, Alvarez-Buylla A. 2009. Epidermal growth factor induces the progeny of subventricular zone type B cells to migrate and differentiate into oligodendrocytes. *stem cells* 27: 2032-43
- Gopinath S, Malla R, Alapati K, Gorantla B, Gujrati M, et al. 2013. Cathepsin B and uPAR regulate self-renewal of glioma-initiating cells through GLI-regulated Sox2 and Bmi1 expression. *Carcinogenesis* 34: 550-59
- Gordon MN, King DL, Diamond DM, Jantzen PT, Boyett KV, et al. 2001. Correlation between cognitive deficits and Abeta deposits in transgenic APP+PS1 mice. *Neurobiol Aging* 22: 377-85
- Gorelick PB, Scuteri A, Black SE, Decarli C, Greenberg SM, et al. 2011. Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the american heart association/american stroke association. *Stroke* 42: 2672-713
- Gotz J, Chen F, van Dorpe J, Nitsch RM. 2001. Formation of neurofibrillary tangles in P301 τ transgenic mice induced by A β 42 fibrils. *Science* 293: 1491-5
- Gotz M, Huttner WB. 2005. The cell biology of neurogenesis. *Nat Rev Mol Cell Biol* 6: 777-88
- Gould E, Tanapat P, McEwen BS, Flugge G, Fuchs E. 1998. Proliferation of granule cell precursors in the dentate gyrus of adult monkeys is diminished by stress. *Proc Natl Acad Sci U S A* 95: 3168-71
- Goulet B, Sansregret L, Leduy L, Bogyo M, Weber E, et al. 2007. Increased expression and activity of nuclear cathepsin L in cancer cells suggests a novel mechanism of cell transformation. *Mol Cancer Res* 5: 899-907
- Goutte C, Tsunozaki M, Hale VA, Priess JR. 2002. APH-1 is a multipass membrane protein essential for the Notch signaling pathway in *Caenorhabditis elegans* embryos. *Proc Natl Acad Sci U S A* 99: 775-9
- Green KA, Lund LR. 2005. ECM degrading proteases and tissue remodelling in the mammary gland. *Bioessays* 27: 894-903
- Green RC, Schneider LS, Amato DA, Beelen AP, Wilcock G, et al. 2009. Effect of tarenflurbil on cognitive decline and activities of daily living in patients with mild Alzheimer disease: a randomized controlled trial. *Jama* 302: 2557-64
- Greenberg DA. 2007. Neurogenesis and stroke. *CNS Neurol Disord Drug Targets* 6: 321-5
- Greenberg SM, Gurol ME, Rosand J, Smith EE. 2004. Amyloid angiopathy-related vascular cognitive impairment. *Stroke* 35: 2616-9
- Grehan S, Tse E, Taylor JM. 2001. Two distal downstream enhancers direct expression of the human apolipoprotein E gene to astrocytes in the brain. *J Neurosci* 21: 812-22

- Gritti A, Parati EA, Cova L, Frolichsthal P, Galli R, et al. 1996. Multipotential stem cells from the adult mouse brain proliferate and self-renew in response to basic fibroblast growth factor. *J Neurosci* 16: 1091-100
- Groves JO, Leslie I, Huang GJ, McHugh SB, Taylor A, et al. 2013. Ablating adult neurogenesis in the rat has no effect on spatial processing: evidence from a novel pharmacogenetic model. *PLoS Genet* 9: e1003718
- Grubb MS, Nissant A, Murray K, Lledo PM. 2008. Functional maturation of the first synapse in olfaction: development and adult neurogenesis. *J Neurosci* 28: 2919-32
- Grundke-Iqbal I, Iqbal K, Quinlan M, Tung YC, Zaidi MS, Wisniewski HM. 1986. Microtubule-associated protein tau. A component of Alzheimer paired helical filaments. *J Biol Chem* 261: 6084-9
- Grzelakowska-Sztubert B. 1998. [Molecular mechanisms of apoptosis induced by activation of membrane receptors from the TNF-R superfamily]. *Postepy Biochem* 44: 8-21
- Grzonka Z, Jankowska E, Kasprzykowski F, Kasprzykowska R, Lankiewicz L, et al. 2001. Structural studies of cysteine proteases and their inhibitors. *Acta Biochim Pol* 48: 1-20
- Gu Z, Cheng J, Zhong P, Qin L, Liu W, Yan Z. 2014. Abeta Selectively Impairs mGluR7 Modulation of NMDA Signaling in Basal Forebrain Cholinergic Neurons: Implication in Alzheimer's Disease. *J Neurosci* 34: 13614-28
- Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, et al. 2013. TREM2 variants in Alzheimer's disease. *N Engl J Med* 368: 117-27
- Guruharsha KG, Kankel MW, Artavanis-Tsakonas S. 2012. The Notch signalling system: recent insights into the complexity of a conserved pathway. *Nat Rev Genet* 13: 654-66
- Haan J, Maat-Schieman ML, van Duinen SG, Jensson O, Thorsteinsson L, Roos RA. 1994. Co-localization of beta/A4 and cystatin C in cortical blood vessels in Dutch, but not in Icelandic hereditary cerebral hemorrhage with amyloidosis. *Acta neurologica Scandinavica* 89: 367-71
- Haass C, Hung AY, Selkoe DJ. 1991. Processing of beta-amyloid precursor protein in microglia and astrocytes favors an internal localization over constitutive secretion. *J Neurosci* 11: 3783-93
- Hack MA, Sugimori M, Lundberg C, Nakafuku M, Gotz M. 2004. Regionalization and fate specification in neurospheres: the role of Olig2 and Pax6. *Mol Cell Neurosci* 25: 664-78
- Hakansson K, Huh C, Grubb A, Karlsson S, Abrahamson M. 1996. Mouse and rat cystatin C: Escherichia coli production, characterization and tissue distribution. *Comp Biochem Physiol B Biochem Mol Biol* 114: 303-11
- Hamos JE, DeGennaro LJ, Drachman DA. 1989. Synaptic loss in Alzheimer's disease and other dementias. *Neurology* 39: 355-61
- Han J, Wang B, Xiao Z, Gao Y, Zhao Y, et al. 2008. Mammalian target of rapamycin (mTOR) is involved in the neuronal differentiation of neural progenitors induced by insulin. *Mol Cell Neurosci* 39: 118-24
- Hanada K, Tamai M, Ohmura S, Sawada J, Seki T, Tanaka I. 1978a. Inhibitory activities of E-64 derivatives on papain. *Agric. Biol. Chem.* 42
- Hanada K, Tamai M, Yamagishi M, Ohmura S, Sawada J, Tanaka I. 1978b. Structure and synthesis of E-64, a new thiol protease inhibitor *Agric. Biol. Chem.* 42: 523-28

- Hand R, Bortone D, Mattar P, Nguyen L, Heng JI, et al. 2005. Phosphorylation of Neurogenin2 specifies the migration properties and the dendritic morphology of pyramidal neurons in the neocortex. *Neuron* 48: 45-62
- Hand R, Polleux F. 2011. Neurogenin2 regulates the initial axon guidance of cortical pyramidal neurons projecting medially to the corpus callosum. *Neural Dev* 6: 30
- Hanley J, Rastegarlar G, Nathwani AC. 2010. An introduction to induced pluripotent stem cells. *Br J Haematol* 151: 16-24
- Hansson SF, Andreasson U, Wall M, Skoog I, Andreasen N, et al. 2009. Reduced levels of amyloid-beta-binding proteins in cerebrospinal fluid from Alzheimer's disease patients. *Journal of Alzheimer's disease : JAD* 16: 389-97
- Hardy J, Selkoe DJ. 2002. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353-6
- Hargus G, Cooper O, Deleidi M, Levy A, Lee K, et al. 2010. Differentiated Parkinson patient-derived induced pluripotent stem cells grow in the adult rodent brain and reduce motor asymmetry in Parkinsonian rats. *Proc Natl Acad Sci U S A* 107: 15921-6
- Hartfuss E, Galli R, Heins N, Gotz M. 2001. Characterization of CNS precursor subtypes and radial glia. *Dev Biol* 229: 15-30
- Hasebe N, Fujita Y, Ueno M, Yoshimura K, Fujino Y, Yamashita T. 2013. Soluble beta-amyloid Precursor Protein Alpha binds to p75 neurotrophin receptor to promote neurite outgrowth. *PLoS One* 8: e82321
- Hasegawa A, Naruse M, Hitoshi S, Iwasaki Y, Takebayashi H, Ikenaka K. 2007. Regulation of glial development by cystatin C. *Journal of neurochemistry* 100: 12-22
- Haughey NJ, Liu D, Nath A, Borchard AC, Mattson MP. 2002. Disruption of neurogenesis in the subventricular zone of adult mice, and in human cortical neuronal precursor cells in culture, by amyloid beta-peptide: implications for the pathogenesis of Alzheimer's disease. *Neuromolecular Med* 1: 125-35
- Hayashi Y, Kashiwagi K, Ohta J, Nakajima M, Kawashima T, Yoshikawa K. 1994. Alzheimer amyloid protein precursor enhances proliferation of neural stem cells from fetal rat brain. *Biochem Biophys Res Commun* 205: 936-43
- Hayon Y, Dashevsky O, Shai E, Varon D, Leker RR. 2013. Platelet lysates stimulate angiogenesis, neurogenesis and neuroprotection after stroke. *Thromb Haemost* 110: 323-30
- He N, Jin WL, Lok KH, Wang Y, Yin M, Wang ZJ. 2013. Amyloid-beta(1-42) oligomer accelerates senescence in adult hippocampal neural stem/progenitor cells via formylpeptide receptor 2. *Cell Death Dis* 4: e924
- He X, Cooley K, Chung CH, Dashti N, Tang J. 2007. Apolipoprotein receptor 2 and X11 alpha/beta mediate apolipoprotein E-induced endocytosis of amyloid-beta precursor protein and beta-secretase, leading to amyloid-beta production. *J Neurosci* 27: 4052-60
- Heber S, Herms J, Gajic V, Hainfellner J, Aguzzi A, et al. 2000. Mice with combined gene knock-outs reveal essential and partially redundant functions of amyloid precursor protein family members. *J Neurosci* 20: 7951-63

- Hebert LE, Weuve J, Scherr PA, Evans DA. 2013. Alzheimer disease in the United States (2010-2050) estimated using the 2010 Census. *Neurology* 19: 1778-83
- Hebert SS, Serneels L, Tolia A, Craessaerts K, Derks C, et al. 2006. Regulated intramembrane proteolysis of amyloid precursor protein and regulation of expression of putative target genes. *EMBO Rep* 7: 739-45
- Heinrich PC, Behrmann I, Haan S, Hermanns HM, Muller-Newen G, Schaper F. 2003. Principles of interleukin (IL)-6-type cytokine signalling and its regulation. *Biochem J* 374: 1-20
- Heins N, Malatesta P, Cecconi F, Nakafuku M, Tucker KL, et al. 2002. Glial cells generate neurons: the role of the transcription factor Pax6. *Nature neuroscience* 5: 308-15
- Helgason CD, Miller CL. 2013. *Basic cell culture protocols*. New York: Humana Press. xiv, 550 p. pp.
- Hendriksen H, Datson NA, Ghijsen WE, van Vliet EA, da Silva FH, et al. 2001. Altered hippocampal gene expression prior to the onset of spontaneous seizures in the rat post-status epilepticus model. *Eur J Neurosci* 14: 1475-84
- Heng JI, Nguyen L, Castro DS, Zimmer C, Wildner H, et al. 2008. Neurogenin 2 controls cortical neuron migration through regulation of Rnd2. *Nature* 455: 114-8
- Heo C, Chang K-A, Choi HS, Kim H-S, Kim S, et al. 2007. Effects of the monomeric, oligomeric, and fibrillar A beta(42) peptides on the proliferation and differentiation of adult neural stem cells from subventricular zone. *J. Neurochem.* 102: 493-500
- Herman JP, Schafer MK, Young EA, Thompson R, Douglass J, et al. 1989. Evidence for hippocampal regulation of neuroendocrine neurons of the hypothalamo-pituitary-adrenocortical axis. *J Neurosci* 9: 3072-82
- Herms J, Anliker B, Heber S, Ring S, Fuhrmann M, et al. 2004. Cortical dysplasia resembling human type 2 lissencephaly in mice lacking all three APP family members. *Embo J* 23: 4106-15
- Hernandez-Rabaza V, Llorens-Martin M, Velazquez-Sanchez C, Ferragud A, Arcusa A, et al. 2009. Inhibition of adult hippocampal neurogenesis disrupts contextual learning but spares spatial working memory, long-term conditional rule retention and spatial reversal. *Neuroscience* 159: 59-68
- Herrmann N, Gauthier S. 2008. Diagnosis and treatment of dementia: 6. Management of severe Alzheimer disease. *Cmaj* 12: 1279-87
- Herzig MC, Winkler DT, Burgermeister P, Pfeifer M, Kohler E, et al. 2004. Abeta is targeted to the vasculature in a mouse model of hereditary cerebral hemorrhage with amyloidosis. *Nat Neurosci* 7: 954-60
- Hesse L, Behr D, Masters CL, Multhaup G. 1994. The beta A4 amyloid precursor protein binding to copper. *FEBS Lett* 349: 109-16
- Hetman M, Cavanaugh JE, Kimelman D, Xia Z. 2000. Role of glycogen synthase kinase-3beta in neuronal apoptosis induced by trophic withdrawal. *J Neurosci* 20: 2567-74
- Heuckeroth RO, Lampe PA, Johnson EM, Milbrandt J. 1998. Neurturin and GDNF promote proliferation and survival of enteric neuron and glial progenitors in vitro. *Dev Biol* 200: 116-29

- Hiroi M, Ohmori Y. 2003. The transcriptional coactivator CREB-binding protein cooperates with STAT1 and NF-kappa B for synergistic transcriptional activation of the CXC ligand 9/monokine induced by interferon-gamma gene. *J Biol Chem* 278: 651-60
- Ho A, Sudhof TC. 2004. Binding of F-spondin to amyloid-beta precursor protein: a candidate amyloid-beta precursor protein ligand that modulates amyloid-beta precursor protein cleavage. *Proc Natl Acad Sci U S A* 101: 2548-53
- Ho R, Ortiz D, Shea TB. 2001. Amyloid-beta promotes calcium influx and neurodegeneration via stimulation of L voltage-sensitive calcium channels rather than NMDA channels in cultured neurons. *J Alzheimers Dis* 3: 479-83
- Hoareau C, Borrell V, Soriano E, Krebs MO, Prochiantz A, Allinquant B. 2008. Amyloid precursor protein cytoplasmic domain antagonizes reelin neurite outgrowth inhibition of hippocampal neurons. *Neurobiol Aging* 29: 542-53
- Hodge RD, Kowalczyk TD, Wolf SA, Encinas JM, Rippey C, et al. 2008. Intermediate progenitors in adult hippocampal neurogenesis: Tbr2 expression and coordinate regulation of neuronal output. *J Neurosci* 28: 3707-17
- Hoe HS, Fu Z, Makarova A, Lee JY, Lu C, et al. 2009a. The effects of amyloid precursor protein on postsynaptic composition and activity. *J Biol Chem* 284: 8495-506
- Hoe HS, Lee HK, Pak DT. 2012. The upside of APP at synapses. *CNS Neurosci Ther* 18: 47-56
- Hoe HS, Lee KJ, Carney RS, Lee J, Markova A, et al. 2009b. Interaction of reelin with amyloid precursor protein promotes neurite outgrowth. *J Neurosci* 29: 7459-73
- Hoe HS, Tran TS, Matsuoka Y, Howell BW, Rebeck GW. 2006. DAB1 and Reelin effects on amyloid precursor protein and ApoE receptor 2 trafficking and processing. *J Biol Chem* 281: 35176-85
- Hoey SE, Williams RJ, Perkinton MS. 2009. Synaptic NMDA receptor activation stimulates alpha-secretase amyloid precursor protein processing and inhibits amyloid-beta production. *J Neurosci* 29: 4442-60
- Hoglinger GU, Rizk P, Muriel MP, Duyckaerts C, Oertel WH, et al. 2004. Dopamine depletion impairs precursor cell proliferation in Parkinson disease. *Nature neuroscience* 7: 726-35
- Holscher C. 1998. Possible causes of Alzheimer's disease: amyloid fragments, free radicals, and calcium homeostasis. *Neurobiol Dis* 5: 129-41
- Holsinger RM, Lee JS, Boyd A, Masters CL, Collins SJ. 2006. CSF BACE1 activity is increased in CJD and Alzheimer disease versus [corrected] other dementias. *Neurology* 67: 710-2
- Holtzman DM, Bales KR, Tenkova T, Fagan AM, Parsadanian M, et al. 2000. Apolipoprotein E isoform-dependent amyloid deposition and neuritic degeneration in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 97: 2892-7
- Holtzman DM, Herz J, Bu G. 2012. Apolipoprotein E and apolipoprotein E receptors: normal biology and roles in Alzheimer disease. *Cold Spring Harb Perspect Med* 2: a006312

- Homayouni R, Rice DS, Sheldon M, Curran T. 1999. Disabled-1 binds to the cytoplasmic domain of amyloid precursor-like protein 1. *J Neurosci* 19: 7507-15
- Hong J, Yoshida K, Rosner MR. 2002. Characterization of a cysteine proteinase inhibitor induced during neuronal cell differentiation. *J Neurochem* 81: 922-34
- Hook G, Hook V, Kindy M. 2011. The cysteine protease inhibitor, E64d, reduces brain amyloid-beta and improves memory deficits in Alzheimer's disease animal models by inhibiting cathepsin B, but not BACE1, beta-secretase activity. *Journal of Alzheimer's disease : JAD* 26: 387-408
- Hook V, Funkelstein L, Wegrzyn J, Bark S, Kindy M, Hook G. 2012. Cysteine Cathepsins in the secretory vesicle produce active peptides: Cathepsin L generates peptide neurotransmitters and cathepsin B produces beta-amyloid of Alzheimer's disease. *Biochim Biophys Acta* 1824: 89-104
- Hornsten A, Lieberthal J, Fadia S, Malins R, Ha L, et al. 2007. APL-1, a *Caenorhabditis elegans* protein related to the human beta-amyloid precursor protein, is essential for viability. *Proc Natl Acad Sci U S A* 104: 1971-6
- Hou WS, Li Z, Buttner FH, Bartnik E, Bromme D. 2003. Cleavage site specificity of cathepsin K toward cartilage proteoglycans and protease complex formation. *Biol Chem* 384: 891-7
- Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, et al. 1996. Correlative memory deficits, A β elevation, and amyloid plaques in transgenic mice. *Science* 274: 99-102
- Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, et al. 2006. AMPAR removal underlies A β -induced synaptic depression and dendritic spine loss. *Neuron* 52: 831-43
- Hsieh J. 2012. Orchestrating transcriptional control of adult neurogenesis. *Genes Dev* 26: 1010-21
- Hsieh J, Aimone JB, Kaspar BK, Kuwabara T, Nakashima K, Gage FH. 2004. IGF-I instructs multipotent adult neural progenitor cells to become oligodendrocytes. *J Cell Biol* 164: 111-22
- Hu X, Hicks CW, He W, Wong P, Macklin WB, et al. 2006. Bace1 modulates myelination in the central and peripheral nervous system. *Nature neuroscience* 9: 1520-5
- Hu X, Zhou X, He W, Yang J, Xiong W, et al. 2010. BACE1 deficiency causes altered neuronal activity and neurodegeneration. *J Neurosci* 30: 8819-29
- Hu Y, Fortini ME. 2003. Different cofactor activities in gamma-secretase assembly: evidence for a nicastrin-Aph-1 subcomplex. *J Cell Biol* 161: 685-90
- Hu Y, Hung AC, Cui H, Dawkins E, Bolos M, et al. 2013. Role of cystatin C in amyloid precursor protein-induced proliferation of neural stem/progenitor cells. *J Biol Chem* 288: 18853-62
- Hua Y, Zhao H, Lu X, Kong Y, Jin H. 2012. Meta-analysis of the cystatin C (CST3) gene G73A polymorphism and susceptibility to Alzheimer's disease. *Int J Neurosci* 122: 431-8
- Huang X, Liu J, Ketova T, Fleming JT, Grover VK, et al. 2010. Transventricular delivery of Sonic hedgehog is essential to cerebellar ventricular zone development. *Proc Natl Acad Sci U S A* 107: 8422-7

- Huangfu D, Anderson KV. 2005. Cilia and Hedgehog responsiveness in the mouse. *Proc Natl Acad Sci U S A* 102: 11325-30
- Huh CG, Hakansson K, Nathanson CM, Thorgeirsson UP, Jonsson N, et al. 1999. Decreased metastatic spread in mice homozygous for a null allele of the cystatin C protease inhibitor gene. *Mol Pathol* 52: 332-40
- Hung AY, Koo EH, Haass C, Selkoe DJ. 1992. Increased expression of beta-amyloid precursor protein during neuronal differentiation is not accompanied by secretory cleavage. *Proc Natl Acad Sci U S A* 89: 9439-43
- Hung AY, Selkoe DJ. 1994. Selective ectodomain phosphorylation and regulated cleavage of beta-amyloid precursor protein. *Embo J* 13: 534-42
- Huovila AP, Turner AJ, Peltto-Huikko M, Karkkainen I, Ortiz RM. 2005. Shedding light on ADAM metalloproteinases. *Trends Biochem Sci* 30: 413-22
- Hussain I, Powell D, Howlett DR, Tew DG, Meek TD, et al. 1999. Identification of a novel aspartic protease (Asp 2) as beta-secretase. *Mol Cell Neurosci* 14: 419-27
- Hyman BT, Marzloff K, Arriagada PV. 1993. The lack of accumulation of senile plaques or amyloid burden in Alzheimer's disease suggests a dynamic balance between amyloid deposition and resolution. *J Neuropathol Exp Neurol* 6: 594-600
- Imayoshi I, Sakamoto M, Ohtsuka T, Takao K, Miyakawa T, et al. 2008. Roles of continuous neurogenesis in the structural and functional integrity of the adult forebrain. *Nature neuroscience* 11: 1153-61
- Imayoshi I, Sakamoto M, Yamaguchi M, Mori K, Kageyama R. 2010. Essential roles of Notch signaling in maintenance of neural stem cells in developing and adult brains. *J Neurosci* 30: 3489-98
- Iosif RE, Ekdahl CT, Ahlenius H, Pronk CJ, Bonde S, et al. 2006. Tumor necrosis factor receptor 1 is a negative regulator of progenitor proliferation in adult hippocampal neurogenesis. *J Neurosci* 26: 9703-12
- Iqbal K, Alonso Adel C, Grundke-Iqbal I. 2008. Cytosolic abnormally hyperphosphorylated tau but not paired helical filaments sequester normal MAPs and inhibit microtubule assembly. *J Alzheimers Dis* 4: 365-70
- Iqbal K, Zaidi T, Bancher C, Grundke-Iqbal I. 1994. Alzheimer paired helical filaments. Restoration of the biological activity by dephosphorylation. *FEBS Lett* 349: 104-8
- Isaac JT, Ashby MC, McBain CJ. 2007. The role of the GluR2 subunit in AMPA receptor function and synaptic plasticity. *Neuron* 54: 859-71
- Islam O, Loo TX, Heese K. 2009. Brain-derived neurotrophic factor (BDNF) has proliferative effects on neural stem cells through the truncated TRK-B receptor, MAP kinase, AKT, and STAT-3 signaling pathways. *Curr Neurovasc Res* 6: 42-53
- Itagaki S, McGeer PL, Akiyama H, Zhu S, Selkoe D. 1989. Relationship of microglia and astrocytes to amyloid deposits of Alzheimer disease. *J Neuroimmunol* 24: 173-82
- Itoh T, Satou T, Nishida S, Tsubaki M, Hashimoto S, Ito H. 2009. Expression of amyloid precursor protein after rat traumatic brain injury. *Neurol Res* 31: 103-9
- Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. 1994. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-

- specific A beta monoclonals: evidence that an initially deposited species is A beta 42. *Neuron* 13: 45-53
- Jack CR, Lowe VJ, Weigand SD, Wiste HJ, Senjem ML, et al. 2009. Serial PIB and MRI in normal, mild cognitive impairment and Alzheimer's disease: implications for sequence of pathological events in Alzheimer's disease. *Brain* 132: 1365-65
- Jacobsen JS, Muenkel HA, Blume AJ, Vitek MP. 1991. A novel species-specific RNA related to alternatively spliced amyloid precursor protein mRNAs. *Neurobiol Aging* 12: 575-83
- Jacobsen KT, Iverfeldt K. 2009. Amyloid precursor protein and its homologues: a family of proteolysis-dependent receptors. *Cell Mol Life Sci* 66: 2299-318
- Jarrett JT, Berger EP, Lansbury PTJ. 1993 The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry* 32: 4693-7
- Jarriault S, Brou C, Logeat F, Schroeter EH, Kopan R, Israel A. 1995. Signalling downstream of activated mammalian Notch. *Nature* 377: 355-8
- Jayatissa MN, Henningsen K, West MJ, Wiborg O. 2009. Decreased cell proliferation in the dentate gyrus does not associate with development of anhedonic-like symptoms in rats. *Brain Res* 1290: 133-41
- Jellinger KA. 2002. Alzheimer disease and cerebrovascular pathology: an update. *J Neural Transm* 109: 813-36
- Jensen JB, Parmar M. 2006. Strengths and limitations of the neurosphere culture system. *Mol Neurobiol* 34: 153-61
- Jiang Q, Lee CY, Mandrekar S, Wilkinson B, Cramer P, et al. 2008. ApoE promotes the proteolytic degradation of Abeta. *Neuron* 58: 681-93
- Jiang T, Tan L, Zhu XC, Zhang QQ, Cao L, et al. 2014. Upregulation of TREM2 Ameliorates Neuropathology and Rescues Spatial Cognitive Impairment in a Transgenic Mouse Model of Alzheimer's Disease. *Neuropsychopharmacology* 39: 2949-62
- Jin K, Galvan V, Xie L, Mao XO, Gorostiza OF, et al. 2004a. Enhanced neurogenesis in Alzheimer's disease transgenic (PDGF-APP^{Sw,Ind}) mice. *Proc Natl Acad Sci U S A* 101: 13363-7
- Jin K, Mao XO, Sun Y, Xie L, Jin L, et al. 2002. Heparin-binding epidermal growth factor-like growth factor: hypoxia-inducible expression in vitro and stimulation of neurogenesis in vitro and in vivo. *J Neurosci* 22: 5365-73
- Jin K, Minami M, Xie L, Sun Y, Mao XO, et al. 2004b. Ischemia-induced neurogenesis is preserved but reduced in the aged rodent brain. *Aging Cell* 3: 373-7
- Jin K, Peel AL, Mao XO, Xie L, Cottrell BA, et al. 2004c. Increased hippocampal neurogenesis in Alzheimer's disease. *Proc Natl Acad Sci U S A* 101: 343-7
- Jin M, Shepardson N, Yang T, Chen G, Walsh D, Selkoe DJ. 2011. Soluble amyloid beta-protein dimers isolated from Alzheimer cortex directly induce Tau hyperphosphorylation and neuritic degeneration. *Proc Natl Acad Sci U S A*. 108: 5819-24
- Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, et al. 2013. Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med* 368: 107-16

- Joshi P, Liang JO, DiMonte K, Sullivan J, Pimplikar SW. 2009. Amyloid precursor protein is required for convergent-extension movements during Zebrafish development. *Dev Biol* 335: 1-11
- Joyce JA, Baruch A, Chehade K, Meyer-Morse N, Giraudo E, et al. 2004. Cathepsin cysteine proteases are effectors of invasive growth and angiogenesis during multistage tumorigenesis. *Cancer Cell* 5: 443-53
- Ju P, Zhang S, Yeap Y, Feng Z. 2012. Induction of neuronal phenotypes from NG2+ glial progenitors by inhibiting epidermal growth factor receptor in mouse spinal cord injury. *Glia* 60: 1801-14
- Judson RL, Babiarz JE, Venere M, Billelloch R. 2009. Embryonic stem cell-specific microRNAs promote induced pluripotency. *Nat Biotechnol* 27: 459-61
- Kadowaki T, Hara K, Yamauchi T, Terauchi Y, Tobe K, Nagai R. 2003. Molecular mechanism of insulin resistance and obesity. *Exp Biol Med (Maywood)* 228: 1111-7
- Kaesler SA, Herzig MC, Coomaraswamy J, Kilger E, Selenica ML, et al. 2007. Cystatin C modulates cerebral beta-amyloidosis. *Nat Genet* 39: 1437-9
- Kaether C, Lammich S, Edbauer D, Ertl M, Rietdorf J, et al. 2002. Presenilin-1 affects trafficking and processing of betaAPP and is targeted in a complex with nicastrin to the plasma membrane. *J Cell Biol* 158: 551-61
- Kajiwara K, Ogata S, Tanihara M. 2005. Promotion of neurite outgrowth from fetal hippocampal cells by TNF-alpha receptor 1-derived peptide. *Cell Transplant* 14: 665-72
- Kalyani A, Hobson K, Rao MS. 1997. Neuroepithelial stem cells from the embryonic spinal cord: isolation, characterization, and clonal analysis. *Dev Biol* 186: 202-23
- Kamei N, Tanaka N, Oishi Y, Hamasaki T, Nakanishi K, et al. 2007. BDNF, NT-3, and NGF released from transplanted neural progenitor cells promote corticospinal axon growth in organotypic cocultures. *Spine (Phila Pa 1976)* 32: 1272-8
- Kamimura D, Ishihara K, Hirano T. 2003. IL-6 signal transduction and its physiological roles: the signal orchestration model. *Rev Physiol Biochem Pharmacol* 149: 1-38
- Kandasamy M, Couillard-Despres S, Raber KA, Stephan M, Lehner B, et al. 2010. Stem cell quiescence in the hippocampal neurogenic niche is associated with elevated transforming growth factor-beta signaling in an animal model of Huntington disease. *J Neuropathol Exp Neurol* 69: 717-28
- Kaneko N, Okano H, Sawamoto K. 2006. Role of the cholinergic system in regulating survival of newborn neurons in the adult mouse dentate gyrus and olfactory bulb. *Genes Cells* 11: 1145-59
- Kanemoto S, Griffin J, Markham-Coultes K, Aubert I, Tandon A, et al. 2014. Proliferation, differentiation and amyloid-beta production in neural progenitor cells isolated from TgCRND8 mice. *Neuroscience* 261: 52-9
- Kang DE, Pietrzik CU, Baum L, Chevallier N, Merriam DE, et al. 2000. Modulation of amyloid beta-protein clearance and Alzheimer's disease susceptibility by the LDL receptor-related protein pathway. *The Journal of clinical investigation* 106: 1159-66
- Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, et al. 1987. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 325: 733-6

- Kaplan MS, Hinds JW. 1977. Neurogenesis in the adult rat: electron microscopic analysis of light radioautographs. *Science* 197: 1092-4
- Kaplan MS, McNelly NA, Hinds JW. 1985. Population dynamics of adult-formed granule neurons of the rat olfactory bulb. *J Comp Neurol* 239: 117-25
- Kato T, Heike T, Okawa K, Haruyama M, Shiraishi K, et al. 2006. A neurosphere-derived factor, cystatin C, supports differentiation of ES cells into neural stem cells. *Proc Natl Acad Sci U S A* 103: 6019-24
- Katsimpardi L, Litterman NK, Schein PA, Miller CM, Loffredo FS, et al. 2014. Vascular and neurogenic rejuvenation of the aging mouse brain by young systemic factors. *Science* 344: 630-4
- Kaur G, Levy E. 2012. Cystatin C in Alzheimer's disease. *Front Mol Neurosci* 5: 79
- Kawarabayashi T, Younkin LH, Saido TC, Shoji M, Ashe KH, Younkin SG. 2001. Age-dependent changes in brain, CSF, and plasma amyloid (beta) protein in the Tg2576 transgenic mouse model of Alzheimer's disease. *J Neurosci* 21: 372-81
- Kee N, Teixeira CM, Wang AH, Frankland PW. 2007. Preferential incorporation of adult-generated granule cells into spatial memory networks in the dentate gyrus. *Nat Neurosci* 10: 355-62
- Kele J, Simpicio N, Ferri AL, Mira H, Guillemot F, et al. 2006. Neurogenin 2 is required for the development of ventral midbrain dopaminergic neurons. *Development* 133: 495-505
- Kelsch W, Lin CW, Mosley CP, Lois C. 2009. A critical period for activity-dependent synaptic development during olfactory bulb adult neurogenesis. *J Neurosci* 29: 11852-8
- Kempermann G, Gast D, Kronenberg G, Yamaguchi M, Gage FH. 2003. Early determination and long-term persistence of adult-generated new neurons in the hippocampus of mice. *Development* 130: 391-9
- Kempermann G, Jessberger S, Steiner B, Kronenberg G. 2004. Milestones of neuronal development in the adult hippocampus. *Trends Neurosci* 27: 447-52
- Kempermann G, Kuhn HG, Gage FH. 1998. Experience-induced neurogenesis in the senescent dentate gyrus. *J Neurosci* 18: 3206-12
- Keohane A, Ryan S, Maloney E, Sullivan AM, Nolan YM. 2010. Tumour necrosis factor- α impairs neuronal differentiation but not proliferation of hippocampal neural precursor cells: Role of Hes1. *Mol Cell Neurosci* 43: 127-35
- Kernie SG, Parent JM. 2010. Forebrain neurogenesis after focal Ischemic and traumatic brain injury. *Neurobiol Dis* 37: 267-74
- Khandekar N, Lie KH, Sachdev PS, Sidhu KS. 2012. Amyloid precursor proteins, neural differentiation of pluripotent stem cells and its relevance to Alzheimer's disease. *Stem cells and development* 21: 997-1006
- Khatoon S, Grundke-Iqbal I, Iqbal K. 1992. Brain levels of microtubule associated protein tau are elevated in Alzheimer's disease: a radioimmunoslot-blot assay for nanograms of the protein. *J Neurochem* 2: 750-3
- Kibbey MC, Jucker M, Weeks BS, Neve RL, Van Nostrand WE, Kleinman HK. 1993. beta-Amyloid precursor protein binds to the neurite-promoting IKVAV site of laminin. *Proc Natl Acad Sci U S A* 90: 10150-3
- Kidd M. 1964. Alzheimer's Disease--an Electron Microscopical Study. *Brain* 87: 307-20

- Kiialainen A, Hovanes K, Paloneva J, Kopra O, Peltonen L. 2005. ap12 and Trem2, molecules involved in innate immunity and neurodegeneration, are co expressed in the CNS. *Neurobiol Dis* 18: 314-22
- Kim DY, Carey BW, Wang H, Ingano LA, Binshtok AM, et al. 2007a. BACE1 regulates voltage-gated sodium channels and neuronal activity. *Nat Cell Biol* 9: 755-64
- Kim EJ, Ables JL, Dickel LK, Eisch AJ, Johnson JE. 2011. Ascl1 (Mash1) defines cells with long-term neurogenic potential in subgranular and subventricular zones in adult mouse brain. *PLoS One* 6: e18472
- Kim EJ, Leung CT, Reed RR, Johnson JE. 2007b. In vivo analysis of Ascl1 defined progenitors reveals distinct developmental dynamics during adult neurogenesis and gliogenesis. *J Neurosci* 27: 12764-74
- Kim HS, Kim EM, Lee JP, Park CH, Kim S, et al. 2003. C-terminal fragments of amyloid precursor protein exert neurotoxicity by inducing glycogen synthase kinase-3 β expression. *Faseb J* 17: 1951-3
- Kim JY, Kim DH, Kim JH, Lee D, Jeon HB, et al. 2012. Soluble intracellular adhesion molecule-1 secreted by human umbilical cord blood-derived mesenchymal stem cell reduces amyloid-beta plaques. *Cell Death Differ* 19: 680-91
- Kim SU, de Vellis J. 2009. Stem cell-based cell therapy in neurological diseases: a review. *Journal of neuroscience research* 87: 2183-200
- Kim Y, Wang WZ, Comte I, Pastrana E, Tran PB, et al. 2010. Dopamine stimulation of postnatal murine subventricular zone neurogenesis via the D3 receptor. *Journal of neurochemistry* 114: 750-60
- Kimberly WT, Esler WP, Ye W, Ostaszewski BL, Gao J, et al. 2003a. Notch and the amyloid precursor protein are cleaved by similar gamma-secretase(s). *Biochemistry* 42: 137-44
- Kimberly WT, LaVoie MJ, Ostaszewski BL, Ye W, Wolfe MS, Selkoe DJ. 2003b. Gamma-secretase is a membrane protein complex comprised of presenilin, nicastrin, Aph-1, and Pen-2. *Proc Natl Acad Sci U S A* 100: 6382-7
- Kimberly WT, Zheng JB, Guenette SY, Selkoe DJ. 2001. The intracellular domain of the beta-amyloid precursor protein is stabilized by Fe65 and translocates to the nucleus in a notch-like manner. *J Biol Chem* 276: 40288-92
- Kimberly WT, Zheng JB, Town T, Flavell RA, Selkoe DJ. 2005. Physiological regulation of the beta-amyloid precursor protein signaling domain by c-Jun N-terminal kinase JNK3 during neuronal differentiation. *J Neurosci* 25: 5533-43
- Kingham PJ, Pocock JM. 2001. Microglial secreted cathepsin B induces neuronal apoptosis. *Journal of neurochemistry* 76: 1475-84
- Kiskinis E, Eggan K. 2010. Progress toward the clinical application of patient-specific pluripotent stem cells. *The Journal of clinical investigation* 120: 51-9
- Kitaguchi N, Takahashi Y, Tokushima Y, Shiojiri S, Ito H. 1988. Novel precursor of Alzheimer's disease amyloid protein shows protease inhibitory activity. *Nature* 331: 530-2
- Kiyota T, Ingraham KL, Jacobsen MT, Xiong H, Ikezu T. 2011. FGF2 gene transfer restores hippocampal functions in mouse models of Alzheimer's disease

- and has therapeutic implications for neurocognitive disorders. *Proc Natl Acad Sci U S A* 108: E1339-48
- Klaver DW, Wilce MC, Cui H, Hung AC, Gasperini R, et al. 2010. Is BACE1 a suitable therapeutic target for the treatment of Alzheimer's disease? Current strategies and future directions. *Biol Chem* 391: 849-59
- Klein C, Butt SJ, Machold RP, Johnson JE, Fishell G. 2005. Cerebellum- and forebrain-derived stem cells possess intrinsic regional character. *Development* 132: 4497-508
- Klein DM, Felsenstein KM, Brenneman DE. 2009. Cathepsins B and L differentially regulate amyloid precursor protein processing. *J Pharmacol Exp Ther* 328: 813-21
- Klevanski M, Saar M, Baumkotter F, Weyer SW, Kins S, Muller UC. 2014. Differential role of APP and APLPs for neuromuscular synaptic morphology and function. *Mol Cell Neurosci* 61: 201-10
- Knops J, Suomensaaari S, Lee M, McConlogue L, Seubert P, Sinha S. 1995. Cell-type and amyloid precursor protein-type specific inhibition of A beta release by bafilomycin A1, a selective inhibitor of vacuolar ATPases. *J Biol Chem* 270: 2419-22
- Knowles RB, Wyart C, Buldyrev SV, Cruz L, Urbanc B, et al. 1999. Plaque-induced neurite abnormalities: implications for disruption of neural networks in Alzheimer's disease. *Proc Natl Acad Sci U S A* 96: 5274-9
- Ko SY, Lin SC, Chang KW, Wong YK, Liu CJ, et al. 2004. Increased expression of amyloid precursor protein in oral squamous cell carcinoma. *Int J Cancer* 111: 727-32
- Kohl Z, Kandasamy M, Winner B, Aigner R, Gross C, et al. 2007. Physical activity fails to rescue hippocampal neurogenesis deficits in the R6/2 mouse model of Huntington's disease. *Brain Res* 1155: 24-33
- Kohli BM, Pflieger D, Mueller LN, Carbonetti G, Aebersold R, et al. 2012. Interactome of the amyloid precursor protein APP in brain reveals a protein network involved in synaptic vesicle turnover and a close association with Synaptotagmin-1. *J Proteome Res* 11: 4075-90
- Kohman RA, Rhodes JS. 2013. Neurogenesis, inflammation and behavior. *Brain Behav Immun* 27: 22-32
- Kohwi M, Osumi N, Rubenstein JL, Alvarez-Buylla A. 2005. Pax6 is required for making specific subpopulations of granule and periglomerular neurons in the olfactory bulb. *J Neurosci* 25: 6997-7003
- Koike H, Tomioka S, Sorimachi H, Saido TC, Maruyama K, et al. 1999. Membrane-anchored metalloprotease MDC9 has an alpha-secretase activity responsible for processing the amyloid precursor protein. *Biochem J* 343 Pt 2: 371-5
- Konietzko U. 2012. AICD nuclear signaling and its possible contribution to Alzheimer's disease. *Curr Alzheimer Res* 9: 200-16
- Koo EH, Sisodia SS, Archer DR, Martin LJ, Weidemann A, et al. 1990. Precursor of amyloid protein in Alzheimer disease undergoes fast anterograde axonal transport. *Proc Natl Acad Sci U S A* 87: 1561-5
- Koo EH, Squazzo SL. 1994. Evidence that production and release of amyloid beta-protein involves the endocytic pathway. *J Biol Chem* 269: 17386-9

- Koo EH, Squazzo SL, Selkoe DJ, Koo CH. 1996. Trafficking of cell-surface amyloid beta-protein precursor. I. Secretion, endocytosis and recycling as detected by labeled monoclonal antibody. *J Cell Sci* 109 (Pt 5): 991-8
- Kopitar-Jerala N. 2006. The role of cystatins in cells of the immune system. *FEBS Lett* 580: 6295-301
- Kornhuber J, Quack G. 1995. Cerebrospinal fluid and serum concentrations of the N-methyl-D-aspartate (NMDA) receptor antagonist memantine in man. *Neurosci Lett* 195: 137-9
- Korte M, Herrmann U, Zhang X, Draguhn A. 2012. The role of APP and APLP for synaptic transmission, plasticity, and network function: lessons from genetic mouse models. *Exp Brain Res* 217: 435-40
- Kosaka K, Aika Y, Toida K, Kosaka T. 2001. Structure of intraglomerular dendritic tufts of mitral cells and their contacts with olfactory nerve terminals and calbindin-immunoreactive type 2 periglomerular neurons. *J Comp Neurol* 440: 219-35
- Kosaka T, Kosaka K. 2011. "Interneurons" in the olfactory bulb revisited. *Neurosci Res* 69: 93-9
- Kosik KS, Joachim CL, Selkoe DJ. 1986. Microtubule-associated protein tau (tau) is a major antigenic component of paired helical filaments in Alzheimer disease. *Proc Natl Acad Sci U S A* 83: 4044-8
- Kouroupi G, Lavdas AA, Gaitanou M, Thomaïdou D, Stylianopoulou F, Matsas R. 2010. Lentivirus-mediated expression of insulin-like growth factor-I promotes neural stem/precursor cell proliferation and enhances their potential to generate neurons. *Journal of neurochemistry* 115: 460-74
- Kronenberg G, Reuter K, Steiner B, Brandt MD, Jessberger S, et al. 2003. Subpopulations of proliferating cells of the adult hippocampus respond differently to physiologic neurogenic stimuli. *J Comp Neurol* 467: 455-63
- Kuhn HG, Dickinson-Anson H, Gage FH. 1996. Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *J Neurosci* 16: 2027-33
- Kuhn HG, Winkler J, Kempermann G, Thal LJ, Gage FH. 1997. Epidermal growth factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain. *J Neurosci* 17: 5820-9
- Kuhn PH, Wang H, Dislich B, Colombo A, Zeitschel U, et al. 2010. ADAM10 is the physiologically relevant, constitutive alpha-secretase of the amyloid precursor protein in primary neurons. *Embo J* 29: 3020-32
- Kullmann DM, Lamsa KP. 2007. Long-term synaptic plasticity in hippocampal interneurons. *Nat. Rev. Neurosci.* 8: 687-99
- Kumada T, Hasegawa A, Iwasaki Y, Baba H, Ikenaka K. 2004. Isolation of cystatin C via functional cloning of astrocyte differentiation factors. *Dev Neurosci* 26: 68-76
- Kurtz A, Oh SJ. 2012. Age related changes of the extracellular matrix and stem cell maintenance. *Prev Med* 54 Suppl: S50-6
- Kuszczyk MA, Sanchez S, Pankiewicz J, Kim J, Duszczek M, et al. 2013. Blocking the interaction between apolipoprotein E and Abeta reduces intraneuronal accumulation of Abeta and inhibits synaptic degeneration. *Am J Pathol* 182: 1750-68

- Kwak YD, Brannen CL, Qu T, Kim HM, Dong X, et al. 2006a. Amyloid precursor protein regulates differentiation of human neural stem cells. *Stem cells and development* 15: 381-9
- Kwak YD, Choumkin E, Sugaya K. 2006b. Amyloid precursor protein is involved in staurosporine induced glial differentiation of neural progenitor cells. *Biochem Biophys Res Commun*. 344: 431-37
- Kwak YD, Hendrix BJ, Sugaya K. 2014. Secreted type of amyloid precursor protein induces glial differentiation by stimulating the BMP/Smad signaling pathway. *Biochem Biophys Res Commun* 447: 394-9
- Laaris N, Puche A, Ennis M. 2007. Complementary postsynaptic activity patterns elicited in olfactory bulb by stimulation of mitral/tufted and centrifugal fiber inputs to granule cells. *J Neurophysiol* 97: 296-306
- Lagace DC, Donovan MH, DeCarolis NA, Farnbauch LA, Malhotra S, et al. 2010. Adult hippocampal neurogenesis is functionally important for stress-induced social avoidance. *Proc Natl Acad Sci U S A* 107: 4436-41
- Lai A, Gibson A, Hopkins CR, Trowbridge IS. 1998. Signal-dependent trafficking of beta-amyloid precursor protein-transferrin receptor chimeras in madin-darby canine kidney cells. *J Biol Chem* 273: 3732-9
- Lai K, Kaspar BK, Gage FH, Schaffer DV. 2003. Sonic hedgehog regulates adult neural progenitor proliferation in vitro and in vivo. *Nature neuroscience* 6: 21-7
- Laird FM, Cai H, Savonenko AV, Farah MH, He K, et al. 2005. BACE1, a major determinant of selective vulnerability of the brain to amyloid-beta amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J Neurosci* 25: 11693-709
- Lamb BT, Sisodia SS, Lawler AM, Slunt HH, Kitt CA, et al. 1993. Introduction and expression of the 400 kilobase amyloid precursor protein gene in transgenic mice [corrected]. *Nat Genet* 5: 22-30
- Lammich S, Kojro E, Postina R, Gilbert S, Pfeiffer R, et al. 1999. Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc Natl Acad Sci U S A* 96: 3922-7
- Lan X, Chen Q, Wang Y, Jia B, Sun L, et al. 2012. TNF-alpha affects human cortical neural progenitor cell differentiation through the autocrine secretion of leukemia inhibitory factor. *PLoS One* 7: e50783
- Landen JW, Zhao Q, Cohen S, Borrie M, Woodward M, et al. 2013. Safety and pharmacology of a single intravenous dose of ponezumab in subjects with mild-to-moderate Alzheimer disease: a phase I, randomized, placebo-controlled, double-blind, dose-escalation study. *Clin Neuropharmacol* 36: 14-23
- Lannfelt L. 1996. Genetics of Alzheimer's disease. *Acta Neurol Scand Suppl* 168: 25-7
- Lannfelt L, Blennow K, Zetterberg H, Batsman S, Ames D, et al. 2008. Safety, efficacy, and biomarker findings of PBT2 in targeting Abeta as a modifying therapy for Alzheimer's disease: a phase IIa, double-blind, randomised, placebo-controlled trial. *Lancet Neurol* 7: 779-86
- Lannfelt L, Relkin NR, Siemers ER. 2014. Amyloid-ss-directed immunotherapy for Alzheimer's disease. *J Intern Med* 275: 284-95

- Lansbury PTJ. 1999. Evolution of amyloid: What normal protein folding may tell us about fibrillogenesis and disease. *Proc Natl Acad Sci U S A* 96: 3342-44
- Laplagne DA, Esposito MS, Piatti VC, Morgenstern NA, Zhao C, et al. 2006. Functional convergence of neurons generated in the developing and adult hippocampus. *PLoS Biol* 4: e409
- Laplagne DA, Kamienkowski JE, Esposito MS, Piatti VC, Zhao C, et al. 2007. Similar GABAergic inputs in dentate granule cells born during embryonic and adult neurogenesis. *Eur J Neurosci* 25: 2973-81
- Laske C. 2014. Phase 3 trials of solanezumab and bapineuzumab for Alzheimer's disease. *N Engl J Med* 370: 1459
- Lathia JD, Mattson MP, Cheng A. 2008. Notch: from neural development to neurological disorders. *Journal of neurochemistry* 107: 1471-81
- Lau WM, Qiu G, Helmeke DM, Lee TM, Tang SW, So KF. 2007. Corticosteroid decreases subventricular zone cell proliferation, which could be reversed by paroxetine. *Restor Neurol Neurosci* 25: 17-23
- Launer LJ, Petrovitch H, Ross GW, Markesbery W, White LR. 2008. AD brain pathology: vascular origins? Results from the HAAS autopsy study. *Neurobiol Aging* 29: 1587-90
- LaVoie MJ, Fraering PC, Ostaszewski BL, Ye W, Kimberly WT, et al. 2003. Assembly of the gamma-secretase complex involves early formation of an intermediate subcomplex of Aph-1 and nicastrin. *J Biol Chem* 278: 37213-22
- Lazarini F, Gabellec MM, Moigneu C, de Chaumont F, Olivo-Marin JC, Lledo PM. 2014. Adult neurogenesis restores dopaminergic neuronal loss in the olfactory bulb. *J Neurosci* 34: 14430-42
- Lazarov O, Demars MP. 2012. All in the Family: How the APPs Regulate Neurogenesis. *Front Neurosci* 6: 81
- Lazarov O, Marr RA. 2010. Neurogenesis and Alzheimer's disease: at the crossroads. *Exp Neurol* 223: 267-81
- Lee AS, Tang C, Rao MS, Weissman IL, Wu JC. 2013a. Tumorigenicity as a clinical hurdle for pluripotent stem cell therapies. *Nature medicine* 19: 998-1004
- Lee HW, Park JW, Sandagsuren EU, Kim KB, Yoo JJ, Chung SH. 2008a. Overexpression of APP stimulates basal and constitutive exocytosis in PC12 cells. *Neurosci Lett* 436: 245-9
- Lee IS, Jung K, Kim IS, Park KI. 2013b. Amyloid- β oligomers regulate the properties of human neural stem cells through GSK-3 β signaling. *Exp Mol Med*. 45: doi: 10.1038/emm.2013.125.
- Lee JA, Cole GJ. 2007. Generation of transgenic zebrafish expressing green fluorescent protein under control of zebrafish amyloid precursor protein gene regulatory elements. *Zebrafish* 4: 277-86
- Lee KJ, Moussa CE, Lee Y, Sung Y, Howell BW, et al. 2010. Beta amyloid-independent role of amyloid precursor protein in generation and maintenance of dendritic spines. *Neuroscience* 169: 344-56
- Lee MK, Borchelt DR, Kim G, Thinakaran G, Slunt HH, et al. 1997. Hyperaccumulation of FAD-linked presenilin 1 variants in vivo. *Nature medicine* 3: 756-60
- Lee MS, Kao SC, Lemere CA, Xia W, Tseng HC, et al. 2003. APP processing is regulated by cytoplasmic phosphorylation. *J Cell Biol* 163: 83-95

- Lee SF, Shah S, Li H, Yu C, Han W, Yu G. 2002. Mammalian A β -1 interacts with presenilin and nicastrin and is required for intramembrane proteolysis of amyloid-beta precursor protein and Notch. *J Biol Chem* 277: 45013-9
- Lee YH, Martin JM, Maple RL, Tharp WG, Pratley RE. 2009. Plasma amyloid-beta peptide levels correlate with adipocyte amyloid precursor protein gene expression in obese individuals. *Neuroendocrinology* 90: 383-90
- Lee YH, Tharp WG, Maple RL, Nair S, Permana PA, Pratley RE. 2008b. Amyloid precursor protein expression is upregulated in adipocytes in obesity. *Obesity (Silver Spring)* 16: 1493-500
- Lefebvre V, Dumitriu B, Penzo-Mendez A, Han Y, Pallavi B. 2007. Control of cell fate and differentiation by Sry-related high-mobility-group box (Sox) transcription factors. *Int J Biochem Cell Biol* 39: 2195-214
- Lefterov I, Fitz NF, Cronican A, Lefterov P, Staufenbiel M, Koldamova R. 2009. Memory deficits in APP23/Abca1+/- mice correlate with the level of A β oligomers. *ASN Neuro* 1
- Lehmann ML, Brachman RA, Martinowich K, Schloesser RJ, Herkenham M. 2013. Glucocorticoids orchestrate divergent effects on mood through adult neurogenesis. *J Neurosci* 33: 2961-72
- Lehre KP, Levy LM, Ottersen OP, Storm-Mathisen J, Danbolt NC. 1995. Differential expression of two glial glutamate transporters in the rat brain: quantitative and immunocytochemical observations. *J Neurosci* 15: 1835-53
- Lehtinen MK, Zappaterra MW, Chen X, Yang YJ, Hill AD, et al. 2011. The cerebrospinal fluid provides a proliferative niche for neural progenitor cells. *Neuron* 69: 893-905
- Lemaire HG, Salbaum JM, Multhaup G, Kang J, Bayney RM, et al. 1989. The PreA4(695) precursor protein of Alzheimer's disease A4 amyloid is encoded by 16 exons. *Nucleic Acids Res* 17: 517-22
- Lenarcic B, Krasovec M, Ritonja A, Olafsson I, Turk V. 1991. Inactivation of human cystatin C and kininogen by human cathepsin D. *FEBS Lett* 280: 211-5
- Lendahl U, Zimmerman LB, McKay RD. 1990. CNS stem cells express a new class of intermediate filament protein. *Cell* 60: 585-95
- Lepousez G, Nissant A, Bryant AK, Gheusi G, Greer CA, Lledo PM. 2014. Olfactory learning promotes input-specific synaptic plasticity in adult-born neurons. *Proc Natl Acad Sci U S A* 111: 13984-9
- Lepousez G, Valley MT, Lledo PM. 2013. The impact of adult neurogenesis on olfactory bulb circuits and computations. *Annu Rev Physiol* 75: 339-63
- Levy E, Carman MD, Fernandez-Madrid IJ, Power MD, Lieberburg I, et al. 1990. Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science* 248: 1124-6
- Levy E, Sastre M, Kumar A, Gallo G, Piccardo P, et al. 2001. Codeposition of cystatin C with amyloid-beta protein in the brain of Alzheimer disease patients. *J Neuropathol Exp Neurol* 60: 94-104
- Levy-Lahad E, Wasco W, Poorkaj P, Romano DM, Oshima J, et al. 1995a. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science* 269: 973-7
- Levy-Lahad E, Wijsman EM, Nemens E, Anderson L, Goddard KA, et al. 1995b. A familial Alzheimer's disease locus on chromosome 1. *Science* 269: 970-3

- Li J, Kanekiyo T, Shinohara M, Zhang Y, LaDu MJ, et al. 2012a. Differential regulation of amyloid-beta endocytic trafficking and lysosomal degradation by apolipoprotein E isoforms. *J Biol Chem* 287: 44593-601
- Li L, Liu C, Biechele S, Zhu Q, Song L, et al. 2013a. Location of transient ectodermal progenitor potential in mouse development. *Development* 140: 4533-43
- Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D. 2009. Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 62: 788-801
- Li S, Mattar P, Zinyk D, Singh K, Chaturvedi CP, et al. 2012b. GSK3 temporally regulates neurogenin 2 proneural activity in the neocortex. *J Neurosci* 32: 7791-805
- Li X, Xiao Z, Han J, Chen L, Xiao H, et al. 2013b. Promotion of neuronal differentiation of neural progenitor cells by using EGFR antibody functionalized collagen scaffolds for spinal cord injury repair. *Biomaterials* 34: 5107-16
- Lichtenthaler SF, Haass C, Steiner H. 2011. Regulated intramembrane proteolysis--lessons from amyloid precursor protein processing. *Journal of neurochemistry* 117: 779-96
- Lichtenthaler SF, Steiner H. 2007. Sheddases and intramembrane-cleaving proteases: RIPpers of the membrane. Symposium on regulated intramembrane proteolysis. *EMBO Rep* 8: 537-41
- Lie DC, Colamarino SA, Song HJ, Desire L, Mira H, et al. 2005. Wnt signalling regulates adult hippocampal neurogenesis. *Nature* 437: 1370-5
- Lim DA, Alvarez-Buylla A. 2014. Adult neural stem cells stake their ground. *Trends Neurosci* 37: 563-71
- Lim DA, Tramontin AD, Trevejo JM, Herrera DG, Garcia-Verdugo JM, Alvarez-Buylla A. 2000. Noggin antagonizes BMP signaling to create a niche for adult neurogenesis. *Neuron* 28: 713-26
- Lin H, Bhatia R, Lal R. 2001. Amyloid beta protein forms ion channels: implications for Alzheimer's disease pathophysiology. *Faseb J* 15: 2433-44
- Lin L, Georgievska B, Mattsson A, Isacson O. 1999. Cognitive changes and modified processing of amyloid precursor protein in the cortical and hippocampal system after cholinergic synapse loss and muscarinic receptor activation. *Proc Natl Acad Sci U S A* 96: 12108-13
- Lin X, Koelsch G, Wu S, Downs D, Dashti A, Tang J. 2000. Human aspartic protease memapsin 2 cleaves the beta-secretase site of beta-amyloid precursor protein. *Proc Natl Acad Sci U S A* 97: 1456-60
- Lindberg OR, Brederlau A, Jansson A, Nannmark U, Cooper-Kuhn C, Kuhn HG. 2012. Characterization of epidermal growth factor-induced dysplasia in the adult rat subventricular zone. *Stem cells and development* 21: 1356-66
- Lindvall O, Barker RA, Brustle O, Isacson O, Svendsen CN. 2012. Clinical translation of stem cells in neurodegenerative disorders. *Cell Stem Cell* 10: 151-5
- Lindvall O, Kokaia Z. 2010. Stem cells in human neurodegenerative disorders--time for clinical translation? *The Journal of clinical investigation* 120: 29-40
- Liu B, Rasool S, Yang Z, Glabe CG, Schreiber SS, et al. 2009. Amyloid-peptide vaccinations reduce {beta}-amyloid plaques but exacerbate vascular

- deposition and inflammation in the retina of Alzheimer's transgenic mice. *Am J Pathol* 175: 2099-110
- Liu ML, Zang T, Zou Y, Chang JC, Gibson JR, et al. 2013. Small molecules enable neurogenin 2 to efficiently convert human fibroblasts into cholinergic neurons. *Nat Commun* 4: 2183
- Liu Q, Zerbinatti CV, Zhang J, Hoe HS, Wang B, et al. 2007. Amyloid precursor protein regulates brain apolipoprotein E and cholesterol metabolism through lipoprotein receptor LRP1. *Neuron* 56: 66-78
- Liu X, Wang Q, Haydar TF, Bordey A. 2005. Nonsynaptic GABA signaling in postnatal subventricular zone controls proliferation of GFAP-expressing progenitors. *Nature neuroscience* 8: 1179-87
- Liu Z, Martin LJ. 2003. Olfactory bulb core is a rich source of neural progenitor and stem cells in adult rodent and human. *J Comp Neurol* 459: 368-91
- Lledo PM, Alonso M, Grubb MS. 2006. Adult neurogenesis and functional plasticity in neuronal circuits. *Nat Rev Neurosci* 7: 179-93
- Lois C, Alvarez-Buylla A. 1993. Proliferating subventricular zone cells in the adult mammalian forebrain can differentiate into neurons and glia. *Proc Natl Acad Sci U S A* 90: 2074-7
- Lois C, Alvarez-Buylla A. 1994. Long-distance neuronal migration in the adult mammalian brain. *Science* 264: 1145-8
- Lois C, Garcia-Verdugo JM, Alvarez-Buylla A. 1996. Chain migration of neuronal precursors. *Science* 271: 978-81
- Lopez-Juarez A, Remaud S, Hassani Z, Jolivet P, Pierre Simons J, et al. 2012. Thyroid hormone signaling acts as a neurogenic switch by repressing Sox2 in the adult neural stem cell niche. *Cell Stem Cell* 10: 531-43
- Lopez-Toledano MA, Shelanski ML. 2004. Neurogenic effect of beta-amyloid peptide in the development of neural stem cells. *J Neurosci* 24: 5439-44
- Lopez-Toledano MA, Shelanski ML. 2007. Increased neurogenesis in young transgenic mice overexpressing human APP(Sw, Ind). *Journal of Alzheimer's disease : JAD* 12: 229-40
- Louis SA, Reynolds BA. 2005. Generation and differentiation of neurospheres from murine embryonic day 14 central nervous system tissue. *Methods Mol Biol* 290: 265-80
- Lourenco FC, Galvan V, Fombonne J, Corset V, Llambi F, et al. 2009. Netrin-1 interacts with amyloid precursor protein and regulates amyloid-beta production. *Cell Death Differ* 16: 655-63
- Lozzo R. 1998. Matrix proteoglycans: from molecular design to cellular function. *Annu Rev Biochem* 67: 609-52
- Lu DC, Rabizadeh S, Chandra S, Shayya RF, Ellerby LM, et al. 2000. A second cytotoxic proteolytic peptide derived from amyloid beta-protein precursor. *Nature medicine* 6: 397-404
- Lu KT, Sun CL, Wo PY, Yen HH, Tang TH, et al. 2011. Hippocampal neurogenesis after traumatic brain injury is mediated by vascular endothelial growth factor receptor-2 and the Raf/MEK/ERK cascade. *J Neurotrauma* 28: 441-50
- Lue LF, Schmitz C, Walker DG. 2014a. What happens to microglial TREM2 in Alzheimer's disease: Immunoregulatory turned into immunopathogenic? *Neuroscience*

- Lue LF, Schmitz CT, Serrano G, Sue LI, Beach TG, Walker DG. 2014b. TREM2 Protein Expression Changes Correlate with Alzheimer's Disease Neurodegenerative Pathologies in Post-Mortem Temporal Cortices. *Brain Pathol*
- Lugert S, Basak O, Knuckles P, Haussler U, Fabel K, et al. 2010. Quiescent and active hippocampal neural stem cells with distinct morphologies respond selectively to physiological and pathological stimuli and aging. *Cell Stem Cell* 6: 445-56
- Lukasiuk K, Pirttila TJ, Pitkanen A. 2002. Upregulation of cystatin C expression in the rat hippocampus during epileptogenesis in the amygdala stimulation model of temporal lobe epilepsy. *Epilepsia* 43 Suppl 5: 137-45
- Lum M, Croze E, Wagner C, McLenachan S, Mitrovic B, Turnley AM. 2009. Inhibition of neurosphere proliferation by IFNgamma but not IFNbeta is coupled to neuronal differentiation. *J Neuroimmunol* 206: 32-8
- Lunn JS, Sakowski SA, Hur J, Feldman EL. 2011. Stem cell technology for neurodegenerative diseases. *Annals of neurology* 70: 353-61
- Luo WJ, Wang H, Li H, Kim BS, Shah S, et al. 2003. PEN-2 and APH-1 coordinately regulate proteolytic processing of presenilin 1. *J Biol Chem* 278: 7850-4
- Luo Y, Bolon B, Kahn S, Bennett BD, Babu-Khan S, et al. 2001. Mice deficient in BACE1, the Alzheimer's beta-secretase, have normal phenotype and abolished beta-amyloid generation. *Nature neuroscience* 4: 231-2
- Luu P, Sill OC, Gao L, Becker S, Wojtowicz JM, Smith DM. 2012. The role of adult hippocampal neurogenesis in reducing interference. *Behav Neurosci* 126: 381-91
- Lynch JR, Morgan D, Mance J, Matthew WD, Laskowitz DT. 2001. Apolipoprotein E modulates glial activation and the endogenous central nervous system inflammatory response. *J Neuroimmunol* 114: 107-13
- Ma DK, Ming GL, Song H. 2005. Glial influences on neural stem cell development: cellular niches for adult neurogenesis. *Curr Opin Neurobiol* 15: 514-20
- Ma HJ, Shan L, Zhou JL, Liu QH, Lu T, Sun S. 2012a. [Case-control study on cable-pin system in the treatment of olecranon fractures]. *Zhongguo Gu Shang* 25: 393-6
- Ma L, Hu B, Liu Y, Vermilyea SC, Liu H, et al. 2012b. Human embryonic stem cell-derived GABA neurons correct locomotion deficits in quinolinic acid-lesioned mice. *Cell Stem Cell* 10: 455-64
- Ma Q, Kintner C, Anderson DJ. 1996. Identification of neurogenin, a vertebrate neuronal determination gene. *Cell* 87: 43-52
- Ma QH, Bagnard D, Xiao ZC, Dawe GS. 2008a. A TAG on to the neurogenic functions of APP. *Cell Adh Migr* 2: 2-8
- Ma QH, Futagawa T, Yang WL, Jiang XD, Zeng L, et al. 2008b. A TAG1-APP signalling pathway through Fe65 negatively modulates neurogenesis. *Nat Cell Biol* 10: 283-94
- Ma YC, Song MR, Park JP, Henry Ho HY, Hu L, et al. 2008c. Regulation of motor neuron specification by phosphorylation of neurogenin 2. *Neuron* 58: 65-77
- Machado VM, Morte MI, Carreira BP, Azevedo MM, Takano J, et al. 2015. Involvement of calpains in adult neurogenesis: implications for stroke. *Front Cell Neurosci* 9: 22

- Maciewicz RA, Wotton SF. 1991. Degradation of cartilage matrix components by the cysteine proteinases, cathepsins B and L. *Biomed Biochim Acta* 50: 561-4
- Magara F, Muller U, Li ZW, Lipp HP, Weissmann C, et al. 1999. Genetic background changes the pattern of forebrain commissure defects in transgenic mice underexpressing the beta-amyloid-precursor protein. *Proc Natl Acad Sci U S A* 96: 4656-61
- Magister S, Kos J. 2013. Cystatins in immune system. *J Cancer* 4: 45-56
- Mahley RW, Weisgraber KH, Huang Y. 2006. Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci U S A* 103: 5644-51
- Mak AC, Pullinger CR, Tang LF, Wong JS, Deo RC, et al. 2014. Effects of the absence of apolipoprotein e on lipoproteins, neurocognitive function, and retinal function. *JAMA Neurol* 71: 1228-36
- Malatesta P, Hack MA, Hartfuss E, Kettenmann H, Klinkert W, et al. 2003. Neuronal or glial progeny: regional differences in radial glia fate. *Neuron* 37: 751-64
- Malatesta P, Hartfuss E, Gotz M. 2000. Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* 127: 5253-63
- Malberg JE, Eisch AJ, Nestler EJ, Duman RS. 2000. Chronic antidepressant treatment increases neurogenesis in adult rat hippocampus. *J Neurosci* 20: 9104-10
- Mandairon N, Sacquet J, Jourdan F, Didier A. 2006. Long-term fate and distribution of newborn cells in the adult mouse olfactory bulb: Influences of olfactory deprivation. *Neuroscience* 141: 443-51
- Mandelkow E, Mandelkow EM. 1995. Microtubules and microtubule-associated proteins. *Curr Opin Cell Biol* 7: 72-81
- Marei HE, Ahmed AE. 2013. Transcription factors expressed in embryonic and adult olfactory bulb neural stem cells reveal distinct proliferation, differentiation and epigenetic control. *Genomics* 101: 12-9
- Marlatt MW, Lucassen PJ. 2010. Neurogenesis and Alzheimer's disease: Biology and pathophysiology in mice and men. *Curr Alzheimer Res* 7: 113-25
- Marshak D, Gardner R, Gottlieb D. 2001. Stem cell biology. *Cold Spring Harbor Laboratory Press* 18: 399-400
- Marshall CA, Novitsch BG, Goldman JE. 2005. Olig2 directs astrocyte and oligodendrocyte formation in postnatal subventricular zone cells. *J Neurosci* 25: 7289-98
- Martinez-Canabal A. 2014. Reconsidering hippocampal neurogenesis in Alzheimer's disease. *Front Neurosci* 8: 147
- Martinez-Coria H, Green KN, Billings LM, Kitazawa M, Albrecht M, et al. 2010. Memantine improves cognition and reduces Alzheimer's-like neuropathology in transgenic mice. *Am J Pathol* 176: 870-80
- Martinez-Serrano A, Bjorklund A. 1996. Protection of the neostriatum against excitotoxic damage by neurotrophin-producing, genetically modified neural stem cells. *J Neurosci* 16: 4604-16
- Maruyama K, Ikeda S, Ishihara T, Allsop D, Yanagisawa N. 1990. Immunohistochemical characterization of cerebrovascular amyloid in 46

- autopsied cases using antibodies to beta protein and cystatin C. *Stroke* 21: 397-403
- Masliah E. 1995. Mechanisms of synaptic dysfunction in Alzheimer's disease. *Histol Histopathol* 10: 509-19
- Masliah E, Hansen L, Adame A, Crews L, Bard F, et al. 2005. Abeta vaccination effects on plaque pathology in the absence of encephalitis in Alzheimer disease. *Neurology* 64: 129-31
- Masliah E, Mallory M, Ge N, Saitoh T. 1992. Amyloid precursor protein is localized in growing neurites of neonatal rat brain. *Brain Res* 593: 323-8
- Masliah E, Terry RD, Mallory M, Alford M, Hansen LA. 1990. Diffuse plaques do not accentuate synapse loss in Alzheimer's disease. *Am J Pathol* 137: 1293-7
- Mason RW, Massey SD. 1992. Surface activation of pro-cathepsin L. *Biochem Biophys Res Commun* 189: 1659-66
- Masters CL, Simms G, Weinman NA, Multhaup G, McDonald BL, Beyreuther K. 1985. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci U S A* 82: 4245-9
- Matsumoto K, Mizoue K, Kitamura K, Tse WC, Huber CP, Ishida T. 1999. Structural basis of inhibition of cysteine proteases by E-64 and its derivatives. *Biopolymers* 51: 99-107
- Matsumoto K, Yamamoto D, Ohishi H, Tomoo K, Ishida T, et al. 1989. Mode of binding of E-64-c, a potent thiol protease inhibitor, to papain as determined by X-ray crystal analysis of the complex. *FEBS Lett* 245: 177-80
- Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE. 1992. beta-Amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity. *J Neurosci* 12: 376-89
- Mattson MP, Pedersen WA, Duan W, Culmsee C, Camandola S. 1999. Cellular and molecular mechanisms underlying perturbed energy metabolism and neuronal degeneration in Alzheimer's and Parkinson's diseases. *Ann N Y Acad Sci* 893: 154-75
- May PC, Lampert-Etchells M, Johnson SA, Poirier J, Masters JN, Finch CE. 1990. Dynamics of gene expression for a hippocampal glycoprotein elevated in Alzheimer's disease and in response to experimental lesions in rat. *Neuron* 5: 831-9
- McCormick WC, Kukull WA, van Belle G, Bowen JD, Teri L, Larson EB. 1994. Symptom patterns and comorbidity in the early stages of Alzheimer's disease. *J Am Geriatr Soc* 42: 517-21
- McGowan EB, Becker E, Detwiler TC. 1989. Inhibition of calpain in intact platelets by the thiol protease inhibitor E-64d. *Biochem Biophys Res Commun* 158: 432-5
- McNay EC. 2007. Insulin and ghrelin: peripheral hormones modulating memory and hippocampal function. *Curr Opin Pharmacol* 7: 628-32
- Mega MS, Cummings JL, Fiorello T, Gornbein J. 1996. The spectrum of behavioral changes in Alzheimer's disease. *Neurology* 46: 130-5
- Melchior B, Garcia AE, Hsiung BK, Lo KM, Doose JM, et al. 2010. Dual induction of TREM2 and tolerance-related transcript, Tmem176b, in amyloid transgenic mice: implications for vaccine-based therapies for Alzheimer's disease. *ASN Neuro* 2: e00037

- Menn B, Garcia-Verdugo JM, Yaschine C, Gonzalez-Perez O, Rowitch D, Alvarez-Buylla A. 2006. Origin of oligodendrocytes in the subventricular zone of the adult brain. *J Neurosci* 26: 7907-18
- Merkle FT, Tramontin AD, Garcia-Verdugo JM, Alvarez-Buylla A. 2004. Radial glia give rise to adult neural stem cells in the subventricular zone. *Proc Natl Acad Sci U S A* 101: 17528-32
- Messersmith DJ, Murtie JC, Le TQ, Frost EE, Armstrong RC. 2000. Fibroblast growth factor 2 (FGF2) and FGF receptor expression in an experimental demyelinating disease with extensive remyelination. *Journal of neuroscience research* 62: 241-56
- Mi W, Pawlik M, Sastre M, Jung SS, Radvinsky DS, et al. 2007. Cystatin C inhibits amyloid-beta deposition in Alzheimer's disease mouse models. *Nat Genet* 39: 1440-2
- Mich JK, Signer RA, Nakada D, Pineda A, Burgess RJ, et al. 2014. Prospective identification of functionally distinct stem cells and neurosphere-initiating cells in adult mouse forebrain. *Elife* 3: e02669
- Michaelson DM. 2014. ApoE4: The most prevalent yet understudied risk factor for Alzheimer's disease. *Alzheimers Dement*
- Michaud JP, Bellavance MA, Prefontaine P, Rivest S. 2013. Real-time in vivo imaging reveals the ability of monocytes to clear vascular amyloid beta. *Cell Rep* 5: 646-53
- Miles DK, Kernie SG. 2008. Hypoxic-ischemic brain injury activates early hippocampal stem/progenitor cells to replace vulnerable neuroblasts. *Hippocampus* 18: 793-806
- Milward EA, Papadopoulos R, Fuller SJ, Moir RD, Small D, et al. 1992. The amyloid protein precursor of Alzheimer's disease is a mediator of the effects of nerve growth factor on neurite outgrowth. *Neuron* 9: 129-37
- Miners JS, Baig S, Palmer J, Palmer LE, Kehoe PG, Love S. 2008. Abeta-degrading enzymes in Alzheimer's disease. *Brain Pathol* 18: 240-52
- Minogue AM, Stubbs AK, Frigerio CS, Boland B, Fadeeva JV, et al. 2009. gamma-secretase processing of APLP1 leads to the production of a p3-like peptide that does not aggregate and is not toxic to neurons. *Brain Res* 1262: 89-99
- Minokadeh A, Funkelstein L, Toneff T, Hwang SR, Beinfeld M, et al. 2010. Cathepsin L participates in dynorphin production in brain cortex, illustrated by protease gene knockout and expression. *Mol Cell Neurosci* 43: 98-107
- Mira H, Andreu Z, Suh H, Lie DC, Jessberger S, et al. 2010. Signaling through BMPR-IA regulates quiescence and long-term activity of neural stem cells in the adult hippocampus. *Cell Stem Cell* 7: 78-89
- Mirochnic S, Wolf S, Staufenbiel M, Kempermann G. 2009. Age effects on the regulation of adult hippocampal neurogenesis by physical activity and environmental enrichment in the APP23 mouse model of Alzheimer disease. *Hippocampus* 19: 1008-18
- Mirzadeh Z, Merkle FT, Soriano-Navarro M, Garcia-Verdugo JM, Alvarez-Buylla A. 2008. Neural stem cells confer unique pinwheel architecture to the ventricular surface in neurogenic regions of the adult brain. *Cell Stem Cell* 3: 265-78
- Mission JP, Takahashi T, Caviness VS, Jr. 1991. Ontogeny of radial and other astroglial cells in murine cerebral cortex. *Glia* 4: 138-48

- Miura K, Okada Y, Aoi T, Okada A, Takahashi K, et al. 2009. Variation in the safety of induced pluripotent stem cell lines. *Nat Biotechnol* 27: 743-5
- Miyake T, Gahara Y, Nakayama M, Yamada H, Uwabe K, Kitamura T. 1996. Up-regulation of cystatin C by microglia in the rat facial nucleus following axotomy. *Brain Res Mol Brain Res* 37: 273-82
- Miyamoto A, Wake H, Moorhouse AJ, Nabekura J. 2013. Microglia and synapse interactions: fine tuning neural circuits and candidate molecules. *Front Cell Neurosci* 7: 70
- Mizoguchi Y, Ishibashi H, Nabekura J. 2003. The action of BDNF on GABA(A) currents changes from potentiating to suppressing during maturation of rat hippocampal CA1 pyramidal neurons. *J Physiol* 548: 703-9
- Mok SS, Sberna G, Heffernan D, Cappai R, Galatis D, et al. 1997. Expression and analysis of heparin-binding regions of the amyloid precursor protein of Alzheimer's disease. *FEBS Lett* 415: 303-7
- Moles A, Tarrats N, Fernandez-Checa JC, Mari M. 2009. Cathepsins B and D drive hepatic stellate cell proliferation and promote their fibrogenic potential. *Hepatology* 49: 1297-307
- Monastero R, Camarda C, Cefalu AB, Caldarella R, Camarda LK, et al. 2005. No association between the cystatin C gene polymorphism and Alzheimer's disease: a case-control study in an Italian population. *Journal of Alzheimer's disease : JAD* 7: 291-5
- Monje ML, Toda H, Palmer TD. 2003. Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302: 1760-5
- Moody SA, Miller V, Spanos A, Frankfurter A. 1996. Developmental expression of a neuron-specific beta-tubulin in frog (*Xenopus laevis*): a marker for growing axons during the embryonic period. *J Comp Neurol* 364: 219-30
- Moolman DL, Vitolo OV, Vonsattel JP, Shelanski ML. 2004. Dendrite and dendritic spine alterations in Alzheimer models. *J Neurocytol* 33: 377-87
- Moon M, Cha MY, Mook-Jung I. 2014. Impaired hippocampal neurogenesis and its enhancement with ghrelin in 5XFAD mice. *Journal of Alzheimer's disease : JAD* 41: 233-41
- Moreno MM, Linster C, Escanilla O, Sacquet J, Didier A, Mandairon N. 2009. Olfactory perceptual learning requires adult neurogenesis. *Proc Natl Acad Sci U S A* 106: 17980-5
- Morley JE, Farr SA, Banks WA, Johnson SN, Yamada KA, Xu L. 2010. A physiological role for amyloid-beta protein: enhancement of learning and memory. *Journal of Alzheimer's disease : JAD* 19: 441-9
- Morris JC, Storandt M, McKeel DWJ, Rubin EH, Price JL, et al. 1996. Cerebral amyloid deposition and diffuse plaques in "normal" aging: Evidence for presymptomatic and very mild Alzheimer's disease. *Neurology* 46: 707-19
- Morris JC, Storandt M, Miller JP, McKeel DW, Price JL, et al. 2001. Mild cognitive impairment represents early-stage Alzheimer disease. *Arch Neurol* 58: 397-405
- Morshead CM, Reynolds BA, Craig CG, McBurney MW, Staines WA, et al. 1994. Neural stem cells in the adult mammalian forebrain: a relatively quiescent subpopulation of subependymal cells. *Neuron* 13: 1071-82
- Moss ML, Powell G, Miller MA, Edwards L, Qi B, et al. 2011. ADAM9 inhibition increases membrane activity of ADAM10 and controls alpha-secretase processing of amyloid precursor protein. *J Biol Chem* 286: 40443-51

- Moya KL, Benowitz LI, Schneider GE, Allinquant B. 1994. The amyloid precursor protein is developmentally regulated and correlated with synaptogenesis. *Dev Biol* 161: 597-603
- Moyse E, Segura S, Liard O, Mahaut S, Mechawar N. 2008. Microenvironmental determinants of adult neural stem cell proliferation and lineage commitment in the healthy and injured central nervous system. *Curr Stem Cell Res Ther* 3: 163-84
- Mu Y, Gage FH. 2011. Adult hippocampal neurogenesis and its role in Alzheimer's disease. *Mol Neurodegener* 6: 85
- Mucke L, Masliah E, Johnson WB, Ruppe MD, Alford M, et al. 1994. Synaptotrophic effects of human amyloid beta protein precursors in the cortex of transgenic mice. *Brain Res* 666: 151-67
- Mueller-Steiner S, Zhou Y, Arai H, Roberson ED, Sun B, et al. 2006. Anti-amyloidogenic and neuroprotective functions of cathepsin B: implications for Alzheimer's disease. *Neuron* 51: 703-14
- Mullan M, Crawford F, Axelman K, Houlden H, Lilius L, et al. 1992. A pathogenic mutation for probable Alzheimer's disease in the APP gene at the N-terminus of beta-amyloid. *Nat Genet* 1: 345-7
- Muller FJ, Snyder EY, Loring JF. 2006. Gene therapy: can neural stem cells deliver? *Nat Rev Neurosci* 7: 75-84
- Muller U, Cristina N, Li ZW, Wolfer DP, Lipp HP, et al. 1994. Behavioral and anatomical deficits in mice homozygous for a modified beta-amyloid precursor protein gene. *Cell* 79: 755-65
- Muller UC, Zheng H. 2012. Physiological functions of APP family proteins. *Cold Spring Harb Perspect Med* 2: a006288
- Murayama Y, Takeda S, Yonezawa K, Giambarella U, Nishimoto I, Ogata E. 1996. Cell surface receptor function of amyloid precursor protein that activates Ser/Thr kinases. *Gerontology* 42 Suppl 1: 2-11
- Murray F, Smith DW, Hutson PH. 2008. Chronic low dose corticosterone exposure decreased hippocampal cell proliferation, volume and induced anxiety and depression like behaviours in mice. *Eur J Pharmacol* 583: 115-27
- Murrell J, Farlow M, Ghetti B, Benson MD. 1991. A mutation in the amyloid precursor protein associated with hereditary Alzheimer's disease. *Science* 254: 97-9
- Naiki H, Gejyo F, Nakakuki K. 1997. Concentration-dependent inhibitory effects of apolipoprotein E on Alzheimer's beta-amyloid fibril formation in vitro. *Biochemistry* 36: 6243-50
- Nairismagi J, Grohn OH, Kettunen MI, Nissinen J, Kauppinen RA, Pitkanen A. 2004. Progression of brain damage after status epilepticus and its association with epileptogenesis: a quantitative MRI study in a rat model of temporal lobe epilepsy. *Epilepsia* 45: 1024-34
- Nakagawa M, Koyanagi M, Tanabe K, Takahashi K, Ichisaka T, et al. 2008. Generation of induced pluripotent stem cells without Myc from mouse and human fibroblasts. *Nat Biotechnol* 26: 101-6
- Nakamura Y, Takeda M, Suzuki H, Hattori H, Tada K, et al. 1991. Abnormal distribution of cathepsins in the brain of patients with Alzheimer's disease. *Neurosci Lett* 130: 195-8

- Nakatomi H, Kuriu T, Okabe S, Yamamoto S, Hatano O, et al. 2002. Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors. *Cell* 110: 429-41
- Nalivaeva NN, Belyaev ND, Zhuravin IA, Turner AJ. 2012. The Alzheimer's amyloid-degrading peptidase, neprilysin: can we control it? *Int J Alzheimers Dis* 2012: 383796
- Namba Y, Tomonaga M, Kawasaki H, Otomo E, Ikeda K. 1991. Apolipoprotein E immunoreactivity in cerebral amyloid deposits and neurofibrillary tangles in Alzheimer's disease and kuru plaque amyloid in Creutzfeldt-Jakob disease. *Brain Res* 541: 163-6
- Naruse S, Thinakaran G, Luo JJ, Kusiak JW, Tomita T, et al. 1998. Effects of PS1 deficiency on membrane protein trafficking in neurons. *Neuron* 21: 1213-21
- Needham BE, Wlodek ME, Ciccotosto GD, Fam BC, Masters CL, et al. 2008. Identification of the Alzheimer's disease amyloid precursor protein (APP) and its homologue APLP2 as essential modulators of glucose and insulin homeostasis and growth. *J Pathol* 215: 155-63
- Neumann H, Kotter MR, Franklin RJ. 2009. Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain* 132: 288-95
- Neumann H, Takahashi K. 2007. Essential role of the microglial triggering receptor expressed on myeloid cells-2 (TREM2) for central nervous tissue immune homeostasis. *J Neuroimmunol* 184: 92-9
- Nicholson DW, Thornberry NA. 2003. Apoptosis. Life and death decisions. *Science* 299: 214-5
- Nicolas M, Hassan BA. 2014. Amyloid precursor protein and neural development. *Development* 141: 2543-8
- Nieto M, Schuurmans C, Britz O, Guillemot F. 2001. Neural bHLH genes control the neuronal versus glial fate decision in cortical progenitors. *Neuron* 29: 401-13
- Nieto-Estevez V, Pignatelli J, Arauzo-Bravo MJ, Hurtado-Chong A, Vicario-Abejon C. 2013. A global transcriptome analysis reveals molecular hallmarks of neural stem cell death, survival, and differentiation in response to partial FGF-2 and EGF deprivation. *PLoS One* 8: e53594
- Nilsberth C, Westlind-Danielsson A, Eckman CB, Condron MM, Axelman K, et al. 2001. The 'Arctic' APP mutation (E693G) causes Alzheimer's disease by enhanced Aβ protofibril formation. *Nature neuroscience* 4: 887-93
- Noctor SC, Martínez-Cerdeño V, Ivic L, Kriegstein AR. 2004. Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci* 7: 136-44
- Nomura T, Katunuma N. 2005. Involvement of cathepsins in the invasion, metastasis and proliferation of cancer cells. *J Med Invest* 52: 1-9
- Nuber S, Petrasch-Parwez E, Winner B, Winkler J, von Horsten S, et al. 2008. Neurodegeneration and motor dysfunction in a conditional model of Parkinson's disease. *J Neurosci* 28: 2471-84
- Nunan J, Small DH. 2000. Regulation of APP cleavage by alpha-, beta- and gamma-secretases. *FEBS Lett* 483: 6-10
- O'Keefe GC, Tyers P, Aarsland D, Dalley JW, Barker RA, Caldwell MA. 2009. Dopamine-induced proliferation of adult neural precursor cells in the

- mammalian subventricular zone is mediated through EGF. *Proc Natl Acad Sci U S A* 106: 8754-9
- O'Leary CJ, Bradford D, Chen M, White A, Blackmore DG, Cooper HM. 2014. The Netrin/RGM Receptor, Neogenin, Controls Adult Neurogenesis by Promoting Neuroblast Migration and Cell Cycle Exit. *stem cells*
- Octave JN, Essalmani R, Tasiaux B, Menager J, Czech C, Mercken L. 2000. The role of presenilin-1 in the gamma-secretase cleavage of the amyloid precursor protein of Alzheimer's disease. *J Biol Chem* 275: 1525-8
- Oddo S, Caccamo A, Green KN, Liang K, Tran L, et al. 2005. Chronic nicotine administration exacerbates tau pathology in a transgenic model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 102: 3046-51
- Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, et al. 2003. Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron* 39: 409-21
- Ohkawara T, Nagase H, Koh CS, Nakayama K. 2011. The amyloid precursor protein intracellular domain alters gene expression and induces neuron-specific apoptosis. *Gene* 475: 1-9
- Ohno M, Chang L, Tseng W, Oakley H, Citron M, et al. 2006. Temporal memory deficits in Alzheimer's mouse models: rescue by genetic deletion of BACE1. *Eur J Neurosci* 23: 251-60
- Ohsawa I, Takamura C, Morimoto T, Ishiguro M, Kohsaka S. 1999. Amino-terminal region of secreted form of amyloid precursor protein stimulates proliferation of neural stem cells. *Eur J Neurosci* 11: 1907-13
- Okamoto T, Takeda S, Murayama Y, Ogata E, Nishimoto I. 1995. Ligand-dependent G protein coupling function of amyloid transmembrane precursor. *J Biol Chem* 270: 4205-8
- Olcese JM, Cao C, Mori T, Mamcarz MB, Maxwell A, et al. 2009. Protection against cognitive deficits and markers of neurodegeneration by long-term oral administration of melatonin in a transgenic model of Alzheimer disease. *J Pineal Res* 47: 82-96
- Olgiati P, Politis AM, Papadimitriou GN, De Ronchi D, Serretti A. 2011. Genetics of late-onset Alzheimer's disease: update from the alzgene database and analysis of shared pathways. *Int J Alzheimers Dis* 2011: 832379
- Oltersdorf T, Ward PJ, Henriksson T, Beattie EC, Neve R, et al. 1990. The Alzheimer amyloid precursor protein. Identification of a stable intermediate in the biosynthetic/degradative pathway. *J Biol Chem* 265: 4492-7
- Orgogozo JM, Gilman S, Dartigues JF, Laurent B, Puel M, et al. 2003. Subacute meningoencephalitis in a subset of patients with AD after Abeta42 immunization. *Neurology* 61: 46-54
- Ortega JA, Alcantara S. 2010. BDNF/MAPK/ERK-induced BMP7 expression in the developing cerebral cortex induces premature radial glia differentiation and impairs neuronal migration. *Cereb Cortex* 20: 2132-44
- Otto HH, Schirmeister T. 1997. Cysteine Proteases and Their Inhibitors. *Chem Rev* 97: 133-72
- Otto T, Eichenbaum H, Wiener SI, Wible CG. 1991. Learning-related patterns of CA1 spike trains parallel stimulation parameters optimal for inducing hippocampal long-term potentiation. *Hippocampus* 1: 181-92

- Oya S, Yoshikawa G, Takai K, Tanaka J, Higashiyama S, et al. 2008. Region-specific proliferative response of neural progenitors to exogenous stimulation by growth factors following ischemia. *Neuroreport* 19: 805-9
- Ozaki T, Li Y, Kikuchi H, Tomita T, Iwatsubo T, Nakagawara A. 2006. The intracellular domain of the amyloid precursor protein (AICD) enhances the p53-mediated apoptosis. *Biochem Biophys Res Commun* 351: 57-63
- Ozen I, Galichet C, Watts C, Parras C, Guillemot F, Raineteau O. 2007. Proliferating neuronal progenitors in the postnatal hippocampus transiently express the proneural gene Ngn2. *Eur J Neurosci* 25: 2591-603
- Paez-Gonzalez P, Asrican B, Rodriguez E, Kuo CT. 2014. Identification of distinct ChAT(+) neurons and activity-dependent control of postnatal SVZ neurogenesis. *Nature neuroscience* 17: 934-42
- Pallotto M, Nissant A, Fritschy JM, Rudolph U, Sassoe-Pognetto M, et al. 2012. Early formation of GABAergic synapses governs the development of adult-born neurons in the olfactory bulb. *J Neurosci* 32: 9103-15
- Palm DE, Knuckey NW, Primiano MJ, Spangenberg AG, Johanson CE. 1995. Cystatin C, a protease inhibitor, in degenerating rat hippocampal neurons following transient forebrain ischemia. *Brain Res* 691: 1-8
- Palmer TD, Markakis EA, Willhoite AR, Safar F, Gage FH. 1999. Fibroblast growth factor-2 activates a latent neurogenic program in neural stem cells from diverse regions of the adult CNS. *J Neurosci* 19: 8487-97
- Palmer TD, Takahashi J, Gage FH. 1997. The adult rat hippocampus contains primordial neural stem cells. *Mol Cell Neurosci* 8: 389-404
- Palmer TD, Willhoite AR, Gage FH. 2000. Vascular niche for adult hippocampal neurogenesis. *J Comp Neurol* 425: 479-94
- Palmert MR, Golde TE, Cohen ML, Kovacs DM, Tanzi RE, et al. 1988. Amyloid protein precursor messenger RNAs: differential expression in Alzheimer's disease. *Science* 241: 1080-4
- Paloneva J, Manninen T, Christman G, Hovanes K, Mandelin J, et al. 2002. Mutations in two genes encoding different subunits of a receptor signaling complex result in an identical disease phenotype. *Am J Hum Genet* 71: 656-62
- Parameshwaran K, Dhanasekaran M, Suppiramaniam V. 2008. Amyloid beta peptides and glutamatergic synaptic dysregulation. *Exp Neurol* 210: 7-13
- Pardossi-Piquard R, Petit A, Kawarai T, Sunyach C, Alves da Costa C, et al. 2005. Presenilin-dependent transcriptional control of the Abeta-degrading enzyme neprilysin by intracellular domains of betaAPP and APLP. *Neuron* 46: 541-54
- Parent JM, Yu TW, Leibowitz RT, Geschwind DH, Sloviter RS, Lowenstein DH. 1997. Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J Neurosci* 17: 3727-38
- Park CH, Kang JS, Yoon EH, Shim JW, Suh-Kim H, Lee SH. 2008a. Proneural bHLH neurogenin 2 differentially regulates Nurr1-induced dopamine neuron differentiation in rat and mouse neural precursor cells in vitro. *FEBS Lett* 582: 537-42
- Park DH, Eve DJ. 2009. Regenerative medicine: advances in new methods and technologies. *Med Sci Monit* 15: RA233-51

- Park IH, Arora N, Huo H, Maherali N, Ahfeldt T, et al. 2008b. Disease-specific induced pluripotent stem cells. *Cell* 134: 877-86
- Park IH, Lerou PH, Zhao R, Huo H, Daley GQ. 2008c. Generation of human-induced pluripotent stem cells. *Nat Protoc* 3: 1180-6
- Parras CM, Schuurmans C, Scardigli R, Kim J, Anderson DJ, Guillemot F. 2002. Divergent functions of the proneural genes Mash1 and Ngn2 in the specification of neuronal subtype identity. *Genes Dev* 16: 324-38
- Parrish-Aungst S, Shipley MT, Erdelyi F, Szabo G, Puche AC. 2007. Quantitative analysis of neuronal diversity in the mouse olfactory bulb. *J Comp Neurol* 501: 825-36
- Parsons CG, Danysz W, Quack G. 1998. Glutamate in CNS disorders as a target for drug development: an update. *Drug News Perspect* 11: 523-69
- Parvathy S, Hussain I, Karran EH, Turner AJ, Hooper NM. 1999. Cleavage of Alzheimer's amyloid precursor protein by alpha-secretase occurs at the surface of neuronal cells. *Biochemistry* 38: 9728-34
- Pasinetti GM, Ungar LH, Lange DJ, Yemul S, Deng H, et al. 2006. Identification of potential CSF biomarkers in ALS. *Neurology* 66: 1218-22
- Pasternak SH, Callahan JW, Mahuran DJ. 2004. The role of the endosomal/lysosomal system in amyloid-beta production and the pathophysiology of Alzheimer's disease: reexamining the spatial paradox from a lysosomal perspective. *Journal of Alzheimer's disease : JAD* 6: 53-65
- Pastorino L, Sun A, Lu PJ, Zhou XZ, Balastik M, et al. 2006. The prolyl isomerase Pin1 regulates amyloid precursor protein processing and amyloid-beta production. *Nature* 440: 528-34
- Patterson D, Gardiner K, Kao FT, Tanzi R, Watkins P, Gusella JF. 1988. Mapping of the gene encoding the beta-amyloid precursor protein and its relationship to the Down syndrome region of chromosome 21. *Proc Natl Acad Sci U S A* 85: 8266-70
- Perera TD, Coplan JD, Lisanby SH, Lipira CM, Arif M, et al. 2007. Antidepressant-induced neurogenesis in the hippocampus of adult nonhuman primates. *J Neurosci* 27: 4894-901
- Perez RG, Soriano S, Hayes JD, Ostaszewski B, Xia W, et al. 1999. Mutagenesis identifies new signals for beta-amyloid precursor protein endocytosis, turnover, and the generation of secreted fragments, including Abeta42. *J Biol Chem* 274: 18851-6
- Perry EK, Johnson M, Ekonomou A, Perry RH, Ballard C, Attems J. 2012. Neurogenic abnormalities in Alzheimer's disease differ between stages of neurogenesis and are partly related to cholinergic pathology. *Neurobiol Dis.* 47: 155-62
- Peterziel H, Sackmann T, Strelau J, Kuhn PH, Lichtenthaler SF, et al. 2011. F-spondin regulates neuronal survival through activation of disabled-1 in the chicken ciliary ganglion. *Mol Cell Neurosci* 46: 483-97
- Petreanu L, Alvarez-Buylla A. 2002. Maturation and death of adult-born olfactory bulb granule neurons: role of olfaction. *J Neurosci* 22: 6106-13
- Pezzini A, Del Zotto E, Volonghi I, Giossi A, Costa P, Padovani A. 2009. Cerebral amyloid angiopathy: a common cause of cerebral hemorrhage. *Curr Med Chem* 16: 2498-513

- Pfeifer LA, White LR, Ross GW, Petrovitch H, Launer LJ. 2002. Cerebral amyloid angiopathy and cognitive function: the HAAS autopsy study. *Neurology* 58: 1629-34
- Pfriegeer FW. 2003a. Cholesterol homeostasis and function in neurons of the central nervous system. *Cell Mol Life Sci* 60: 1158-71
- Pfriegeer FW. 2003b. Role of cholesterol in synapse formation and function. *Biochim Biophys Acta* 1610: 271-80
- Picard-Riera N, Decker L, Delarasse C, Goude K, Nait-Oumesmar B, et al. 2002. Experimental autoimmune encephalomyelitis mobilizes neural progenitors from the subventricular zone to undergo oligodendrogenesis in adult mice. *Proc Natl Acad Sci U S A* 99: 13211-6
- Pike CJ, Cummings BJ, Cotman CW. 1995. Early association of reactive astrocytes with senile plaques in Alzheimer's disease. *Exp Neurol* 132: 172-9
- Pirttila TJ, Lukasiuk K, Hakansson K, Grubb A, Abrahamson M, Pitkanen A. 2005. Cystatin C modulates neurodegeneration and neurogenesis following status epilepticus in mouse. *Neurobiol Dis* 20: 241-53
- Pitas RE, Boyles JK, Lee SH, Foss D, Mahley RW. 1987. Astrocytes synthesize apolipoprotein E and metabolize apolipoprotein E-containing lipoproteins. *Biochim Biophys Acta* 917: 148-61
- Platel JC, Dave KA, Gordon V, Lacar B, Rubio ME, Bordey A. 2010. NMDA receptors activated by subventricular zone astrocytic glutamate are critical for neuroblast survival prior to entering a synaptic network. *Neuron* 65: 859-72
- Platel JC, Gordon V, Heintz T, Bordey A. 2009. GFAP-GFP neural progenitors are antigenically homogeneous and anchored in their enclosed mosaic niche. *Glia* 57: 66-78
- Pomeroy SL, LaMantia AS, Purves D. 1990. Postnatal construction of neural circuitry in the mouse olfactory bulb. *J Neurosci* 10: 1952-66
- Postina R, Schroeder A, Dewachter I, Bohl J, Schmitt U, et al. 2004. A disintegrin-metalloproteinase prevents amyloid plaque formation and hippocampal defects in an Alzheimer disease mouse model. *The Journal of clinical investigation* 113: 1456-64
- Potten CS, Loeffler M. 1990. Stem cells: attributes, cycles, spirals, pitfalls and uncertainties. Lessons for and from the crypt. *Development* 110: 1001-20
- Pottier C, Wallon D, Rousseau S, Rovelet-Lecrux A, Richard AC, et al. 2013. TREM2 R47H variant as a risk factor for early-onset Alzheimer's disease. *J Alzheimers Dis* 35: 45-9
- Powers CJ, McLeskey SW, Wellstein A. 2000. Fibroblast growth factors, their receptors and signaling. *Endocr Relat Cancer* 7: 165-97
- Pressler RT, Inoue T, Strowbridge BW. 2007. Muscarinic receptor activation modulates granule cell excitability and potentiates inhibition onto mitral cells in the rat olfactory bulb. *J Neurosci* 27: 10969-81
- Price SA, Held B, Pearson HA. 1998. Amyloid beta protein increases Ca²⁺ currents in rat cerebellar granule neurones. *Neuroreport* 9: 539-45
- Priller C, Bauer T, Mitteregger G, Krebs B, Kretzschmar HA, Herms J. 2006. Synapse formation and function is modulated by the amyloid precursor protein. *J Neurosci* 26: 7212-21

- Priester A, Saraulli D, Farioli-Vecchioli S, Strimpakos G, Costanzi M, et al. 2013. Impact of N-tau on adult hippocampal neurogenesis, anxiety, and memory. *Neurobiol Aging* 34: 2551-63
- Prvulovic D, Hampel H, Pantel J. 2010. Galantamine for Alzheimer's disease. *Expert Opin Drug Metab Toxicol* 6: 345-54
- Puente XS, Sanchez LM, Overall CM, Lopez-Otin C. 2003. Human and mouse proteases: a comparative genomic approach. *Nat Rev Genet* 4: 544-58
- Puig KL, Swigost AJ, Zhou X, Sens MA, Combs CK. 2012. Amyloid precursor protein expression modulates intestine immune phenotype. *J Neuroimmune Pharmacol* 7: 215-30
- Pungercar JR, Caglic D, Sajid M, Dolinar M, Vasiljeva O, et al. 2009. Autocatalytic processing of procathepsin B is triggered by proenzyme activity. *Febs J* 276: 660-8
- Puzzo D, Privitera L, Fa' M, Staniszewski A, Hashimoto G, et al. 2011. Endogenous amyloid- β is necessary for hippocampal synaptic plasticity and memory. *Ann Neurol* 69: 819-30
- Puzzo D, Privitera L, Leznik E, Fa M, Staniszewski A, et al. 2008. Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. *J Neurosci* 28: 14537-45
- Qi-Takahara Y, Morishima-Kawashima M, Tanimura Y, Dolios G, Hirotsu N, et al. 2005. Longer forms of amyloid beta protein: implications for the mechanism of intramembrane cleavage by gamma-secretase. *J Neurosci* 25: 436-45
- Qiu WQ, Ferreira A, Miller C, Koo EH, Selkoe DJ. 1995. Cell-surface beta-amyloid precursor protein stimulates neurite outgrowth of hippocampal neurons in an isoform-dependent manner. *J Neurosci* 15: 2157-67
- Radde R, Bolmont T, Kaeser SA, Coomaraswamy J, Lindau D, et al. 2006. Abeta42-driven cerebral amyloidosis in transgenic mice reveals early and robust pathology. *EMBO Rep* 7: 940-6
- Radde R, Duma C, Goedert M, Jucker M. 2008. The value of incomplete mouse models of Alzheimer's disease. *Eur J Nucl Med Mol Imaging* 35 Suppl 1: S70-4
- Rafalski VA, Brunet A. 2011. Energy metabolism in adult neural stem cell fate. *Prog Neurobiol* 93: 182-203
- Rahmani A, Kheradmand D, Keyhanvar P, Shoaee-Hassani A, Darbandi-Azar A. 2013. Neurogenesis and increase in differentiated neural cell survival via phosphorylation of Akt1 after fluoxetine treatment of stem cells. *Biomed Res Int* 2013: 582526
- Rama N, Goldschneider D, Corset V, Lambert J, Pays L, Mehlen P. 2012. Amyloid precursor protein regulates netrin-1-mediated commissural axon outgrowth. *J Biol Chem* 287: 30014-23
- Ranganathan R, Sawin ER, Trent C, Horvitz HR. 2001. Mutations in the *Caenorhabditis elegans* serotonin reuptake transporter MOD-5 reveal serotonin-dependent and -independent activities of fluoxetine. *J Neurosci* 21: 5871-84
- Ray J, Peterson DA, Schinstine M, Gage FH. 1993. Proliferation, differentiation, and long-term culture of primary hippocampal neurons. *Proc Natl Acad Sci U S A* 90: 3602-6

- Rayaprolu S, Mullen B, Baker M, Lynch T, Finger E, et al. 2013. TREM2 in neurodegeneration: evidence for association of the p.R47H variant with frontotemporal dementia and Parkinson's disease. *Mol Neurodegener* 8: 19
- Reaume AG, Howland DS, Trusko SP, Savage MJ, Lang DM, et al. 1996. Enhanced amyloidogenic processing of the beta-amyloid precursor protein in gene-targeted mice bearing the Swedish familial Alzheimer's disease mutations and a "humanized" A β sequence. *J Biol Chem* 271: 23380-8
- Rebeck GW, Reiter JS, Strickland DK, Hyman BT. 1993. Apolipoprotein E in sporadic Alzheimer's disease: allelic variation and receptor interactions. *Neuron* 11: 575-80
- Rebok GW, Rovner BW, Folstein MF. 1991. Sleep disturbance and Alzheimer's disease: relationship to behavioral problems. *Aging (Milano)* 2: 193-6
- Reif A, Fritzen S, Finger M, Strobel A, Lauer M, et al. 2006. Neural stem cell proliferation is decreased in schizophrenia, but not in depression. *Mol Psychiatry* 11: 514-22
- Reinhard C, Hébert SS, De Strooper B. 2005. The amyloid- β precursor protein: integrating structure with biological function. *Embo J* 24: 3996-4006
- Reinhardt RR, Bondy CA. 1994. Insulin-like growth factors cross the blood-brain barrier. *Endocrinology* 135: 1753-61
- Reisberg B, Doody R, Stöffler A, Schmitt F, Ferris S, Möbius HJ. 2003. Memantine in moderate-to-severe Alzheimer's disease. *New Engl. J. Med.* 348: 1333-41
- Reitz C, Brayne C, Mayeux R. 2011. Epidemiology of Alzheimer disease. *Nat Rev Neurol* 7: 137-52
- Reuss B, Dermietzel R, Unsicker K. 1998. Fibroblast growth factor 2 (FGF-2) differentially regulates connexin (cx) 43 expression and function in astroglial cells from distinct brain regions. *Glia* 22: 19-30
- Revesz T, Ghiso J, Lashley TP, G. Rostagno, A., Frangione B, Holton JL. 2003. Cerebral amyloid angiopathies: a pathologic, biochemical, and genetic view. *J Neuropathol Exp Neurol* 9: 885-98
- Reynolds BA, Rietze RL. 2005. Neural stem cells and neurospheres--re-evaluating the relationship. *Nat Methods* 2: 333-6
- Reynolds BA, Weiss S. 1992. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255: 1707-10
- Reynolds BA, Weiss S. 1996. Clonal and population analyses demonstrate that an EGF-responsive mammalian embryonic CNS precursor is a stem cell. *Dev Biol* 175: 1-13
- Richards LJ, Kilpatrick TJ, Bartlett PF. 1992. De novo generation of neuronal cells from the adult mouse brain. *Proc Natl Acad Sci U S A* 89: 8591-5
- Rietze RL, Valcanis H, Brooker GF, Thomas T, Voss AK, Bartlett PF. 2001. Purification of a pluripotent neural stem cell from the adult mouse brain. *Nature* 412: 736-9
- Ring S, Weyer SW, Kilian SB, Waldron E, Pietrzik CU, et al. 2007. The secreted beta-amyloid precursor protein ectodomain APPs α is sufficient to rescue the anatomical, behavioral, and electrophysiological abnormalities of APP-deficient mice. *J Neurosci* 27: 7817-26

- Ringheim GE, Szczepanik AM, Petko W, Burgher KL, Zhu SZ, Chao CC. 1998. Enhancement of beta-amyloid precursor protein transcription and expression by the soluble interleukin-6 receptor/interleukin-6 complex. *Brain Res Mol Brain Res* 55: 35-44
- Riordan KC, Hoffman Snyder CR, Wellik KE, Caselli RJ, Wingerchuk DM, Demaerschalk BM. 2011. Effectiveness of adding memantine to an Alzheimer dementia treatment regimen which already includes stable donepezil therapy: a critically appraised topic. *Neurologist* 17: 121-3
- Rippon HJ, Bishop AE. 2004. Embryonic stem cells. *Cell Prolif* 37: 23-34
- Roberds SL, Anderson J, Basi G, Bienkowski MJ, Branstetter DG, et al. 2001. BACE knockout mice are healthy despite lacking the primary beta-secretase activity in brain: implications for Alzheimer's disease therapeutics. *Hum Mol Genet* 10: 1317-24
- Rochefort C, Gheusi G, Vincent JD, Lledo PM. 2002. Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory. *J Neurosci* 22: 2679-89
- Rodgers SP, Born HA, Das P, Jankowsky JL. 2012. Transgenic APP expression during postnatal development causes persistent locomotor hyperactivity in the adult. *Mol Neurodegener* 7: 28
- Rodriguez JJ, Jones VC, Tabuchi M, Allan SM, Knight EM, et al. 2008. Impaired adult neurogenesis in the dentate gyrus of a triple transgenic mouse model of Alzheimer's disease. *PLoS One* 3: e2935
- Rogelius N, Hebsgaard JB, Lundberg C, Parmar M. 2008. Reprogramming of neonatal SVZ progenitors by islet-1 and neurogenin-2. *Mol Cell Neurosci* 38: 453-9
- Roher AE, Lowenson JD, Clarke S, Woods AS, Cotter RJ, et al. 1993. beta-Amyloid-(1-42) is a major component of cerebrovascular amyloid deposits: implications for the pathology of Alzheimer disease. *Proc Natl Acad Sci U S A* 90: 10836-40
- Ross SE, Greenberg ME, Stiles CD. 2003. Basic helix-loop-helix factors in cortical development. *Neuron* 39: 13-25
- Rossjohn J, Cappai R, Feil SC, Henry A, McKinsty WJ, et al. 1999. Crystal structure of the N-terminal, growth factor-like domain of Alzheimer amyloid precursor protein. *Nat Struct Biol* 6: 327-31
- Rousselot P, Lois C, Alvarez-Buylla A. 1995. Embryonic (PSA) N-CAM reveals chains of migrating neuroblasts between the lateral ventricle and the olfactory bulb of adult mice. *J Comp Neurol* 351: 51-61
- Rousselot P, Nottebohm F. 1995. Expression of polysialylated N-CAM in the central nervous system of adult canaries and its possible relation to function. *J Comp Neurol* 356: 629-40
- Roy NS, Cleren C, Singh SK, Yang L, Beal MF, Goldman SA. 2006. Functional engraftment of human ES cell-derived dopaminergic neurons enriched by coculture with telomerase-immortalized midbrain astrocytes. *Nature medicine* 12: 1259-68
- Roybon L, Deierborg T, Brundin P, Li JY. 2009a. Involvement of Ngn2, Tbr and NeuroD proteins during postnatal olfactory bulb neurogenesis. *Eur J Neurosci* 29: 232-43
- Roybon L, Hjalt T, Stott S, Guillemot F, Li JY, Brundin P. 2009b. Neurogenin2 directs granule neuroblast production and amplification while NeuroD1

- specifies neuronal fate during hippocampal neurogenesis. *PLoS One* 4: e4779
- Roybon L, Mastracci TL, Ribeiro D, Sussel L, Brundin P, Li JY. 2010. GABAergic differentiation induced by Mash1 is compromised by the bHLH proteins Neurogenin2, NeuroD1, and NeuroD2. *Cereb Cortex* 20: 1234-44
- Ruiz i Altaba A, Palma V, Dahmane N. 2002. Hedgehog-Gli signalling and the growth of the brain. *Nat Rev Neurosci* 3: 24-33
- Ryan KA, Pimplikar SW. 2005. Activation of GSK-3 and phosphorylation of CRMP2 in transgenic mice expressing APP intracellular domain. *J Cell Biol* 171: 327-35
- Ryu JK, Cho T, Wang YT, McLarnon JG. 2009. Neural progenitor cells attenuate inflammatory reactivity and neuronal loss in an animal model of inflamed AD brain. *J Neuroinflammation* 6: 39
- Rzychon M, Chmiel D, Stec-Niemczyk J. 2004. Modes of inhibition of cysteine proteases. *Acta Biochim Pol* 51: 861-73
- Sabo SL, Ikin AF, Buxbaum JD, Greengard P. 2003. The amyloid precursor protein and its regulatory protein, FE65, in growth cones and synapses in vitro and in vivo. *J Neurosci* 23: 5407-15
- Sabo SL, Lanier LM, Ikin AF, Khorkova O, Sahasrabudhe S, et al. 1999. Regulation of beta-amyloid secretion by FE65, an amyloid protein precursor-binding protein. *J Biol Chem* 274: 7952-7
- Sahin U, Weskamp G, Kelly K, Zhou HM, Higashiyama S, et al. 2004. Distinct roles for ADAM10 and ADAM17 in ectodomain shedding of six EGFR ligands. *J Cell Biol* 164: 769-79
- Sahlin C, Lord A, Magnusson K, Englund H, Almeida CG, et al. 2007. The Arctic Alzheimer mutation favors intracellular amyloid-beta production by making amyloid precursor protein less available to alpha-secretase. *Journal of neurochemistry* 101: 854-62
- Saino-Saito S, Cave JW, Akiba Y, Sasaki H, Goto K, et al. 2007. ER81 and CaMKIV identify anatomically and phenotypically defined subsets of mouse olfactory bulb interneurons. *J Comp Neurol* 502: 485-96
- Saitoh T, Sundsmo M, Roch JM, Kimura N, Cole G, et al. 1989. Secreted form of amyloid beta protein precursor is involved in the growth regulation of fibroblasts. *Cell* 58: 615-22
- Sakamoto M, Ieki N, Miyoshi G, Mochimaru D, Miyachi H, et al. 2014. Continuous postnatal neurogenesis contributes to formation of the olfactory bulb neural circuits and flexible olfactory associative learning. *J Neurosci* 34: 5788-99
- Sala Frigerio C, Fadeeva JV, Minogue AM, Citron M, Van Leuven F, et al. 2010. β -secretase cleavage is not required for generation of the intracellular C-terminal domain of the amyloid precursor family of proteins. *FEBS J.* 277: 1503-18
- Salbaum JM, Ruddle FH. 1994. Embryonic expression pattern of amyloid protein precursor suggests a role in differentiation of specific subsets of neurons. *J Exp Zool* 269: 116-27
- Salewski RP, Eftekharpour E, Fehlings MG. 2010. Are induced pluripotent stem cells the future of cell-based regenerative therapies for spinal cord injury? *J Cell Physiol* 222: 515-21

- Salloway S, Sperling R, Fox NC, Blennow K, Klunk W, et al. 2014. Two phase 3 trials of bapineuzumab in mild-to-moderate Alzheimer's disease. *N Engl J Med* 370: 322-33
- Salmena L, Lemmers B, Hakem A, Matysiak-Zablocki E, Murakami K, et al. 2003. Essential role for caspase 8 in T-cell homeostasis and T-cell-mediated immunity. *Genes Dev* 17: 883-95
- Salvioli R, Ricci-Vitiani L, Tatti M, Scarpa S, De Maria R, Vaccaro AM. 2008. The secretion and maturation of prosaposin and procathepsin D are blocked in embryonic neural progenitor cells. *Biochim Biophys Acta* 1783: 1480-9
- Sanai N, Nguyen T, Ihrie RA, Mirzadeh Z, Tsai HH, et al. 2011. Corridors of migrating neurons in the human brain and their decline during infancy. *Nature* 478: 382-6
- Sanan DA, Weisgraber KH, Russell SJ, Mahley RW, Huang DS, A., et al. 1994. Apolipoprotein E associates with beta amyloid peptide of Alzheimer's disease to form novel monofibrils. Isoform apoE4 associates more efficiently than apoE3. *J Clin Invest* 94
- Sanchez-Mejia RO, Newman JW, Toh S, Yu GQ, Zhou Y, et al. 2008. Phospholipase A2 reduction ameliorates cognitive deficits in a mouse model of Alzheimer's disease. *Nat Neurosci* 11: 1311-8
- Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, et al. 2003. Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 301: 805-9
- Santos DM, Xavier JM, Morgado AL, Sola S, Rodrigues CM. 2012. Distinct regulatory functions of calpain 1 and 2 during neural stem cell self-renewal and differentiation. *PLoS One* 7: e33468
- Sarkar A, Hochedlinger K. 2013. The sox family of transcription factors: versatile regulators of stem and progenitor cell fate. *Cell Stem Cell* 12: 15-30
- Sarter M, Bruno JP. 2004. Developmental origins of the age-related decline in cortical cholinergic function and associated cognitive abilities. *Neurobiol Aging* 25: 1127-39
- Sastre M, Steiner H, Fuchs K, Capell A, Multhaup G, et al. 2001. Presenilin-dependent gamma-secretase processing of beta-amyloid precursor protein at a site corresponding to the S3 cleavage of Notch. *EMBO Rep* 2: 835-41
- Satija NK, Singh VK, Verma YK, Gupta P, Sharma S, et al. 2009. Mesenchymal stem cell-based therapy: a new paradigm in regenerative medicine. *J Cell Mol Med* 13: 4385-402
- Saura CA, Choi SY, Beglopoulos V, Malkani S, Zhang D, et al. 2004. Loss of presenilin function causes impairments of memory and synaptic plasticity followed by age-dependent neurodegeneration. *Neuron* 42: 23-36
- Saxe MD, Battaglia F, Wang JW, Malleret G, David DJ, et al. 2006. Ablation of hippocampal neurogenesis impairs contextual fear conditioning and synaptic plasticity in the dentate gyrus. *Proc Natl Acad Sci U S A* 103: 17501-6
- Sberna G, Saez-Valero J, Beyreuther K, Masters CL, Small DH. 1997. The amyloid beta-protein of Alzheimer's disease increases acetylcholinesterase expression by increasing intracellular calcium in embryonal carcinoma P19 cells. *J Neurochem* 69: 1177-84

- Scandura JM, Zhang Y, Van Nostrand WE, Walsh PN. 1997. Progress curve analysis of the kinetics with which blood coagulation factor XIa is inhibited by protease nexin-2. *Biochemistry* 36: 412-20
- Scardigli R, Capelli P, Vignone D, Brandi R, Ceci M, et al. 2014. Neutralization of nerve growth factor impairs proliferation and differentiation of adult neural progenitors in the subventricular zone. *stem cells* 32: 2516-28
- Scardigli R, Schuurmans C, Gradwohl G, Guillemot F. 2001. Crossregulation between Neurogenin2 and pathways specifying neuronal identity in the spinal cord. *Neuron* 31: 203-17
- Schabitz WR, Steigleder T, Cooper-Kuhn CM, Schwab S, Sommer C, et al. 2007. Intravenous brain-derived neurotrophic factor enhances poststroke sensorimotor recovery and stimulates neurogenesis. *Stroke* 38: 2165-72
- Scheff SW, Price DA, Schmitt FA, Mufson EJ. 2006. Hippocampal synaptic loss in early Alzheimer's disease and mild cognitive impairment. *Neurobiol Aging* 27: 1372-84
- Scheinfeld MH, Gherzi E, Laky K, Fowlkes BJ, D'Adamio L. 2002a. Processing of beta-amyloid precursor-like protein-1 and -2 by gamma-secretase regulates transcription. *J Biol Chem* 277: 44195-201
- Scheinfeld MH, Roncarati R, Vito P, Lopez PA, Abdallah M, D'Adamio L. 2002b. Jun NH2-terminal kinase (JNK) interacting protein 1 (JIP1) binds the cytoplasmic domain of the Alzheimer's beta-amyloid precursor protein (APP). *J Biol Chem* 277: 3767-75
- Schenk D, Barbour R, Dunn W, Gordon G, Grajeda H, et al. 1999. Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse. *Nature* 400: 173-7
- Scheuner D, Eckman C, Jensen M, Song X, Citron M, et al. 1996. Secreted amyloid β -protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat Med* 2: 864-70
- Schindowski K, Belarbi K, Bretteville A, Ando K, Buee L. 2008. Neurogenesis and cell cycle-reactivated neuronal death during pathogenic tau aggregation. *Genes Brain Behav* 7 Suppl 1: 92-100
- Schmechel DE, Saunders AM, Strittmatter WJ, Crain BJ, Hulette CM, et al. 1993. Increased amyloid beta-peptide deposition in cerebral cortex as a consequence of apolipoprotein E genotype in late-onset Alzheimer disease. *Proc Natl Acad Sci U S A* 90: 9649-53
- Schmidt-Hieber C, Jonas P, Bischofberger J. 2004. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature* 429: 184-7
- Schmidt-Salzmann C, Li L, Bischofberger J. 2014. Functional properties of extrasynaptic AMPA and NMDA receptors during postnatal hippocampal neurogenesis. *J Physiol* 592: 125-40
- Schneider LS, Dagerman KS, Higgins JP, McShane R. 2011. Lack of evidence for the efficacy of memantine in mild Alzheimer disease. *Arch Neurol* 68: 991-8
- Schneider MR, Werner S, Paus R, Wolf E. 2008. Beyond wavy hairs: the epidermal growth factor receptor and its ligands in skin biology and pathology. *Am J Pathol* 173: 14-24

- Schwartz MW, Bergman RN, Kahn SE, Taborsky GJ, Jr., Fisher LD, et al. 1991. Evidence for entry of plasma insulin into cerebrospinal fluid through an intermediate compartment in dogs. Quantitative aspects and implications for transport. *The Journal of clinical investigation* 88: 1272-81
- Scotto-Lomassese S, Strambi C, Strambi A, Aouane A, Augier R, et al. 2003. Suppression of adult neurogenesis impairs olfactory learning and memory in an adult insect. *J Neurosci* 23: 9289-96
- Seaberg RM, van der Kooy D. 2003. Stem and progenitor cells: the premature desertion of rigorous definitions. *Trends Neurosci* 26: 125-31
- Seabrook GR, Smith DW, Bowery BJ, Easter A, Reynolds T, et al. 1999. Mechanisms contributing to the deficits in hippocampal synaptic plasticity in mice lacking amyloid precursor protein. *Neuropharmacology* 38: 349-59
- Selkoe D, Kopan R. 2003. Notch and Presenilin: regulated intramembrane proteolysis links development and degeneration. *Annu Rev Neurosci* 26: 565-97
- Selkoe DJ. 1993. Physiological production of the beta-amyloid protein and the mechanism of Alzheimer's disease. *Trends Neurosci* 16: 403-9
- Selkoe DJ. 2001. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 81: 741-66
- Selkoe DJ, Abraham CR, Podlisny MB, Duffy LK. 1986. Isolation of low-molecular-weight proteins from amyloid plaque fibers in Alzheimer's disease. *Journal of neurochemistry* 46: 1820-34
- Seo JS, Leem YH, Lee KW, Kim SW, Lee JK, Han PL. 2010. Severe motor neuron degeneration in the spinal cord of the Tg2576 mouse model of Alzheimer disease. *Journal of Alzheimer's disease : JAD* 21: 263-76
- Seo Y, Yang SR, Jee MK, Joo EK, Roh KH, et al. 2011. Human umbilical cord blood-derived mesenchymal stem cells protect against neuronal cell death and ameliorate motor deficits in Niemann Pick type C1 mice. *Cell Transplant* 20: 1033-47
- Seri B, Alvarez-Buylla A. 2002. Neural stem cell and the regulation of neurogenesis in adult hippocampus. *Clin Neurosci Res* 2: 11-16
- Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Buylla A. 2001. Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 21: 7153-60
- Seri B, Herrera DG, Gritti A, Ferron S, Collado L, et al. 2006. Composition and organization of the SCZ: a large germinal layer containing neural stem cells in the adult mammalian brain. *Cereb Cortex* 16 Suppl 1: i103-11
- Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT. 2011. Neuropathological alterations in Alzheimer disease. *Cold Spring Harb Perspect Med* 1: a006189
- Serrano-Pozo A, William CM, Ferrer I, Uro-Coste E, Delisle MB, et al. 2010. Beneficial effect of human anti-amyloid-beta active immunization on neurite morphology and tau pathology. *Brain* 133: 1312-27
- Sessa G, Podini P, Mariani M, Meroni A, Spreafico R, et al. 2004. Distribution and signaling of TREM2/DAP12, the receptor system mutated in human polycystic lipomembraneous osteodysplasia with sclerosing leukoencephalopathy dementia. *Eur J Neurosci* 20: 2617-28

- Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, et al. 1992. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* 359: 325-7
- Shamblott MJ, Axelman J, Wang S, Bugg EM, Littlefield JW, et al. 1998. Derivation of pluripotent stem cells from cultured human primordial germ cells. *Proc Natl Acad Sci U S A* 95: 13726-31
- Shankar GM, Leissring MA, Adame A, Sun X, Spooner E, et al. 2009. Biochemical and immunohistochemical analysis of an Alzheimer's disease mouse model reveals the presence of multiple cerebral Abeta assembly forms throughout life. *Neurobiol Dis* 36: 293-302
- Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, et al. 2008. Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nature medicine* 14: 837-42
- Shariati SA, Lau P, Hassan BA, Muller U, Dotti CG, et al. 2013. APLP2 regulates neuronal stem cell differentiation during cortical development. *J Cell Sci* 126: 1268-77
- Shen J, Bronson RT, Chen DF, Xia W, Selkoe DJ, Tonegawa S. 1997. Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell* 89: 629-39
- Shen Q, Goderie SK, Jin L, Karanth N, Sun Y, et al. 2004. Endothelial cells stimulate self-renewal and expand neurogenesis of neural stem cells. *Science* 304: 1338-40
- Shen Q, Wang Y, Kokovay E, Lin G, Chuang SM, et al. 2008. Adult SVZ stem cells lie in a vascular niche: a quantitative analysis of niche cell-cell interactions. *Cell Stem Cell* 3: 289-300
- Sheng JG, Zhou XQ, Mrak RE, Griffin WS. 1998. Progressive neuronal injury associated with amyloid plaque formation in Alzheimer disease. *J Neuropathol Exp Neurol* 57: 714-7
- Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, et al. 1995. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature* 375: 754-60
- Shetty AK, Hattiangady B, Rao MS, Shuai B. 2012. Neurogenesis response of middle-aged hippocampus to acute seizure activity. *PLoS One* 7: e43286
- Shigemoto-Mogami Y, Hoshikawa K, Goldman JE, Sekino Y, Sato K. 2014. Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal subventricular zone. *J Neurosci* 34: 2231-43
- Shimozaki K, Zhang CL, Suh H, Denli AM, Evans RM, Gage FH. 2012. SRY-box-containing gene 2 regulation of nuclear receptor tailless (Tlx) transcription in adult neural stem cells. *J Biol Chem* 287: 5969-78
- Shirotani K, Edbauer D, Capell A, Schmitz J, Steiner H, Haass C. 2003. Gamma-secretase activity is associated with a conformational change of nicastrin. *J Biol Chem* 278: 16474-7
- Shirotani K, Edbauer D, Prokop S, Haass C, Steiner H. 2004. Identification of distinct gamma-secretase complexes with different APH-1 variants. *J Biol Chem* 279: 41340-5
- Shirotani K, Tsubuki S, Iwata N, Takaki Y, Harigaya W, et al. 2001. Neprilysin degrades both amyloid beta peptides 1-40 and 1-42 most rapidly and efficiently among thiorphan- and phosphoramidon-sensitive endopeptidases. *J Biol Chem* 276: 21895-901

- Shu R, Wong W, Ma QH, Yang ZZ, Zhu H, et al. 2015. APP intracellular domain acts as a transcriptional regulator of miR-663 suppressing neuronal differentiation. *Cell Death Dis* 6
- Sierra A, Encinas JM, Deudero JJ, Chancey JH, Enikolopov G, et al. 2010. Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell* 7: 483-95
- Siman R, Card JP, Nelson RB, Davis LG. 1989. Expression of beta-amyloid precursor protein in reactive astrocytes following neuronal damage. *Neuron* 3: 275-85
- Simmons AD, Horton S, Abney AL, Johnson JE. 2001. Neurogenin2 expression in ventral and dorsal spinal neural tube progenitor cells is regulated by distinct enhancers. *Dev Biol* 229: 327-39
- Simonsen AH, McGuire J, Podust VN, Hagnelius NO, Nilsson TK, et al. 2007. A novel panel of cerebrospinal fluid biomarkers for the differential diagnosis of Alzheimer's disease versus normal aging and frontotemporal dementia. *Dement Geriatr Cogn Disord* 24: 434-40
- Singec I, Knoth R, Meyer RP, Maciarczyk J, Volk B, et al. 2006. Defining the actual sensitivity and specificity of the neurosphere assay in stem cell biology. *Nat Methods* 3: 801-6
- Sipe JD, Benson MD, Buxbaum JN, Ikeda S, Merlini G, et al. 2010. Amyloid fibril protein nomenclature: 2010 recommendations from the nomenclature committee of the International Society of Amyloidosis. *Amyloid* 17: 101-4
- Sisodia SS. 1992. Beta-amyloid precursor protein cleavage by a membrane-bound protease. *Proc Natl Acad Sci U S A* 89: 6075-9
- Sisodia SS, Koo EH, Beyreuther K, Unterbeck A, Price DL. 1990. Evidence that beta-amyloid protein in Alzheimer's disease is not derived by normal processing. *Science* 248: 492-5
- Sisodia SS, Koo EH, Hoffman PN, Perry G, Price DL. 1993. Identification and transport of full-length amyloid precursor proteins in rat peripheral nervous system. *J Neurosci* 13: 3136-42
- Small DH, Clarriss HL, Williamson TG, Reed G, Key B, et al. 1999. Neurite-outgrowth regulating functions of the amyloid protein precursor of Alzheimer's disease. *Journal of Alzheimer's disease : JAD* 1: 275-85
- Small DH, Losic D, Martin LL, Turner BJ, Friedhuber A, Aguilar MI. 2004. Alzheimer's disease therapeutics: new approaches to an ageing problem. *IUBMB Life* 56: 203-8
- Small DH, Mok SS, Bornstein JC. 2001. Alzheimer's disease and Abeta toxicity: from top to bottom. *Nat Rev Neurosci* 2: 595-8
- Small DH, Nurcombe V, Moir R, Michaelson S, Monard D, et al. 1992. Association and release of the amyloid protein precursor of Alzheimer's disease from chick brain extracellular matrix. *J Neurosci* 12: 4143-50
- Small DH, Nurcombe V, Reed G, Clarriss H, Moir R, et al. 1994. A heparin-binding domain in the amyloid protein precursor of Alzheimer's disease is involved in the regulation of neurite outgrowth. *J Neurosci* 14: 2117-27
- Small SA, Kent K, Pierce A, Leung C, Kang MS, et al. 2005. Model-guided microarray implicates the retromer complex in Alzheimer's disease. *Ann Neurol* 58: 909-19

- Smith RP, Broze GJ, Jr. 1992. Characterization of platelet-releasable forms of beta-amyloid precursor proteins: the effect of thrombin. *Blood* 80: 2252-60
- Smith RP, Higuchi DA, Broze GJ, Jr. 1990. Platelet coagulation factor XIa-inhibitor, a form of Alzheimer amyloid precursor protein. *Science* 248: 1126-8
- Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, et al. 2005. Regulation of NMDA receptor trafficking by amyloid-beta. *Nature neuroscience* 8: 1051-8
- Snyder JS, Cameron HA. 2012. Could adult hippocampal neurogenesis be relevant for human behavior? *Behav Brain Res* 227: 384-90
- Snyder JS, Soumier A, Brewer M, Pickel J, Cameron HA. 2011. Adult hippocampal neurogenesis buffers stress responses and depressive behaviour. *Nature* 476: 458-61
- Soba P, Eggert S, Wagner K, Zentgraf H, Siehl K, et al. 2005. Homo- and heterodimerization of APP family members promotes intercellular adhesion. *Embo J* 24: 3624-34
- Sommer L, Ma Q, Anderson DJ. 1996. neurogenins, a novel family of atonal-related bHLH transcription factors, are putative mammalian neuronal determination genes that reveal progenitor cell heterogeneity in the developing CNS and PNS. *Mol Cell Neurosci* 8: 221-41
- Song MR, Ghosh A. 2004. FGF2-induced chromatin remodeling regulates CNTF-mediated gene expression and astrocyte differentiation. *Nature neuroscience* 7: 229-35
- Soto C, Branes MC, Alvarez J, Inestrosa NC. 1994. Structural determinants of the Alzheimer's amyloid beta-peptide. *Journal of neurochemistry* 63: 1191-8
- Sotthibundhu A, Li QX, Thangnipon W, Coulson EJ. 2009. Abeta(1-42) stimulates adult SVZ neurogenesis through the p75 neurotrophin receptor. *Neurobiol Aging* 30: 1975-85
- Spassky N, Merkle FT, Flames N, Tramontin AD, Garcia-Verdugo JM, Alvarez-Buylla A. 2005. Adult ependymal cells are postmitotic and are derived from radial glial cells during embryogenesis. *J Neurosci* 25: 10-8
- Spencer B, Masliah E. 2014. Immunotherapy for Alzheimer's disease: past, present and future. *Front Aging Neurosci* 6: 114
- Srivastava AS, Malhotra R, Sharp J, Berggren T. 2008. Potentials of ES cell therapy in neurodegenerative diseases. *Curr Pharm Des* 14: 3873-9
- Steinbach JP, Muller U, Leist M, Li ZW, Nicotera P, Aguzzi A. 1998. Hypersensitivity to seizures in beta-amyloid precursor protein deficient mice. *Cell Death Differ* 5: 858-66
- Steiner H, Winkler E, Edbauer D, Prokop S, Basset G, et al. 2002. PEN-2 is an integral component of the gamma-secretase complex required for coordinated expression of presenilin and nicastrin. *J Biol Chem* 277: 39062-5
- Steinhoff T, Moritz E, Wollmer MA, Mohajeri MH, Kins S, Nitsch RM. 2001. Increased cystatin C in astrocytes of transgenic mice expressing the K670N-M671L mutation of the amyloid precursor protein and deposition in brain amyloid plaques. *Neurobiol Dis* 8: 647-54
- Stoka V, Turk B, Turk V. 2005. Lysosomal cysteine proteases: structural features and their role in apoptosis. *IUBMB Life*. 57: 347-53
- Stone SS, Teixeira CM, Zaslavsky K, Wheeler AL, Martinez-Canabal A, et al. 2011. Functional convergence of developmentally and adult-generated granule

- cells in dentate gyrus circuits supporting hippocampus-dependent memory. *Hippocampus* 21: 1348-62
- Struhl G, Adachi A. 1998. Nuclear access and action of notch in vivo. *Cell* 93: 649-60
- Suarez-Pereira I, Canals S, Carrion AM. 2014. Adult newborn neurons are involved in learning acquisition and long-term memory formation: The distinct demands on temporal neurogenesis of different cognitive tasks. *Hippocampus*
- Suda H, Aoyagi T, Hamada M, Takeuchi T, Umezawa H. 1972. Antipain, a new protease inhibitor isolated from actinomycetes. *J Antibiot (Tokyo)* 25: 263-6
- Suh YH, Checler F. 2002. Amyloid precursor protein, presenilins, and alpha-synuclein: molecular pathogenesis and pharmacological applications in Alzheimer's disease. *Pharmacol Rev* 54: 469-525
- Sultan S, Mandairon N, Kermen F, Garcia S, Sacquet J, Didier A. 2010. Learning-dependent neurogenesis in the olfactory bulb determines long-term olfactory memory. *Faseb J* 24: 2355-63
- Sumioka A, Nagaishi S, Yoshida T, Lin A, Miura M, Suzuki T. 2005. Role of 14-3-3gamma in FE65-dependent gene transactivation mediated by the amyloid beta-protein precursor cytoplasmic fragment. *J Biol Chem* 280: 42364-74
- Sun B, Zhou Y, Halabisky B, Lo I, Cho SH, et al. 2008. Cystatin C-cathepsin B axis regulates amyloid beta levels and associated neuronal deficits in an animal model of Alzheimer's disease. *Neuron* 60: 247-57
- Sun D. 2014. The potential of endogenous neurogenesis for brain repair and regeneration following traumatic brain injury. *Neural Regen Res* 9: 688-92
- Sun D, Bullock MR, Altememi N, Zhou Z, Hagood S, et al. 2010. The effect of epidermal growth factor in the injured brain after trauma in rats. *J Neurotrauma* 27: 923-38
- Sun Q. 1989. Growth stimulation of 3T3 fibroblasts by cystatin. *Exp Cell Res* 180: 150-60
- Sun Y, Nadal-Vicens M, Misono S, Lin MZ, Zubiaga A, et al. 2001. Neurogenin promotes neurogenesis and inhibits glial differentiation by independent mechanisms. *Cell* 104: 365-76
- Sun Y, Shi J, Fu SL, Lu PH, Xu XM. 2003. [Effects of embryonic neural stem cells and glial cell line-derived neurotrophic factor in the repair of spinal cord injury]. *Sheng Li Xue Bao* 55: 349-54
- Sundholm-Peters NL, Yang HK, Goings GE, Walker AS, Szele FG. 2005. Subventricular zone neuroblasts emigrate toward cortical lesions. *J Neuropathol Exp Neurol* 64: 1089-100
- Supeno NE, Pati S, Hadi RA, Ghani AR, Mustafa Z, et al. 2013. IGF-1 acts as controlling switch for long-term proliferation and maintenance of EGF/FGF-responsive striatal neural stem cells. *Int J Med Sci* 10: 522-31
- Surget A, Tanti A, Leonardo ED, Laugeray A, Rainer Q, et al. 2011. Antidepressants recruit new neurons to improve stress response regulation. *Mol Psychiatry* 16: 1177-88
- Suyama T, Furuya M, Nishiyama M, Kasuya Y, Kimura S, et al. 2005. Up-regulation of the interferon gamma (IFN-gamma)-inducible chemokines IFN-inducible T-cell alpha chemoattractant and monokine induced by IFN-

- gamma and of their receptor CXC receptor 3 in human renal cell carcinoma. *Cancer* 103: 258-67
- Suzuki N, Cheung TT, Cai XD, Odaka A, Otvos L, Jr., et al. 1994. An increased percentage of long amyloid beta protein secreted by familial amyloid beta protein precursor (beta APP717) mutants. *Science* 264: 1336-40
- Suzuki Y, Jin C, Yazawa I. 2014a. Cystatin C triggers neuronal degeneration in a model of multiple system atrophy. *Am J Pathol* 184: 790-9
- Suzuki Y, Jin C, Yazawa I. 2014b. Cystatin C triggers neuronal degeneration in a model of multiple system atrophy. *Am J Pathol*. 184: 790-9
- Tabaton M, Cammarata S, Mandybur T, Richey P, Kawai M, et al. 1992. Senile plaques in cerebral amyloid angiopathy show accumulation of amyloid precursor protein without cytoskeletal abnormalities. *Brain Res* 593: 299-303
- Tagliavini F, Giaccone G, Linoli G, Frangione B, Bugiani O. 1989. Cerebral extracellular preamyloid deposits in Alzheimer's disease, Down syndrome and nondemented elderly individuals. *Prog Clin Biol Res* 317: 1001-5
- Takahashi K, Rochford CD, Neumann H. 2005. Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *J Exp Med* 201: 647-57
- Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, et al. 2007. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131: 861-72
- Takasugi N, Tomita T, Hayashi I, Tsuruoka M, Niimura M, et al. 2003. The role of presenilin cofactors in the gamma-secretase complex. *Nature* 422: 438-41
- Takeda A, Loveman E, Clegg A, Kirby J, Picot J, et al. 2006. A systematic review of the clinical effectiveness of donepezil, rivastigmine and galantamine on cognition, quality of life and adverse events in Alzheimer's disease. *Int J Geriatr Psychiatry* 21: 17-28
- Tamai M, Hanada K, Adachi T, Oguma K, Kashiwagi K, et al. 1981. Papain inhibitions by optically active E-64 analogs. *J Biochem* 90: 255-7
- Tarassishin L, Yin YI, Bassit B, Li YM. 2004. Processing of Notch and amyloid precursor protein by gamma-secretase is spatially distinct. *Proc Natl Acad Sci U S A* 101: 17050-5
- Tarawneh R, Holtzman DM. 2012. The clinical problem of symptomatic Alzheimer disease and mild cognitive impairment. *Cold Spring Harb Perspect Med* 2: a006148
- Tarr PE, Roncarati R, Pelicci G, Pelicci PG, D'Adamio L. 2002. Tyrosine phosphorylation of the beta-amyloid precursor protein cytoplasmic tail promotes interaction with Shc. *J Biol Chem* 277: 16798-804
- Taupin P, Ray J, Fischer WH, Suhr ST, Hakansson K, et al. 2000. FGF-2-responsive neural stem cell proliferation requires CCg, a novel autocrine/paracrine cofactor. *Neuron* 28: 385-97
- Tavazoie M, Van der Veken L, Silva-Vargas V, Louissaint M, Colonna L, et al. 2008. A specialized vascular niche for adult neural stem cells. *Cell Stem Cell* 3: 279-88
- Tavera C, Leung-Tack J, Prevot D, Gensac MC, Martinez J, et al. 1992. Cystatin C secretion by rat glomerular mesangial cells: autocrine loop for in vitro growth-promoting activity. *Biochem Biophys Res Commun* 182: 1082-8

- Taylor CJ, Jhaveri DJ, Bartlett PF. 2013. The therapeutic potential of endogenous hippocampal stem cells for the treatment of neurological disorders. *Front Cell Neurosci* 7: 5
- Temple S. 1989. Division and differentiation of isolated CNS blast cells in microculture. *Nature* 340: 471-3
- Temple S, Alvarez-Buylla A. 1999. Stem cells in the adult mammalian central nervous system. *Curr Opin Neurobiol* 9: 135-41
- Teramoto T, Qiu J, Plumier JC, Moskowitz MA. 2003. EGF amplifies the replacement of parvalbumin-expressing striatal interneurons after ischemia. *The Journal of clinical investigation* 111: 1125-32
- Terauchi Y, Kadowaki T. 2002. Insights into molecular pathogenesis of type 2 diabetes from knockout mouse models. *Endocr J* 49: 247-63
- Terry R, Masliah E, Salmon D, Butters N, Deteresa R, et al. 1991a. Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol* 30: 572-80
- Terry RD. 1963. The Fine Structure of Neurofibrillary Tangles in Alzheimer's Disease. *J Neuropathol Exp Neurol* 22: 629-42
- Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, et al. 1991b. Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol* 4: 572-80
- Thal DR, Ghebremedhin E, Orantes M, Wiestler OD. 2003. Vascular pathology in Alzheimer disease: correlation of cerebral amyloid angiopathy and arteriosclerosis/lipohyalinosis with cognitive decline. *J Neuropathol Exp Neurol* 62: 1287-301
- Tham M, Ramasamy S, Gan HT, Ramachandran A, Poonepalli A, et al. 2010. CSPG is a secreted factor that stimulates neural stem cell survival possibly by enhanced EGFR signaling. *PLoS One* 5: e15341
- Thinakaran G, Borchelt DR, Lee MK, Slunt HH, Spitzer L, et al. 1996. Endoproteolysis of presenilin 1 and accumulation of processed derivatives in vivo. *Neuron* 17: 181-90
- Thinakaran G, Kitt CA, Roskams AJ, Slunt HH, Masliah E, et al. 1995. Distribution of an APP homolog, APLP2, in the mouse olfactory system: a potential role for APLP2 in axogenesis. *J Neurosci* 15: 6314-26
- Thoma EC, Wischmeyer E, Offen N, Maurus K, Siren AL, et al. 2012. Ectopic expression of neurogenin 2 alone is sufficient to induce differentiation of embryonic stem cells into mature neurons. *PLoS One* 7: e38651
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, et al. 1998. Embryonic stem cell lines derived from human blastocysts. *Science* 282: 1145-7
- Thornberry NA. 1997. The caspase family of cysteine proteases. *Br Med Bull* 53: 478-90
- Tizon B, Ribe EM, Mi W, Troy CM, Levy E. 2010. Cystatin C protects neuronal cells from amyloid-beta-induced toxicity. *Journal of Alzheimer's disease : JAD* 19: 885-94
- Tokutake T, Kasuga K, Yajima R, Sekine Y, Tezuka T, et al. 2012. Hyperphosphorylation of Tau induced by naturally secreted amyloid- β at nanomolar concentrations is modulated by insulin-dependent Akt-GSK3 β signaling pathway. *J Biol Chem* 287: 35222-33

- Tomita S, Kirino Y, Suzuki T. 1998. Cleavage of Alzheimer's amyloid precursor protein (APP) by secretases occurs after O-glycosylation of APP in the protein secretory pathway. Identification of intracellular compartments in which APP cleavage occurs without using toxic agents that interfere with protein metabolism. *J Biol Chem* 273: 6277-84
- Toni N, Laplagne DA, Zhao C, Lombardi G, Ribak CE, et al. 2008. Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nature neuroscience* 11: 901-7
- Toni N, Teng EM, Bushong EA, Aimone JB, Zhao C, et al. 2007. Synapse formation on neurons born in the adult hippocampus. *Nature neuroscience* 10: 727-34
- Tozuka Y, Fukuda S, Namba T, Seki T, Hisatsune T. 2005. GABAergic excitation promotes neuronal differentiation in adult hippocampal progenitor cells. *Neuron* 47: 803-15
- Tran PB, Ren D, Veldhouse TJ, Miller RJ. 2004. Chemokine receptors are expressed widely by embryonic and adult neural progenitor cells. *Journal of neuroscience research* 76: 20-34
- Trapp BD, Hauer PE. 1994. Amyloid precursor protein is enriched in radial glia: implications for neuronal development. *Journal of neuroscience research* 37: 538-50
- Tropepe V, Craig CG, Morshead CM, van der Kooy D. 1997. Transforming growth factor-alpha null and senescent mice show decreased neural progenitor cell proliferation in the forebrain subependyma. *J Neurosci* 17: 7850-9
- Tropepe V, Hitoshi S, Sirard C, Mak TW, Rossant J, van der Kooy D. 2001. Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. *Neuron* 30: 65-78
- Tropepe V, Sibilio M, Ciruna BG, Rossant J, Wagner EF, van der Kooy D. 1999. Distinct neural stem cells proliferate in response to EGF and FGF in the developing mouse telencephalon. *Dev Biol* 208: 166-88
- Tsai J, Grutzendler J, Duff K, Gan WB. 2004. Fibrillar amyloid deposition leads to local synaptic abnormalities and breakage of neuronal branches. *Nature neuroscience* 7: 1181-3
- Tsuji O, Miura K, Okada Y, Fujiyoshi K, Mukaino M, et al. 2010. Therapeutic potential of appropriately evaluated safe-induced pluripotent stem cells for spinal cord injury. *Proc Natl Acad Sci U S A* 107: 12704-9
- Tsuji-Akimoto S, Yabe I, Niino M, Kikuchi S, Sasaki H. 2009. Cystatin C in cerebrospinal fluid as a biomarker of ALS. *Neurosci Lett* 452: 52-5
- Turbic A, Leong SY, Turnley AM. 2011. Chemokines and inflammatory mediators interact to regulate adult murine neural precursor cell proliferation, survival and differentiation. *PLoS One* 6: e25406
- Tureyen K, Vemuganti R, Bowen KK, Sailor KA, Dempsey RJ. 2005. EGF and FGF-2 infusion increases post-ischemic neural progenitor cell proliferation in the adult rat brain. *Neurosurgery* 57: 1254-63; discussion 54-63
- Turk B, Turk D, Turk V. 2000. Lysosomal cysteine proteases: more than scavengers. *Biochim Biophys Acta* 1477: 98-111
- Turk B, Turk V, Turk D. 1997. Structural and functional aspects of papain-like cysteine proteinases and their protein inhibitors. *Biol Chem* 378: 141-50

- Turk V, Stoka V, Turk D. 2008. Cystatins: biochemical and structural properties, and medical relevance. *Front Biosci* 13: 5406-20
- Turnley AM, Bartlett PF. 2000. Cytokines that signal through the leukemia inhibitory factor receptor-beta complex in the nervous system. *Journal of neurochemistry* 74: 889-99
- Tuszynski MH, Thal L, Pay M, Salmon DP, U HS, et al. 2005. A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer disease. *Nature medicine* 11: 551-5
- Tyan SH, Shih AY, Walsh JJ, Maruyama H, Sarsoza F, et al. 2012. Amyloid precursor protein (APP) regulates synaptic structure and function. *Mol Cell Neurosci* 51: 43-52
- Tycko R, Wickner RB. 2013. Molecular structures of amyloid and prion fibrils: consensus versus controversy. *Acc Chem Res* 46: 1487-96
- Uchida N, Buck DW, He D, Reitsma MJ, Masek M, et al. 2000. Direct isolation of human central nervous system stem cells. *Proc Natl Acad Sci U S A* 97: 14720-5
- Ulrich JD, Finn MB, Wang Y, Shen A, Mahan TE, et al. 2014. Altered microglial response to Aβ plaques in APPS1-21 mice heterozygous for TREM2. *Mol Neurodegener* 9: 20
- Urbanc B, Cruz L, Le R, Sanders J, Ashe KH, et al. 2002. Neurotoxic effects of thioflavin S-positive amyloid deposits in transgenic mice and Alzheimer's disease. *Proc Natl Acad Sci U S A* 99: 13990-5
- Urbizu A, Canet-Pons J, Munoz-Marmol AM, Aldecoa I, Lopez MT, et al. 2014. Cystatin C is differentially involved in multiple system atrophy phenotypes. *Neuropathol Appl Neurobiol*: DOI: 10.1111/nan.12134
- Van Nostrand WE, Schmaier AH, Farrow JS, Cines DB, Cunningham DD. 1991a. Protease nexin-2/amyloid beta-protein precursor in blood is a platelet-specific protein. *Biochem Biophys Res Commun* 175: 15-21
- Van Nostrand WE, Schmaier AH, Farrow JS, Cunningham DD. 1991b. Platelet protease nexin-2/amyloid beta-protein precursor. Possible pathologic and physiologic functions. *Ann N Y Acad Sci* 640: 140-4
- Van Nostrand WE, Wagner SL, Farrow JS, Cunningham DD. 1990. Immunopurification and protease inhibitory properties of protease nexin-2/amyloid beta-protein precursor. *J Biol Chem* 265: 9591-4
- Vandenberghe R, Tournoy J. 2005. Cognitive aging and Alzheimer's disease. *Postgrad Med J* 81: 343-52
- Vasiljeva O, Dolinar M, Pungercar JR, Turk V, Turk B. 2005. Recombinant human procathepsin S is capable of autocatalytic processing at neutral pH in the presence of glycosaminoglycans. *FEBS Lett*. 579: 1285-90
- Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, et al. 1999. Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 286: 735-41
- Vassar R, Kovacs DM, Yan R, Wong PC. 2009. The beta-secretase enzyme BACE in health and Alzheimer's disease: regulation, cell biology, function, and therapeutic potential. *J Neurosci* 29: 12787-94
- Veeraraghavalu K, Choi SH, Sisodia SS. 2010. Expression of familial Alzheimer's disease-linked human presenilin 1 variants impair enrichment-induced adult hippocampal neurogenesis. *Neurodegener Dis* 7: 46-9

- Veeraraghavalu K, Choi SH, Zhang X, Sisodia SS. 2013. Endogenous expression of FAD-linked PS1 impairs proliferation, neuronal differentiation and survival of adult hippocampal progenitors. *Mol Neurodegener* 8: 41
- Vehmas AK, Kawas CH, Stewart WF, Troncoso JC. 2003. Immune reactive cells in senile plaques and cognitive decline in Alzheimer's disease. *Neurobiol Aging* 24: 321-31
- Veillard F, Saidi A, Burden RE, Scott CJ, Gillet L, et al. 2011. Cysteine cathepsins S and L modulate anti-angiogenic activities of human endostatin. *J Biol Chem* 286: 37158-67
- Venkataramani V, Rossner C, Iffland L, Schweyer S, Tamboli IY, et al. 2010. Histone deacetylase inhibitor valproic acid inhibits cancer cell proliferation via down-regulation of the alzheimer amyloid precursor protein. *J Biol Chem* 285: 10678-89
- Vergheze PB, Castellano JM, Garai K, Wang Y, Jiang H, et al. 2013. ApoE influences amyloid-beta (Abeta) clearance despite minimal apoE/Abeta association in physiological conditions. *Proc Natl Acad Sci U S A* 110: E1807-16
- Vescovi AL, Reynolds BA, Fraser DD, Weiss S. 1993. bFGF regulates the proliferative fate of unipotent (neuronal) and bipotent (neuronal/astroglial) EGF-generated CNS progenitor cells. *Neuron* 11: 951-66
- Veyrac A, Didier A, Colpaert F, Jourdan F, Marien M. 2005. Activation of noradrenergic transmission by alpha2-adrenoceptor antagonists counteracts deafferentation-induced neuronal death and cell proliferation in the adult mouse olfactory bulb. *Exp Neurol* 194: 444-56
- Vivar C, Potter MC, Choi J, Lee JY, Stringer TP, et al. 2012. Monosynaptic inputs to new neurons in the dentate gyrus. *Nat Commun* 3: 1107
- Vogt DL, Thomas D, Galvan V, Bredesen DE, Lamb BT, Pimplikar SW. 2011. Abnormal neuronal networks and seizure susceptibility in mice overexpressing the APP intracellular domain. *Neurobiol Aging* 32: 1725-9
- Volicer L, Harper DG, Manning BC, Goldstein R, Satlin A. 2001. Sundowning and circadian rhythms in Alzheimer's disease. *Am J Psychiatry* 158: 704-11
- von Koch CS, Zheng H, Chen H, Trumbauer M, Thinakaran G, et al. 1997. Generation of APLP2 KO mice and early postnatal lethality in APLP2/APP double KO mice. *Neurobiol Aging* 18: 661-9
- von Rotz RC, Kohli BM, Bosset J, Meier M, Suzuki T, et al. 2004. The APP intracellular domain forms nuclear multiprotein complexes and regulates the transcription of its own precursor. *J Cell Sci* 117: 4435-48
- Vrotsos EG, Kolattukudy PE, Sugaya K. 2009. MCP-1 involvement in glial differentiation of neuroprogenitor cells through APP signaling. *Brain Res Bull* 79: 97-103
- Waclaw RR, Allen ZJ, 2nd, Bell SM, Erdelyi F, Szabo G, et al. 2006. The zinc finger transcription factor Sp8 regulates the generation and diversity of olfactory bulb interneurons. *Neuron* 49: 503-16
- Wagner JP, Black IB, DiCicco-Bloom E. 1999. Stimulation of neonatal and adult brain neurogenesis by subcutaneous injection of basic fibroblast growth factor. *J Neurosci* 19: 6006-16
- Wake H, Moorhouse AJ, Nabekura J. 2011. Functions of microglia in the central nervous system--beyond the immune response. *Neuron Glia Biol* 7: 47-53

- Wang B, Wang Z, Sun L, Yang L, Li H, et al. 2014. The amyloid precursor protein controls adult hippocampal neurogenesis through GABAergic interneurons. *J Neurosci* 34: 13314-25
- Wang C, Sun B, Zhou Y, Grubb A, Gan L. 2012. Cathepsin B degrades amyloid-beta in mice expressing wild-type human amyloid precursor protein. *J Biol Chem* 287: 39834-41
- Wang JW, David DJ, Monckton JE, Battaglia F, Hen R. 2008. Chronic fluoxetine stimulates maturation and synaptic plasticity of adult-born hippocampal granule cells. *J Neurosci* 28: 1374-84
- Wang Y, Zhang M, Moon C, Hu Q, Wang B, et al. 2009. The APP-interacting protein FE65 is required for hippocampus-dependent learning and long-term potentiation. *Learn Mem* 16: 537-44
- Wang ZZ, Jensson OT, L., Vinters HV. 1997. Microvascular degeneration in hereditary cystatin C amyloid angiopathy of the brain. *APMIS* 105: 41-7
- Warfel AH, Zucker-Franklin D, Frangione B, Ghiso J. 1987. Constitutive secretion of cystatin C (gamma-trace) by monocytes and macrophages and its downregulation after stimulation. *J Exp Med* 166: 1912-7
- Wasco W, Bupp K, Magendantz M, Gusella JF, Tanzi RE, Solomon F. 1992. Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer disease-associated amyloid beta protein precursor. *Proc Natl Acad Sci U S A* 89: 10758-62
- Wasco W, Gurubhagavatula S, Paradis MD, Romano DM, Sisodia SS, et al. 1993. Isolation and characterization of APLP2 encoding a homologue of the Alzheimer's associated amyloid beta protein precursor. *Nat Genet* 5: 95-100
- Watanabe S, Hayakawa T, Wakasugi K, Yamanaka K. 2014. Cystatin C protects neuronal cells against mutant copper-zinc superoxide dismutase-mediated toxicity. *Cell Death Dis* 5: e1497
- Watkinson A. 1999. Stratum corneum thiol protease (SCTP): a novel cysteine protease of late epidermal differentiation. *Arch Dermatol Res* 291: 260-8
- Webster CI, Burrell M, Olsson LL, Fowler SB, Digby S, et al. 2014. Engineering neprilysin activity and specificity to create a novel therapeutic for Alzheimer's disease. *PLoS One* 9: e104001
- Webster MT, Groome N, Francis PT, Pearce BR, Sherriff FE, et al. 1995. A novel protein, amyloid precursor-like protein 2, is present in human brain, cerebrospinal fluid and conditioned media. *Biochem J* 310 (Pt 1): 95-9
- Weggen S, Eriksen JL, Das P, Sagi SA, Wang R, et al. 2001. A subset of NSAIDs lower amyloidogenic Abeta42 independently of cyclooxygenase activity. *Nature* 414: 212-6
- Weidemann A, Eggert S, Reinhard FB, Vogel M, Paliga K, et al. 2002. A novel epsilon-cleavage within the transmembrane domain of the Alzheimer amyloid precursor protein demonstrates homology with Notch processing. *Biochemistry* 41: 2825-35
- Weidemann A, Konig G, Bunke D, Fischer P, Salbaum JM, et al. 1989. Identification, biogenesis, and localization of precursors of Alzheimer's disease A4 amyloid protein. *Cell* 57: 115-26
- Weiner MF, Hynan LS, Bret ME, White C, 3rd. 2005. Early behavioral symptoms and course of Alzheimer's disease. *Acta Psychiatr Scand* 111: 367-71

- Weisgraber KH, Rall SC, Jr., Mahley RW. 1981. Human E apoprotein heterogeneity. Cysteine-arginine interchanges in the amino acid sequence of the apo-E isoforms. *J Biol Chem* 256: 9077-83
- Weiss S, Reynolds BA, Vescovi AL, Morshead C, Craig CG, van der Kooy D. 1996. Is there a neural stem cell in the mammalian forebrain? *Trends Neurosci* 19: 387-93
- Weissman IL, Anderson DJ, Gage F. 2001. Stem and progenitor cells: origins, phenotypes, lineage commitments, and transdifferentiations. *Annu Rev Cell Dev Biol* 17: 387-403
- Wen PH, Friedrich VL, Jr., Shioi J, Robakis NK, Elder GA. 2002a. Presenilin-1 is expressed in neural progenitor cells in the hippocampus of adult mice. *Neurosci Lett* 318: 53-6
- Wen PH, Shao X, Shao Z, Hof PR, Wisniewski T, et al. 2002b. Overexpression of wild type but not an FAD mutant presenilin-1 promotes neurogenesis in the hippocampus of adult mice. *Neurobiol Dis* 10: 8-19
- Werner S, Unsicker K, von Bohlen und Halbach O. 2011. Fibroblast growth factor-2 deficiency causes defects in adult hippocampal neurogenesis, which are not rescued by exogenous fibroblast growth factor-2. *Journal of neuroscience research* 89: 1605-17
- Wernig M, Zhao JP, Pruszak J, Hedlund E, Fu D, et al. 2008. Neurons derived from reprogrammed fibroblasts functionally integrate into the fetal brain and improve symptoms of rats with Parkinson's disease. *Proc Natl Acad Sci U S A* 105: 5856-61
- Weskamp G, Cai H, Brodie TA, Higashiyama S, Manova K, et al. 2002. Mice lacking the metalloprotease-disintegrin MDC9 (ADAM9) have no evident major abnormalities during development or adult life. *Mol Cell Biol* 22: 1537-44
- Whitehouse PJ, Price DL, Struble RG, Clark AW, Coyle JT, Delon MR. 1982. Alzheimer's disease and senile dementia: loss of neurons in the basal forebrain. *Science* 215: 1237-9
- Whitman MC, Greer CA. 2007a. Adult-generated neurons exhibit diverse developmental fates. *Dev Neurobiol* 67: 1079-93
- Whitman MC, Greer CA. 2007b. Synaptic integration of adult-generated olfactory bulb granule cells: basal axodendritic centrifugal input precedes apical dendrodendritic local circuits. *J Neurosci* 27: 9951-61
- Whitson JS, Selkoe DJ, Cotman CW. 1989. Amyloid beta protein enhances the survival of hippocampal neurons in vitro. *Science* 243: 1488-90
- Wichterle H, Lieberam I, Porter JA, Jessell TM. 2002. Directed differentiation of embryonic stem cells into motor neurons. *Cell* 110: 385-97
- Wilcox D, Mason RW. 1992. Inhibition of cysteine proteinases in lysosomes and whole cells. *Biochem J* 285 (Pt 2): 495-502
- Wilkinson D, Schindler R, Schwam E, Waldemar G, Jones RW, et al. 2009. Effectiveness of donepezil in reducing clinical worsening in patients with mild-to-moderate alzheimer's disease. *Dement Geriatr Cogn Disord* 28: 244-51
- Willem M, Garratt AN, Novak B, Citron M, Kaufmann S, et al. 2006. Control of peripheral nerve myelination by the beta-secretase BACE1. *Science* 314: 664-6
- Williamson TG, Mok SS, Henry A, Cappai R, Lander AD, et al. 1996a. Secreted glypican binds to the amyloid precursor protein of Alzheimer's disease

- (APP) and inhibits APP-induced neurite outgrowth. *J Biol Chem* 271: 31215-21
- Williamson TG, Mok SS, Henry A, Cappai R, Lander AD, et al. 1996b. Secreted glypican binds to the amyloid precursor protein of Alzheimer's disease (APP) and inhibits APP-induced neurite outgrowth. *J Biol Chem* 271: 31215-21
- Winner B, Cooper-Kuhn CM, Aigner R, Winkler J, Kuhn HG. 2002. Long-term survival and cell death of newly generated neurons in the adult rat olfactory bulb. *Eur J Neurosci* 16: 1681-9
- Winner B, Couillard-Despres S, Geyer M, Aigner R, Bogdahn U, et al. 2008. Dopaminergic lesion enhances growth factor-induced striatal neuroblast migration. *J Neuropathol Exp Neurol* 67: 105-16
- Winner B, Kohl Z, Gage FH. 2011. Neurodegenerative disease and adult neurogenesis. *Eur J Neurosci* 33: 1139-51
- Winner B, Lie DC, Rockenstein E, Aigner R, Aigner L, et al. 2004. Human wild-type alpha-synuclein impairs neurogenesis. *J Neuropathol Exp Neurol* 63: 1155-66
- Winocur G, Wojtowicz JM, Sekeres M, Snyder JS, Wang S. 2006. Inhibition of neurogenesis interferes with hippocampus-dependent memory function. *Hippocampus* 16: 296-304
- Wisniewski T, Frangione B. 1992. Apolipoprotein E: a pathological chaperone protein in patients with cerebral and systemic amyloid. *Neurosci Lett* 135: 235-8
- Wojcik L, Sawicka A, Rivera S, Zalewska T. 2009. Neurogenesis in gerbil hippocampus following brain ischemia: focus on the involvement of metalloproteinases. *Acta Neurobiol Exp (Wars)* 69: 52-61
- Wojcik-Stanaszek L, Gregor A, Zalewska T. 2011. Regulation of neurogenesis by extracellular matrix and integrins. *Acta Neurobiol Exp (Wars)* 71: 103-12
- Wolf SA, Melnik A, Kempermann G. 2011. Physical exercise increases adult neurogenesis and telomerase activity, and improves behavioral deficits in a mouse model of schizophrenia. *Brain Behav Immun* 25: 971-80
- Wolfe MS, Xia W, Ostaszewski BL, Diehl TS, Kimberly WT, Selkoe DJ. 1999. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and gamma-secretase activity. *Nature* 398: 513-7
- Wong G, Goldshmit Y, Turnley AM. 2004. Interferon-gamma but not TNF alpha promotes neuronal differentiation and neurite outgrowth of murine adult neural stem cells. *Exp Neurol* 187: 171-7
- Wong PC, Zheng H, Chen H, Becher MW, Sirinathsinghji DJ, et al. 1997. Presenilin 1 is required for Notch1 and DII1 expression in the paraxial mesoderm. *Nature* 387: 288-92
- Wood JG, Mirra SS, Pollock NJ, Binder LI. 1986. Neurofibrillary tangles of Alzheimer disease share antigenic determinants with the axonal microtubule-associated protein tau (tau). *Proc Natl Acad Sci U S A* 11: 4040-43
- Wood SJ, Chan W, Wetzel R. 1996a. Seeding of A beta fibril formation is inhibited by all three isotypes of apolipoprotein E. *Biochemistry* 35: 12623-8
- Wood SJ, Maleeff B, Hart T, Wetzel R. 1996b. Physical, morphological and functional differences between pH 5.8 and 7.4 aggregates of the Alzheimer's amyloid peptide Abeta. *J Mol Biol* 256: 870-7

- Xia J, Wang F, Wang L, Fan Q. 2013. Elevated serine protease HtrA1 inhibits cell proliferation, reduces invasion, and induces apoptosis in esophageal squamous cell carcinoma by blocking the nuclear factor-kappaB signaling pathway. *Tumour Biol* 34: 317-28
- Xiao Q, Yan P, Ma X, Liu H, Perez R, et al. 2014. Enhancing astrocytic lysosome biogenesis facilitates Abeta clearance and attenuates amyloid plaque pathogenesis. *J Neurosci* 34: 9607-20
- Xu PT, Schmechel D, Qiu HL, Herbstreith M, Rothrock-Christian T, et al. 1999. Sialylated human apolipoprotein E (apoEs) is preferentially associated with neuron-enriched cultures from APOE transgenic mice. *Neurobiol Dis* 6: 63-75
- Xu Y, Kim HS, Joo Y, Choi Y, Chang KA, et al. 2007. Intracellular domains of amyloid precursor-like protein 2 interact with CP2 transcription factor in the nucleus and induce glycogen synthase kinase-3beta expression. *Cell Death Differ* 14: 79-91
- Yadaiah M, Sudhamalla B, Rao PN, Roy KR, Ramakrishna D, et al. 2013a. Arrested cell proliferation through cysteine protease activity of eukaryotic ribosomal protein S4. *FASEB* 27: 803-10
- Yadaiah M, Sudhamalla B, Rao PN, Roy KR, Ramakrishna D, et al. 2013b. Arrested cell proliferation through cysteine protease activity of eukaryotic ribosomal protein S4. *Faseb J* 27: 803-10
- Yakubov E, Rechavi G, Rozenblatt S, Givol D. 2010. Reprogramming of human fibroblasts to pluripotent stem cells using mRNA of four transcription factors. *Biochem Biophys Res Commun* 394: 189-93
- Yamada M, Tsukagoshi H, Wada Y, Otomo E, Hayakawa M, et al. 1989. Absence of the cystatin C amyloid in the cerebral amyloid angiopathy, senile plaque, and extra-CNS amyloid deposits of aged Japanese. *Acta neurologica Scandinavica* 79: 504-9
- Yamaguchi H, Hirai S, Morimatsu M, Shoji M, Harigaya Y. 1988. Diffuse type of senile plaques in the brains of Alzheimer-type dementia. *Acta Neuropathol* 77: 113-9
- Yamaguchi H, Hirai S, Morimatsu M, Shoji M, Nakazato Y. 1989. Diffuse type of senile plaques in the cerebellum of Alzheimer-type dementia demonstrated by beta protein immunostain. *Acta Neuropathol* 77: 314-9
- Yamaguchi M, Mori K. 2005. Critical period for sensory experience-dependent survival of newly generated granule cells in the adult mouse olfactory bulb. *Proc Natl Acad Sci U S A* 102: 9697-702
- Yamanaka S. 2008. Induction of pluripotent stem cells from mouse fibroblasts by four transcription factors. *Cell Prolif* 41 Suppl 1: 51-6
- Yamasaki TR, Blurton-Jones M, Morrisette DA, Kitazawa M, Oddo S, LaFerla FM. 2007. Neural stem cells improve memory in an inducible mouse model of neuronal loss. *J Neurosci* 27: 11925-33
- Yamashita T, Ninomiya M, Hernandez Acosta P, Garcia-Verdugo JM, Sunabori T, et al. 2006. Subventricular zone-derived neuroblasts migrate and differentiate into mature neurons in the post-stroke adult striatum. *J Neurosci* 26: 6627-36
- Yamauchi K, Tozuka M, Hidaka H, Nakabayashi T, Sugano M, Katsuyama T. 2002. Isoform-specific effect of apolipoprotein E on endocytosis of beta-amyloid in cultures of neuroblastoma cells. *Ann Clin Lab Sci* 32: 65-74

- Yamazaki T, Koo EH, Selkoe DJ. 1996. Trafficking of cell-surface amyloid beta-protein precursor. II. Endocytosis, recycling and lysosomal targeting detected by immunolocalization. *J Cell Sci* 109 (Pt 5): 999-1008
- Yamazaki T, Koo EH, Selkoe DJ. 1997. Cell surface amyloid beta-protein precursor colocalizes with beta 1 integrins at substrate contact sites in neural cells. *J Neurosci* 17: 1004-10
- Yan R, Bienkowski MJ, Shuck ME, Miao H, Tory MC, et al. 1999. Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity. *Nature* 402: 533-7
- Yan RT, Ma WX, Wang SZ. 2001a. neurogenin2 elicits the genesis of retinal neurons from cultures of nonneural cells. *Proc Natl Acad Sci U S A* 98: 15014-9
- Yan XX, Najbauer J, Woo CC, Dashtipour K, Ribak CE, Leon M. 2001b. Expression of active caspase-3 in mitotic and postmitotic cells of the rat forebrain. *J Comp Neurol* 433: 4-22
- Yang AJ, Chandswangbhuvana D, Margol L, Glabe CG. 1998. loss of endosomal/lysosomal membrane impermeability is an early event in amyloid Abeta1-42 pathogenesis. *J Neurosci Res* 52: 691-8
- Yang DS, Small DH, Seydel U, Smith JD, Hallmayer J, et al. 1999. Apolipoprotein E promotes the binding and uptake of beta-amyloid into Chinese hamster ovary cells in an isoform-specific manner. *Neuroscience* 90: 1217-26
- Yang L, Wang Z, Wang B, Justice NJ, Zheng H. 2009a. Amyloid precursor protein regulates Cav1.2 L-type calcium channel levels and function to influence GABAergic short-term plasticity. *J Neurosci* 29: 15660-8
- Yang LB, Lindholm K, Yan R, Citron M, Xia W, et al. 2003. Elevated beta-secretase expression and enzymatic activity detected in sporadic Alzheimer disease. *Nature medicine* 9: 3-4
- Yang Y, Liu S, Qin Z, Cui Y, Qin Y, Bai S. 2009b. Alteration of cystatin C levels in cerebrospinal fluid of patients with Guillain-Barre Syndrome by a proteomical approach. *Mol Biol Rep* 36: 677-82
- Yang Z. 2008. Postnatal subventricular zone progenitors give rise not only to granular and periglomerular interneurons but also to interneurons in the external plexiform layer of the rat olfactory bulb. *J Comp Neurol* 506: 347-58
- Yasuhara O, Hanai K, Ohkubo I, Sasaki M, McGeer PL, Kimura H. 1993. Expression of cystatin C in rat, monkey and human brains. *Brain Res* 628: 85-92
- Yi SH, Jo AY, Park CH, Koh HC, Park RH, et al. 2008. Mash1 and neurogenin 2 enhance survival and differentiation of neural precursor cells after transplantation to rat brains via distinct modes of action. *Mol Ther* 16: 1873-82
- Ying QL, Nichols J, Evans EP, Smith AG. 2002. Changing potency by spontaneous fusion. *Nature* 416: 545-8
- Ylostalo JH, Bartosh TJ, Coble K, Prockop DJ. 2012. Human mesenchymal stem/stromal cells cultured as spheroids are self-activated to produce prostaglandin E2 that directs stimulated macrophages into an anti-inflammatory phenotype. *stem cells* 30: 2283-96
- Yoneyama M, Shiba T, Yamaguchi T, Ogita K. 2014. Possible involvement of caspases in proliferation of neocortical neural stem/progenitor cells in the developing mouse brain. *Biol Pharm Bull* 37: 1699-703

- Yoshikai S, Sasaki H, Doh-ura K, Furuya H, Sakaki Y. 1990. Genomic organization of the human amyloid beta-protein precursor gene. *Gene* 87: 257-63
- Yoshimura S, Takagi Y, Harada J, Teramoto T, Thomas SS, et al. 2001. FGF-2 regulation of neurogenesis in adult hippocampus after brain injury. *Proc Natl Acad Sci U S A* 98: 5874-9
- Young KM, Psachoulia K, Tripathi RB, Dunn SJ, Cossell L, et al. 2013. Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. *Neuron* 77: 873-85
- Young SZ, Platel JC, Nielsen JV, Jensen NA, Bordey A. 2010. GABA(A) Increases Calcium in Subventricular Zone Astrocyte-Like Cells Through L- and T-Type Voltage-Gated Calcium Channels. *Front Cell Neurosci* 4: 8
- Young-Pearse TL, Bai J, Chang R, Zheng JB, LoTurco JJ, Selkoe DJ. 2007. A critical function for beta-amyloid precursor protein in neuronal migration revealed by in utero RNA interference. *J Neurosci* 27: 14459-69
- Young-Pearse TL, Chen AC, Chang R, Marquez C, Selkoe DJ. 2008. Secreted APP regulates the function of full-length APP in neurite outgrowth through interaction with integrin beta1. *Neural Dev* 3: 15
- Yu DX, Marchetto MC, Gage FH. 2013. Therapeutic translation of iPSCs for treating neurological disease. *Cell Stem Cell* 12: 678-88
- Yu G, Nishimura M, Arawaka S, Levitan D, Zhang L, et al. 2000. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and betaAPP processing. *Nature* 407: 48-54
- Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, et al. 2007. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318: 1917-20
- Yu SW, Baek SH, Brennan RT, Bradley CJ, Park SK, et al. 2008. Autophagic death of adult hippocampal neural stem cells following insulin withdrawal. *stem cells* 26: 2602-10
- Yu Y, He J, Zhang Y, Luo H, Zhu S, et al. 2009. Increased hippocampal neurogenesis in the progressive stage of Alzheimer's disease phenotype in an APP/PS1 double transgenic mouse model. *Hippocampus* 19: 1247-53
- Zagorski MG, Barrow CJ. 1992. NMR studies of amyloid beta-peptides: proton assignments, secondary structure, and mechanism of an alpha-helix----beta-sheet conversion for a homologous, 28-residue, N-terminal fragment. *Biochemistry* 31: 5621-31
- Zannis VI, Breslow JL, Utermann G, Mahley RW, Weisgraber KH, et al. 1982. Proposed nomenclature of apoE isoproteins, apoE genotypes, and phenotypes. *J Lipid Res* 23: 911-4
- Zhang C, McNeil E, Dressler L, Siman R. 2007a. Long-lasting impairment in hippocampal neurogenesis associated with amyloid deposition in a knock-in mouse model of familial Alzheimer's disease. *Exp Neurol* 204: 77-87
- Zhang X, Zhang L, Cheng X, Guo Y, Sun X, et al. 2014. IGF-1 Promotes Brn-4 Expression and Neuronal Differentiation of Neural Stem Cells via the PI3K/Akt Pathway. *PLoS One* 9: e113801
- Zhang YW, Wang R, Liu Q, Zhang H, Liao FF, Xu H. 2007b. Presenilin/gamma-secretase-dependent processing of beta-amyloid precursor protein regulates EGF receptor expression. *Proc Natl Acad Sci U S A* 104: 10613-8

- Zhao C, Teng EM, Summers RG, Jr., Ming GL, Gage FH. 2006. Distinct morphological stages of dentate granule neuron maturation in the adult mouse hippocampus. *J Neurosci* 26: 3-11
- Zhao T, Zhang ZN, Rong Z, Xu Y. 2011. Immunogenicity of induced pluripotent stem cells. *Nature* 474: 212-5
- Zhao W, Dumanis SB, Tamboli IY, Rodriguez GA, Jo Ladu M, et al. 2014. Human APOE genotype affects intraneuronal Abeta1-42 accumulation in a lentiviral gene transfer model. *Hum Mol Genet* 23: 1365-75
- Zheng H, Jiang M, Trumbauer ME, Hopkins R, Sirinathsinghji DJ, et al. 1996. Mice deficient for the amyloid precursor protein gene. *Ann N Y Acad Sci* 777: 421-6
- Zheng H, Jiang M, Trumbauer ME, Sirinathsinghji DJ, Hopkins R, et al. 1995. beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. *Cell* 81: 525-31
- Zheng H, Koo EH. 2006. The amyloid precursor protein: beyond amyloid. *Mol Neurodegener* 1: 5
- Zheng H, Koo EH. 2011. Biology and pathophysiology of the amyloid precursor protein. *Mol Neurodegener* 6: 27
- Zheng M, Liu J, Ruan Z, Tian S, Ma Y, et al. 2013. Intrahippocampal injection of Abeta1-42 inhibits neurogenesis and down-regulates IFN-gamma and NF-kappaB expression in hippocampus of adult mouse brain. *Amyloid* 20: 13-20
- Zhong XM, Hou L, Luo XN, Shi HS, Hu GY, et al. 2013. Alterations of CSF cystatin C levels and their correlations with CSF Alphas40 and Alphas42 levels in patients with Alzheimer's disease, dementia with lewy bodies and the atrophic form of general paresis. *PLoS One* 8: e55328
- Zhou D, Noviello C, D'Ambrosio C, Scaloni A, D'Adamio L. 2004. Growth factor receptor-bound protein 2 interaction with the tyrosine-phosphorylated tail of amyloid beta precursor protein is mediated by its Src homology 2 domain. *J Biol Chem* 279: 25374-80
- Zhou F, Gong K, Song B, Ma T, van Laar T, et al. 2012. The APP intracellular domain (AICD) inhibits Wnt signalling and promotes neurite outgrowth. *Biochim Biophys Acta* 1823: 1233-41
- Zhou Y, Yuan J, Zhou B, Lee AJ, Ghawji M, Jr., Yoo TJ. 2011a. The therapeutic efficacy of human adipose tissue-derived mesenchymal stem cells on experimental autoimmune hearing loss in mice. *Immunology* 133: 133-40
- Zhou ZD, Chan CH, Ma QH, Xu XH, Xiao ZC, Tan EK. 2011b. The roles of amyloid precursor protein (APP) in neurogenesis: Implications to pathogenesis and therapy of Alzheimer disease. *Cell Adh Migr* 5: 280-92
- Zhu SQ, Kum W, Ho SK, Young JD, Cockram CS. 1990. Structure-function relationships of insulin receptor interactions in cultured mouse astrocytes. *Brain Res* 529: 329-32
- Zhu W, Cheng S, Xu G, Ma M, Zhou Z, et al. 2011. Intranasal nerve growth factor enhances striatal neurogenesis in adult rats with focal cerebral ischemia. *Drug Deliv* 18: 338-43
- Zhu Y, Nwabuisi-Heath E, Dumanis SB, Tai LM, Yu C, et al. 2012. APOE genotype alters glial activation and loss of synaptic markers in mice. *Glia* 60: 559-69

Ziegler AN, Chidambaram S, Forbes BE, Wood TL, Levison SW. 2014. Insulin-like growth factor-II (IGF-II) and IGF-II analogs with enhanced insulin receptor- α binding affinity promote neural stem cell expansion. *J Biol Chem* 289: 4626-33